

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁷ : C12Q 1/68, C12N 15/63, 15/85		A2	(11) International Publication Number: WO 00/08209
			(43) International Publication Date: 17 February 2000 (17.02.00)
(21) International Application Number: PCT/IB99/01444			(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 6 August 1999 (06.08.99)			
(30) Priority Data: 60/095,653 7 August 1998 (07.08.98) US			
(71) Applicant (for all designated States except US): GENSET [FR/FR]; 24, rue Royale, F-75008 Paris (FR).			
(72) Inventors; and (75) Inventors/Applicants (for US only): BLUMENFELD, Marta [FR/FR]; 5, rue Tagore, F-75013 Paris (FR). BOUGUEL-ERET, Lydie [FR/FR]; 14, rue Vouillé, F-75015 Paris (FR). CHUMAKOV, Ilya [FR/FR]; 196, rue des Chèvrefeuilles, F-77000 Vaux-le-Pénil (FR).			
(74) Agents: MARTIN, Jean-Jacques et al.; Cabinet Regimbeau, 26, avenue Kléber, F-75116 Paris (FR).			Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: NUCLEIC ACIDS ENCODING HUMAN TBC-1 PROTEIN AND POLYMORPHIC MARKERS THEREOF			
(57) Abstract <p>The invention concerns genomic and cDNA sequences of the human <i>TBC-1</i> Gene. The invention also concerns polypeptides encoded by the <i>TBC-1</i> gene. The invention also deals with antibodies directed specifically against such polypeptides that are useful as diagnostic reagents. The invention further encompasses biallelic markers of the <i>TBC-1</i> gene useful in genetic analysis.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Nucleic acids encoding human TBC-1 protein and polymorphic markers thereof.

FIELD OF THE INVENTION

The invention concerns genomic and cDNA sequences of the human *TBC-1* gene. The invention also concerns polypeptides encoded by the *TBC-1* gene. The invention also deals with antibodies directed specifically against such polypeptides that are useful as diagnostic reagents. The invention further encompasses biallelic markers of the *TBC-1* gene useful in genetic analysis.

BACKGROUND OF THE INVENTION

The incidence of prostate cancer has dramatically increased over the last decades. It averages 30-50/100,000 males in Western European countries as well as within the US White male population. In these countries, it has recently become the most commonly diagnosed malignancy, being one of every four cancers diagnosed in American males. Prostate cancer's incidence is very much population specific, since it varies from 2/100,000 in China, to over 80/100,000 among African-American males.

In France, the incidence of prostate cancer is 35/100,000 males and it is increasing by 10/100,000 per decade. Mortality due to prostate cancer is also growing accordingly. It is the second cause of cancer death among French males, and the first one among French males aged over 70. This makes prostate cancer a serious burden in terms of public health.

Prostate cancer is a latent disease. Many men carry prostate cancer cells without overt signs of disease. Autopsies of individuals dying of other causes show prostate cancer cells in 30 % of men at age 50 and in 60 % of men at age 80. Furthermore, prostate cancer can take up to 10 years to kill a patient after the initial diagnosis.

The progression of the disease usually goes from a well-defined mass within the prostate to a breakdown and invasion of the lateral margins of the prostate, followed by metastasis to regional lymph nodes, and metastasis to the bone marrow. Cancer metastasis to bone is common and often associated with uncontrollable pain.

Unfortunately, in 80 % of cases, diagnosis of prostate cancer is established when the disease has already metastasized to the bones. Of special interest is the observation that prostate cancers frequently grow more rapidly in sites of metastasis than within the prostate itself.

Early-stage diagnosis of prostate cancer mainly relies today on Prostate Specific Antigen (PSA) dosage, and allows the detection of prostate cancer seven years before clinical symptoms become apparent. The effectiveness of PSA dosage diagnosis is however limited, due to its inability to discriminate between malignant and non-malignant affections of the organ and because not all prostate cancers give rise to an elevated serum PSA concentration. Furthermore, PSA dosage and

other currently available approaches such as physical examination, tissue biopsy and bone scans are of limited value in predicting disease progression.

Therefore, there is a strong need for a reliable diagnostic procedure which would enable a more systematic early-stage prostate cancer prognosis.

5 Although an early-stage prostate cancer prognosis is important, the possibility of measuring the period of time during which treatment can be deferred is also interesting as currently available medicaments are expensive and generate important adverse effects. However, the aggressiveness of prostate tumors varies widely. Some tumors are relatively aggressive, doubling every six months whereas others are slow-growing, doubling once every five years. In fact, the majority of prostate
10 cancers grows relatively slowly and never becomes clinically manifest. Very often, affected patients are among the elderly and die from another disease before prostate cancer actually develops. Thus, a significant question in treating prostate carcinoma is how to discriminate between tumors that will progress and those that will not progress during the expected lifetime of the patient.

Hence, there is also a strong need for detection means which may be used to evaluate the
15 aggressiveness or the development potential of prostate cancer tumors once diagnosed.

Furthermore, at the present time, there is no means to predict prostate cancer susceptibility. It would also be very beneficial to detect individual susceptibility to prostate cancer. This could allow preventive treatment and a careful follow up of the development of the tumor.

A further consequence of the slow growth rate of prostate cancer is that few cancer cells are
20 actively dividing at any one time, rendering prostate cancer generally resistant to radiation and chemotherapy. Surgery is the mainstay of treatment but it is largely ineffective and removes the ejaculatory ducts, resulting in impotence. Oral oestrogens and luteinizing releasing hormone analogs are also used for treatment of prostate cancer. These hormonal treatments provide marked improvement for many patients, but they only provide temporary relief. Indeed, most of these
25 cancers soon relapse with the development of hormone-resistant tumor cells and the oestrogen treatment can lead to serious cardiovascular complications. Consequently, there is a strong need for preventive and curative treatment of prostate cancer.

Efficacy/tolerance prognosis could be precious in prostate cancer therapy. Indeed, hormonal therapy, the main treatment currently available, presents important side effects. The use of
30 chemotherapy is limited because of the small number of patients with chemosensitive tumors. Furthermore the age profile of the prostate cancer patient and intolerance to chemotherapy make the systematic use of this treatment very difficult.

Therefore, a valuable assessment of the eventual efficacy of a medicament to be administered to a prostate cancer patient as well as the patient's eventual tolerance to it may permit to
35 enhance the benefit/risk ratio of prostate cancer treatment.

It is known today that there is a familial risk of prostate cancer. Clinical studies in the 1950s had already demonstrated a familial aggregation in prostate cancer. Control-case clinical studies

have been conducted more recently to attempt to evaluate the incidence of the genetic risk factors in the disease. Thus Steinberg et al., 1990, and McWhorter et al., 1992 confirm that the risk of prostate cancer is increased in subjects having one or more relatives already affected by the disease and when forms of early diagnosis in the relatives exist.

5 It is now well established that cancer is a disease caused by the deregulation of the expression of certain genes. In fact, the development of a tumor necessitates an important succession of steps. Each of these steps comprises the deregulation of an important gene intervening in the normal metabolism of the cell and the emergence of an abnormal cellular sub-clone which overwhelms the other cell types because of a proliferative advantage. The genetic origin of this
10 concept has found confirmation in the isolation and the characterization of genes which could be responsible. These genes, commonly called "cancer genes", have an important role in the normal metabolism of the cell and are capable of intervening in carcinogenesis following a change.

Recent studies have identified three groups of genes which are frequently mutated in cancer. The first group of genes, called oncogenes, are genes whose products activate cell
15 proliferation. The normal non-mutant versions are called protooncogenes. The mutated forms are excessively or inappropriately active in promoting cell proliferation, and act in the cell in a dominant way in that a single mutant allele is enough to affect the cell phenotype. Activated oncogenes are rarely transmitted as germline mutations since they may probably be lethal when expressed in all the cells. Therefore oncogenes can only be investigated in tumor tissues.

20 The second group of genes which are frequently mutated in cancer, called tumor suppressor genes, are genes whose products inhibit cell growth. Mutant versions in cancer cells have lost their normal function, and act in the cell in a recessive way in that both copies of the gene must be inactivated in order to change the cell phenotype. Most importantly, the tumor phenotype can be rescued by the wild type allele, as shown by cell fusion experiments first described by Harris and
25 colleagues (1969). Germline mutations of tumor suppressor genes may be transmitted and thus studied in both constitutional and tumor DNA from familial or sporadic cases. The current family of tumor suppressors includes DNA-binding transcription factors (i.e., p53, WT1), transcription regulators (i.e., RB, APC, probably BRCA1), protein kinase inhibitors (i.e., p16), among others (for review, see Haber D & Harlow E, 1997).

30 The third group of genes which are frequently mutated in cancer, called mutator genes, are responsible for maintaining genome integrity and/or low mutation rates. Loss of function of both alleles increases cell mutation rates, and as a consequence, proto-oncogenes and tumor suppressor genes may be mutated. Mutator genes can also be classified as tumor suppressor genes, except for the fact that tumorigenesis caused by this class of genes cannot be suppressed simply by restoration
35 of a wild-type allele, as described above. Genes whose inactivation may lead to a mutator phenotype include mismatch repair genes (i.e., MLH1, MSH2), DNA helicases (i.e., BLM, WRN)

or other genes involved in DNA repair and genomic stability (i.e., p53, possibly BRCA1 and BRCA2) (For review see Haber D & Harlow E, 1997; Fishel R & Wilson T. 1997; Ellis NA, 1997).

There is growing evidence that a critical event in the progression of a tumor cell from a non-metastatic to metastatic phenotype is the loss of function of metastasis-suppressor genes. These
5 genes specifically suppress the ability of a cell to metastasize. Work from several groups has demonstrated that human chromosomes 8, 10, 11 and 17 encode prostate cancer metastasis suppressor activities. However, other human chromosomes such as chromosomes 1, 7, 13, 16, and 18 may also be associated to prostate cancer.

It thus remains to localize and to identify the genes specifically involved in the development
10 and the progression of prostate cancers starting from the genetic analysis of the hereditary and the non-hereditary forms and to define their clinical implications in terms of prognosis and therapeutic innovations.

SUMMARY OF THE INVENTION

The present invention pertains to nucleic acid molecules comprising the genomic sequence
15 of a novel human gene which encodes a TBC-1 protein. The *TBC-1* genomic sequences comprise regulatory sequence located upstream (5'-end) and downstream (3'-end) of the transcribed portion of said gene, these regulatory sequences being also part of the invention. The human *TBC-1* genomic sequence is included in a previously unknown candidate region of prostate cancer located on chromosome 4.

20 The invention also deals with the two complete cDNA sequences encoding the TBC-1 protein, as well as with the corresponding translation product.

Oligonucleotide probes or primers hybridizing specifically with a *TBC-1* genomic or cDNA sequence are also part of the present invention, as well as DNA amplification and detection methods using said primers and probes.

25 A further object of the invention consists of recombinant vectors comprising any of the nucleic acid sequences described above, and in particular of recombinant vectors comprising a *TBC-1* regulatory sequence or a sequence encoding a TBC-1 protein, as well as of cell hosts and transgenic non human animals comprising said nucleic acid sequences or recombinant vectors.

The invention also concerns a *TBC-1*-related biallelic marker and the use thereof.

30 Finally, the invention is directed to methods for the screening of substances or molecules that inhibit the expression of *TBC-1*, as well as with methods for the screening of substances or molecules that interact with a TBC-1 polypeptide.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 : An amino acid alignment of a portion of the amino acid sequence of the TBC-1 protein of SEQ ID No 5 with other proteins sharing amino acid homology with TBC-1. The amino acid numbering refers to the murine TBC-1.

5 **Brief Description of the sequences provided in the Sequence Listing**

SEQ ID No 1 contains a first part of the *TBC-1* genomic sequence comprising the 5' regulatory sequence and the exons 1, 1bis, and 2.

SEQ ID No 2 contains a second part of the *TBC-1* genomic sequence comprising the 12 last exons of the *TBC-1* gene and the 3' regulatory sequence.

10 SEQ ID No 3 contains a first cDNA sequence of the *TBC-1* gene.

SEQ ID No 4 contains a second cDNA sequence of the *TBC-1* gene.

SEQ ID No 5 contains the amino acid sequence encoded by the cDNAs of SEQ ID Nos 3 and 4.

15 SEQ ID No 6 contains a primer containing the additional PU 5' sequence described further in Example 3.

SEQ ID No 7 contains a primer containing the additional RP 5' sequence described further in Example 3.

In accordance with the regulations relating to Sequence Listings, the following codes have been used in the Sequence Listing to indicate the locations of biallelic markers within the sequences and to identify each of the alleles present at the polymorphic base. The code "r" in the sequences indicates that one allele of the polymorphic base is a guanine, while the other allele is an adenine. The code "y" in the sequences indicates that one allele of the polymorphic base is a thymine, while the other allele is a cytosine. The code "m" in the sequences indicates that one allele of the polymorphic base is an adenine, while the other allele is an cytosine. The code "k" in the sequences indicates that one allele of the polymorphic base is a guanine, while the other allele is a thymine. The code "s" in the sequences indicates that one allele of the polymorphic base is a guanine, while the other allele is a cytosine. The code "w" in the sequences indicates that one allele of the polymorphic base is an adenine, while the other allele is an thymine. The nucleotide code of the original allele for each biallelic marker is the following:

30	Biallelic marker	Original allele
	99-430-352	G
	99-20508-456	C
	99-20469-213	C
	5-254-227	A
35	5-257-353	C
	99-20511-32	T

	99-20511-221	A
	99-20504-90	G
	99-20493-238	A
	99-20499-221	G
5	99-20499-364	A
	99-20499-399	A
	5-249-304	G
	99-20485-269	A
	99-20481-131	G
10	99-20481-419	T
	99-20480-233	A

DETAILED DESCRIPTION OF THE INVENTION

The present invention concerns polynucleotides and polypeptides related to the human *TBC-1* gene (also termed "*TBC-1* gene" throughout the present specification), which is potentially involved in the regulation of the differentiation of various cell types in mammals. A deregulation or an alteration of *TBC-1* expression, or alternatively an alteration in the amino acid sequence of the *TBC-1* protein may be involved in the generation of a pathological state related to cell differentiation in a patient, more particularly to abnormal cell proliferation leading to cancer states, such as prostate cancer.

20

Definitions

Before describing the invention in greater detail, the following definitions are set forth to illustrate and define the meaning and scope of the terms used to describe the invention herein.

The term "*TBC-1* gene", when used herein, encompasses mRNA and cDNA sequences encoding the *TBC-1* protein. In the case of a genomic sequence, the *TBC-1* gene also includes native regulatory regions which control the expression of the coding sequence of the *TBC-1* gene.

The term "functionally active fragment" of the *TBC-1* protein is intended to designate a polypeptide carrying at least one of the structural features of the *TBC-1* protein involved in at least one of the biological functions and/or activity of the *TBC-1* protein.

A "heterologous" or "exogenous" polynucleotide designates a purified or isolated nucleic acid that has been placed, by genetic engineering techniques, in the environment of unrelated nucleotide sequences, such as the final polynucleotide construct does not occur naturally. An illustrative, but not limitative, embodiment of such a polynucleotide construct may be represented by a polynucleotide comprising (1) a regulatory polynucleotide derived from the *TBC-1* gene sequence and (2) a polynucleotide encoding a cytokine, for example GM-CSF. The polypeptide

encoded by the heterologous polynucleotide will be termed an heterologous polypeptide for the purpose of the present invention.

By a "biologically active fragment or variant" of a regulatory polynucleotide according to the present invention is intended a polynucleotide comprising or alternatively consisting in a
5 fragment of said polynucleotide which is functional as a regulatory region for expressing a recombinant polypeptide or a recombinant polynucleotide in a recombinant cell host.

For the purpose of the invention, a nucleic acid or polynucleotide is "functional" as a regulatory region for expressing a recombinant polypeptide or a recombinant polynucleotide if said regulatory polynucleotide contains nucleotide sequences which contain transcriptional and
10 translational regulatory information, and such sequences are "operatively linked" to nucleotide sequences which encode the desired polypeptide or the desired polynucleotide. An operable linkage is a linkage in which the regulatory nucleic acid and the DNA sequence sought to be expressed are linked in such a way as to permit gene expression.

A "promoter" refers to a DNA sequence recognized by the synthetic machinery of the cell
15 required to initiate the specific transcription of a gene.

A sequence which is "operably linked" to a regulatory sequence such as a promoter means that said regulatory element is in the correct location and orientation in relation to the nucleic acid to control RNA polymerase initiation and expression of the nucleic acid of interest.

As used herein, the term "operably linked" refers to a linkage of polynucleotide elements in
20 a functional relationship. For instance, a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the coding sequence. More precisely, two DNA molecules (such as a polynucleotide containing a promoter region and a polynucleotide encoding a desired polypeptide or polynucleotide) are said to be "operably linked" if the nature of the linkage between the two polynucleotides does not (1) result in the introduction of a frame-shift mutation or (2)
25 interfere with the ability of the polynucleotide containing the promoter to direct the transcription of the coding polynucleotide. The promoter polynucleotide would be operably linked to a polynucleotide encoding a desired polypeptide or a desired polynucleotide if the promoter is capable of effecting transcription of the polynucleotide of interest.

The term "primer" denotes a specific oligonucleotide sequence which is complementary to
30 a target nucleotide sequence and used to hybridize to the target nucleotide sequence. A primer serves as an initiation point for nucleotide polymerization catalyzed by either DNA polymerase, RNA polymerase or reverse transcriptase.

The term "probe" denotes a defined nucleic acid segment (or nucleotide analog segment, e.g., polynucleotide as defined hereinbelow) which can be used to identify a specific polynucleotide
35 sequence present in samples, said nucleic acid segment comprising a nucleotide sequence complementary of the specific polynucleotide sequence to be identified.

The terms "sample" or "material sample" are used herein to designate a solid or a liquid material suspected to contain a polynucleotide or a polypeptide of the invention. A solid material may be, for example, a tissue slice or biopsy within which is searched the presence of a polynucleotide encoding a TBC-1 protein, either a DNA or RNA molecule or within which is
5 searched the presence of a native or a mutated TBC-1 protein, or alternatively the presence of a desired protein of interest the expression of which has been placed under the control of a *TBC-1* regulatory polynucleotide. A liquid material may be, for example, any body fluid like serum, urine etc., or a liquid solution resulting from the extraction of nucleic acid or protein material of interest from a cell suspension or from cells in a tissue slice or biopsy. The term "biological sample" is also
10 used and is more precisely defined within the Section dealing with DNA extraction.

As used herein, the term "purified" does not require absolute purity; rather, it is intended as a relative definition. Purification of starting material or natural material to at least one order of magnitude, preferably two or three orders, and more preferably four or five orders of magnitude is expressly contemplated. As an example, purification from 0.1% concentration to 10% concentration
15 is two orders of magnitude.

The term "isolated" requires that the material be removed from its original environment (e.g. the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or DNA or polypeptide, separated from some or all of the coexisting materials in the natural system,
20 is isolated. Such polynucleotide could be part of a vector and/or such polynucleotide or polypeptide could be part of a composition and still be isolated in that the vector or composition is not part of its natural environment.

The term "polypeptide" refers to a polymer of amino acids without regard to the length of the polymer; thus, peptides, oligopeptides, and proteins are included within the definition of
25 polypeptide. This term also does not specify or exclude post-expression modifications of polypeptides, for example, polypeptides which include the covalent attachment of glycosyl groups, acetyl groups, phosphate groups, lipid groups and the like are expressly encompassed by the term polypeptide. Also included within the definition are polypeptides which contain one or more analogs of an amino acid (including, for example, non-naturally occurring amino acids, amino acids
30 which only occur naturally in an unrelated biological system, modified amino acids from mammalian systems etc.), polypeptides with substituted linkages, as well as other modifications known in the art, both naturally occurring and non-naturally occurring.

The term "recombinant polypeptide" is used herein to refer to polypeptides that have been artificially designed and which comprise at least two polypeptide sequences that are not found as
35 contiguous polypeptide sequences in their initial natural environment, or to refer to polypeptides which have been expressed from a recombinant polynucleotide.

The term "purified" is used herein to describe a polypeptide of the invention which has been separated from other compounds including, but not limited to nucleic acids, lipids, carbohydrates and other proteins. A polypeptide is substantially pure when at least about 50%, preferably 60 to 75% of a sample exhibits a single polypeptide sequence. A substantially pure polypeptide typically
5 comprises about 50%, preferably 60 to 90% weight/weight of a protein sample, more usually about 95%, and preferably is over about 99% pure. Polypeptide purity or homogeneity is indicated by a number of means well known in the art, such as polyacrylamide gel electrophoresis of a sample, followed by visualizing a single polypeptide band upon staining the gel. For certain purposes higher resolution can be provided by using HPLC or other means well known in the art.

10 As used herein, the term "non-human animal" refers to any non-human vertebrate, birds and more usually mammals, preferably primates, farm animals such as swine, goats, sheep, donkeys, and horses, rabbits or rodents, more preferably rats or mice. As used herein, the term "animal" is used to refer to any vertebrate, preferable a mammal. Both the terms "animal" and "mammal" expressly embrace human subjects unless preceded with the term "non-human".

15 As used herein, the term "antibody" refers to a polypeptide or group of polypeptides which are comprised of at least one binding domain, where an antibody binding domain is formed from the folding of variable domains of an antibody molecule to form three-dimensional binding spaces with an internal surface shape and charge distribution complementary to the features of an antigenic determinant of an antigen, which allows an immunological reaction with the antigen. Antibodies
20 include recombinant proteins comprising the binding domains, as wells as fragments, including Fab, Fab', F(ab)₂, and F(ab')₂ fragments.

As used herein, an "antigenic determinant" is the portion of an antigen molecule, in this case a TBC-1 polypeptide, that determines the specificity of the antigen-antibody reaction. An "epitope" refers to an antigenic determinant of a polypeptide. An epitope can comprise as few as 3
25 amino acids in a spatial conformation which is unique to the epitope. Generally an epitope consists of at least 6 such amino acids, and more usually at least 8-10 such amino acids. Methods for determining the amino acids which make up an epitope include x-ray crystallography, 2-dimensional nuclear magnetic resonance, and epitope mapping e.g. the Pepscan method described by Geysen et al. 1984; PCT Publication No. WO 84/03564; and PCT Publication No. WO
30 84/03506.

Throughout the present specification, the expression "nucleotide sequence" may be employed to designate indifferently a polynucleotide or an oligonucleotide or a nucleic acid. More precisely, the expression "nucleotide sequence" encompasses the nucleic material itself and is thus not restricted to the sequence information (i.e. the succession of letters chosen among the four base
35 letters) that biochemically characterizes a specific DNA or RNA molecule.

As used interchangeably herein, the term "oligonucleotides", and "polynucleotides" include RNA, DNA, or RNA/DNA hybrid sequences of more than one nucleotide in either single chain or

duplex form. The term "nucleotide" as used herein as an adjective to describe molecules comprising RNA, DNA, or RNA/DNA hybrid sequences of any length in single-stranded or duplex form. The term "nucleotide" is also used herein as a noun to refer to individual nucleotides or varieties of nucleotides, meaning a molecule, or individual unit in a larger nucleic acid molecule, comprising a purine or pyrimidine, a ribose or deoxyribose sugar moiety, and a phosphate group, or phosphodiester linkage in the case of nucleotides within an oligonucleotide or polynucleotide. Although the term "nucleotide" is also used herein to encompass "modified nucleotides" which comprise at least one modification (a) an alternative linking group, (b) an analogous form of purine, (c) an analogous form of pyrimidine, or (d) an analogous sugar, for examples of analogous linking groups, purine, pyrimidines, and sugars see for example PCT publication No WO 95/04064. However, the polynucleotides of the invention are preferably comprised of greater than 50% conventional deoxyribose nucleotides, and most preferably greater than 90% conventional deoxyribose nucleotides. The polynucleotide sequences of the invention may be prepared by any known method, including synthetic, recombinant, *ex vivo* generation, or a combination thereof, as well as utilizing any purification methods known in the art.

The term "heterozygosity rate" is used herein to refer to the incidence of individuals in a population which are heterozygous at a particular allele. In a biallelic system, the heterozygosity rate is on average equal to $2P_a(1-P_a)$, where P_a is the frequency of the least common allele. In order to be useful in genetic studies, a genetic marker should have an adequate level of heterozygosity to allow a reasonable probability that a randomly selected person will be heterozygous.

The term "genotype" as used herein refers the identity of the alleles present in an individual or a sample. In the context of the present invention a genotype preferably refers to the description of the biallelic marker alleles present in an individual or a sample. The term "genotyping" a sample or an individual for a biallelic marker consists of determining the specific allele or the specific nucleotide carried by an individual at a biallelic marker.

The term "polymorphism" as used herein refers to the occurrence of two or more alternative genomic sequences or alleles between or among different genomes or individuals. "Polymorphic" refers to the condition in which two or more variants of a specific genomic sequence can be found in a population. A "polymorphic site" is the locus at which the variation occurs. A single nucleotide polymorphism is a single base pair change. Typically a single nucleotide polymorphism is the replacement of one nucleotide by another nucleotide at the polymorphic site. Deletion of a single nucleotide or insertion of a single nucleotide, also give rise to single nucleotide polymorphisms. In the context of the present invention "single nucleotide polymorphism" preferably refers to a single nucleotide substitution. However, the polymorphism can also involve an insertion or a deletion of at least one nucleotide, preferably between 1 and 5 nucleotides. Typically, between different genomes or between different individuals, the polymorphic site may be occupied by two different nucleotides.

The term “biallelic polymorphism” and “biallelic marker” are used interchangeably herein to refer to a single nucleotide polymorphism having two alleles at a fairly high frequency in the population. A “biallelic marker allele” refers to the nucleotide variants present at a biallelic marker site. Typically, the frequency of the less common allele of the biallelic markers of the present invention has been validated to be greater than 1%, preferably the frequency is greater than 10%, more preferably the frequency is at least 20% (i.e. heterozygosity rate of at least 0.32), even more preferably the frequency is at least 30% (i.e. heterozygosity rate of at least 0.42). A biallelic marker wherein the frequency of the less common allele is 30% or more is termed a “high quality biallelic marker”.

10 The location of nucleotides in a polynucleotide with respect to the center of the polynucleotide are described herein in the following manner. When a polynucleotide has an odd number of nucleotides, the nucleotide at an equal distance from the 3' and 5' ends of the polynucleotide is considered to be “at the center” of the polynucleotide, and any nucleotide immediately adjacent to the nucleotide at the center, or the nucleotide at the center itself is
15 considered to be “within 1 nucleotide of the center.” With an odd number of nucleotides in a polynucleotide any of the five nucleotides positions in the middle of the polynucleotide would be considered to be within 2 nucleotides of the center, and so on. When a polynucleotide has an even number of nucleotides, there would be a bond and not a nucleotide at the center of the polynucleotide. Thus, either of the two central nucleotides would be considered to be “within 1
20 nucleotide of the center” and any of the four nucleotides in the middle of the polynucleotide would be considered to be “within 2 nucleotides of the center”, and so on. For polymorphisms which involve the substitution, insertion or deletion of 1 or more nucleotides, the polymorphism, allele or biallelic marker is “at the center” of a polynucleotide if the difference between the distance from the substituted, inserted, or deleted polynucleotides of the polymorphism and the 3' end of the
25 polynucleotide, and the distance from the substituted, inserted, or deleted polynucleotides of the polymorphism and the 5' end of the polynucleotide is zero or one nucleotide. If this difference is 0 to 3, then the polymorphism is considered to be “within 1 nucleotide of the center.” If the difference is 0 to 5, the polymorphism is considered to be “within 2 nucleotides of the center.” If the difference is 0 to 7, the polymorphism is considered to be “within 3 nucleotides of the center,” and
30 so on.

As used herein the terminology “defining a biallelic marker” means that a sequence includes a polymorphic base from a biallelic marker. The sequences defining a biallelic marker may be of any length consistent with their intended use, provided that they contain a polymorphic base from a biallelic marker. The sequence has between 1 and 500 nucleotides in length, preferably
35 between 5, 10, 15, 20, 25, or 40 and 200 nucleotides and more preferably between 30 and 50 nucleotides in length. Each biallelic marker therefore corresponds to two forms of a polynucleotide sequence included in a gene, which, when compared with one another, present a nucleotide

modification at one position. Preferably, the sequences defining a biallelic marker include a polymorphic base selected from the group consisting of the biallelic markers A1 to A19 and the complements thereof. In some embodiments the sequences defining a biallelic marker comprise one of the sequences selected from the group consisting of P1 to P7, P9 to P13, P15 to P19 and the complementary sequences thereto. Likewise, the term "marker" or "biallelic marker" requires that the sequence is of sufficient length to practically (although not necessarily unambiguously) identify the polymorphic allele, which usually implies a length of at least 4, 5, 6, 10, 15, 20, 25, or 40 nucleotides.

The term "upstream" is used herein to refer to a location which is toward the 5' end of the polynucleotide from a specific reference point.

The terms "base paired" and "Watson & Crick base paired" are used interchangeably herein to refer to nucleotides which can be hydrogen bonded to one another by virtue of their sequence identities in a manner like that found in double-helical DNA with thymine or uracil residues linked to adenine residues by two hydrogen bonds and cytosine and guanine residues linked by three hydrogen bonds (See Stryer, L., *Biochemistry*, 4th edition, 1995).

The terms "complementary" or "complement thereof" are used herein to refer to the sequences of polynucleotides which is capable of forming Watson & Crick base pairing with another specified polynucleotide throughout the entirety of the complementary region. For the purpose of the present invention, a first polynucleotide is deemed to be complementary to a second polynucleotide when each base in the first polynucleotide is paired with its complementary base. Complementary bases are, generally, A and T (or A and U), or C and G. "Complement" is used herein as a synonym from "complementary polynucleotide", "complementary nucleic acid" and "complementary nucleotide sequence". These terms are applied to pairs of polynucleotides based solely upon their sequences and not any particular set of conditions under which the two polynucleotides would actually bind.

Variants and fragments

1. Polynucleotides

The invention also relates to variants and fragments of the polynucleotides described herein, particularly of a *TBC-1* gene containing one or more biallelic markers according to the invention.

Variants of polynucleotides, as the term is used herein, are polynucleotides that differ from a reference polynucleotide. A variant of a polynucleotide may be a naturally occurring variant such as a naturally occurring allelic variant, or it may be a variant that is not known to occur naturally. Such non-naturally occurring variants of the polynucleotide may be made by mutagenesis techniques, including those applied to polynucleotides, cells or organisms. Generally, differences are limited so that the nucleotide sequences of the reference and the variant are closely similar overall and, in many regions, identical.

5 Variants of polynucleotides according to the invention include, without being limited to, nucleotide sequences that are at least 95% identical to any of SEQ ID Nos 1-4 or the sequences complementary thereto or to any polynucleotide fragment of at least 8 consecutive nucleotides of any of SEQ ID Nos 1-4 or the sequences complementary thereto, and preferably at least 98% identical, more particularly at least 99.5% identical, and most preferably at least 99.9% identical to any of SEQ ID Nos 1-4 or the sequences complementary thereto or to any polynucleotide fragment of at least 8 consecutive nucleotides of any of SEQ ID Nos 1-4 or the sequences complementary thereto.

10 Changes in the nucleotide of a variant may be silent, which means that they do not alter the amino acids encoded by the polynucleotide.

However, nucleotide changes may also result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence. The substitutions, deletions or additions may involve one or more nucleotides. The variants may be altered in coding or non-coding regions or both. Alterations in the coding regions may produce conservative or non-conservative amino acid substitutions, deletions or additions.

15 In the context of the present invention, particularly preferred embodiments are those in which the polynucleotides encode polypeptides which retain substantially the same biological function or activity as the mature TBC-1 protein.

A polynucleotide fragment is a polynucleotide having a sequence that entirely is the same as part but not all of a given nucleotide sequence, preferably the nucleotide sequence of a *TBC-1* gene, and variants thereof. The fragment can be a portion of an exon or of an intron of a *TBC-1* gene. It can also be a portion of the regulatory sequences of the *TBC-1* gene. Preferably, such fragments comprise the polymorphic base of a biallelic marker selected from the group consisting of the biallelic markers A1 to A19 and the complements thereof.

25 Such fragments may be "free-standing", i.e. not part of or fused to other polynucleotides, or they may be comprised within a single larger polynucleotide of which they form a part or region. However, several fragments may be comprised within a single larger polynucleotide.

As representative examples of polynucleotide fragments of the invention, there may be mentioned those which have from about 4, 6, 8, 15, 20, 25, 40, 10 to 20, 10 to 30, 30 to 55, 50 to 100, 75 to 100 or 100 to 200 nucleotides in length. Preferred are those fragments having about 49 nucleotides in length, such as those of P1 to P7, P9 to P13, P15 to P19 or the sequences complementary thereto and containing at least one of the biallelic markers of a *TBC-1* gene which are described herein.

2. Polypeptides.

35 The invention also relates to variants, fragments, analogs and derivatives of the polypeptides described herein, including mutated TBC-1 proteins.

The variant may be 1) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code, or 2) one in which one or more of the amino acid residues includes a substituent group, or 3) one in which the mutated TBC-1 is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or 4) one in which the additional amino acids are fused to the mutated TBC-1, such as a leader or secretory sequence or a sequence which is employed for purification of the mutated TBC-1 or a preprotein sequence. Such variants are deemed to be within the scope of those skilled in the art.

10 More particularly, a variant TBC-1 polypeptide comprises amino acid changes ranging from 1, 2, 3, 4, 5, 10 to 20 substitutions, additions or deletions of one amino acid, preferably from 1 to 10, more preferably from 1 to 5 and most preferably from 1 to 3 substitutions, additions or deletions of one amino acid. The preferred amino acid changes are those which have little or no influence on the biological activity or the capacity of the variant TBC-1 polypeptide to be recognized by antibodies
15 raised against a native TBC-1 protein.

By homologous peptide according to the present invention is meant a polypeptide containing one or several amino acid additions, deletions and/or substitutions in the amino acid sequence of a TBC-1 polypeptide. In the case of an amino acid substitution, one or several consecutive or non-consecutive amino acids are replaced by « equivalent » amino acids.

20 The expression "equivalent" amino acid is used herein to designate any amino acid that may be substituted for one of the amino acids having similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydrophobic nature of the polypeptide to be substantially unchanged. Generally, the following groups of amino acids represent equivalent changes: (1) Ala, Pro, Gly, Glu, Asp, Gln, Asn, Ser, Thr; (2) Cys, Ser, Tyr, Thr; (3) Val, Ile, Leu, Met, Ala, Phe; (4) Lys, Arg, His; (5) Phe, Tyr, Trp, His.
25

By an equivalent amino acid according to the present invention is also meant the replacement of a residue in the L-form by a residue in the D form or the replacement of a Glutamic acid (E) residue by a Pyro-glutamic acid compound. The synthesis of peptides containing at least one residue in the D-form is, for example, described by Koch (1977).

30 A specific, but not restrictive, embodiment of a modified peptide molecule of interest according to the present invention, which consists in a peptide molecule which is resistant to proteolysis, is a peptide in which the -CONH- peptide bond is modified and replaced by a (CH₂NH) reduced bond, a (NHCO) retro inverso bond, a (CH₂-O) methylene-oxy bond, a (CH₂-S) thiomethylene bond, a (CH₂CH₂) carba bond, a (CO-CH₂) cetomethylene bond, a (CHOH-CH₂) hydroxyethylene bond), a (N-N) bound, a E-alcene bond or also a -CH=CH- bond.
35

The polypeptide according to the invention could have post-translational modifications. For example, it can present the following modifications: acylation, disulfide bond formation, prenylation, carboxymethylation and phosphorylation.

A polypeptide fragment is a polypeptide having a sequence that entirely is the same as part but not all of a given polypeptide sequence, preferably a polypeptide encoded by a *TBC-1* gene and variants thereof. Preferred fragments include those regions possessing antigenic properties and which can be used to raise antibodies against the TBC-1 protein.

Such fragments may be "free-standing", i.e. not part of or fused to other polypeptides, or they may be comprised within a single larger polypeptide of which they form a part or region.

10 However, several fragments may be comprised within a single larger polypeptide.

As representative examples of polypeptide fragments of the invention, there may be mentioned those which comprise at least about 5, 6, 7, 8, 9 or 10 to 15, 10 to 20, 15 to 40, or 30 to 55 amino acids of the TBC-1. In some embodiments, the fragments contain at least one amino acid mutation in the TBC-1 protein.

15 Identity Between Nucleic Acids Or Polypeptides

The terms "percentage of sequence identity" and "percentage homology" are used interchangeably herein to refer to comparisons among polynucleotides and polypeptides, and are determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity. Homology is evaluated using any of the variety of sequence comparison algorithms and programs known in the art. Such algorithms and programs include, but are by no means limited to, TBLASTN, BLASTP, FASTA, TFASTA, and CLUSTALW (Pearson and Lipman, 1988; Altschul et al., 1990; Thompson et al., 1994; Higgins et al., 1996; Altschul et al., 1993). In a particularly preferred embodiment, protein and nucleic acid sequence homologies are evaluated using the Basic Local Alignment Search Tool ("BLAST") which is well known in the art (see, e.g., Karlin and Altschul, 1990; Altschul et al., 1990, 1993, 1997). In particular, five specific BLAST programs are used to perform the following task:

- (1) BLASTP and BLAST3 compare an amino acid query sequence against a protein sequence database;
- 35 (2) BLASTN compares a nucleotide query sequence against a nucleotide sequence database;

(3) BLASTX compares the six-frame conceptual translation products of a query nucleotide sequence (both strands) against a protein sequence database;

(4) TBLASTN compares a query protein sequence against a nucleotide sequence database translated in all six reading frames (both strands); and

5 (5) TBLASTX compares the six-frame translations of a nucleotide query sequence against the six-frame translations of a nucleotide sequence database.

The BLAST programs identify homologous sequences by identifying similar segments, which are referred to herein as "high-scoring segment pairs," between a query amino or nucleic acid sequence and a test sequence which is preferably obtained from a protein or nucleic acid sequence database.

10 High-scoring segment pairs are preferably identified (i.e., aligned) by means of a scoring matrix, many of which are known in the art. Preferably, the scoring matrix used is the BLOSUM62 matrix (Gonnet et al., 1992; Henikoff and Henikoff, 1993). Less preferably, the PAM or PAM250 matrices may also be used (see, e.g., Schwartz and Dayhoff, eds., 1978). The BLAST programs evaluate the statistical significance of all high-scoring segment pairs identified, and preferably
15 selects those segments which satisfy a user-specified threshold of significance, such as a user-specified percent homology. Preferably, the statistical significance of a high-scoring segment pair is evaluated using the statistical significance formula of Karlin (see, e.g., Karlin and Altschul, 1990).

Stringent Hybridization Conditions

20 By way of example and not limitation, procedures using conditions of high stringency are as follows: Prehybridization of filters containing DNA is carried out for 8 h to overnight at 65°C in buffer composed of 6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 µg/ml denatured salmon sperm DNA. Filters are hybridized for 48 h at 65°C, the preferred hybridization temperature, in prehybridization mixture containing 100 µg/ml
25 denatured salmon sperm DNA and 5-20 X 10⁶ cpm of ³²P-labeled probe. Alternatively, the hybridization step can be performed at 65°C in the presence of SSC buffer, 1 x SSC corresponding to 0.15M NaCl and 0.05 M Na citrate. Subsequently, filter washes can be done at 37°C for 1 h in a solution containing 2 x SSC, 0.01% PVP, 0.01% Ficoll, and 0.01% BSA, followed by a wash in 0.1 X SSC at 50°C for 45 min. Alternatively, filter washes can be performed in a solution containing 2
30 x SSC and 0.1% SDS, or 0.5 x SSC and 0.1% SDS, or 0.1 x SSC and 0.1% SDS at 68°C for 15 minute intervals. Following the wash steps, the hybridized probes are detectable by autoradiography. Other conditions of high stringency which may be used are well known in the art and as cited in Sambrook et al., 1989; and Ausubel et al., 1989, are incorporated herein in their entirety. These hybridization conditions are suitable for a nucleic acid molecule of about 20
35 nucleotides in length. There is no need to say that the hybridization conditions described above are to be adapted according to the length of the desired nucleic acid, following techniques well known to the one skilled in the art. The suitable hybridization conditions may for example be adapted

according to the teachings disclosed in the book of Hames and Higgins (1985) or in Sambrook et al.(1989).

Candidate Region On The Chromosome 4 (Linkage Analysis).

In order to localize the prostate cancer gene(s) starting from families, a systematic familial
5 study of genetic link research is carried out using markers of the microsatellite type described at the Genethon laboratory by the Jean Weissenbach team (Dib et al., 1996).

The studies of genetic link or of "linkage" are based on the principle according to which
two neighboring sequences on a chromosome do not present (or very rarely present) recombinations
by crossing-over during meiosis. To do this, microsatellite DNA sequences (chromosomal markers)
10 constantly co-inherited with the disease studied are searched for in a family having a predisposition
for this disease. These DNA sequences organized in the form of a repetition of di-, tri- or
tetranucleotides are systematically present along the genome, and thus allow the identification of
chromosomal fragments harboring them. More than 5000 microsatellite markers, have been
localized with precision on the genome as a result of the first studies on the genetic map carried out
15 at Genethon under the supervision of Jean Weissenbach, and on the physical map (using the "Yeast
Artificial Chromosomes"), work conducted by Daniel Cohen at C.E.P.H. and at Genethon
(Chumakov et al., 1995). Genetic link analysis calculates the probabilities of recombinations of the
target gene with the chromosomal markers used, according to the genealogical tree, the transmission
of the disease, and the transmission of the markers. Thus if a particular allele of a given marker is
20 transmitted with the disease more often than chance would have it (recombination level of between
0 and 0.5), it is possible to deduce that the target gene in question is found in the neighborhood of
the marker. Using this technique, it has been possible to localize several genes of genetic
predisposition to familial cancers. In order to be able to be included in a genetic link study, the
families affected by a hereditary form of the disease must satisfy the "informativeness" criteria:
25 several affected subjects (and whose constitutional DNA is available) per generation, and at best
having a large number of siblings.

By linkage analysis, the inventors have identified a candidate region for prostate cancer on
chromosome 4. Indeed, the LOD scores at 2 points between the disease and the markers on a total
population of approximately fifty families present a value of 2.49 for marker D4S398 which
30 indicates a probable genetic link with this marker. The curve of the variation of the LOD score on a
map of 5 markers is centered on D4S398 and the value higher than 3.3 indicates that a gene
involved in familial prostate cancer is probably found in the region located between markers
D4S2978 and D4S3018, or a space of approximately 9.7 cM.

Homologies Of The Novel Human Gene Translation Product With A Known Murine Protein.

A novel human gene was found in this candidate region. It presents a good probability to be involved in cancer. Database homology searches have allowed the inventors to determine that the translation product of this novel human gene has significant identity with a murine protein called *tbc1*. Therefore, the novel human gene of the invention has thus been called *TBC-1* throughout the present specification. *TBC-1* comprises an open Reading frame that encodes a novel protein, the TBC-1 protein. Based on sequence similarity, an alignment of a portion of the TBC-1 amino acid sequence with the known *tbc1* murine protein, it is expected that TBC1 protein may play a role in the cell cycle and in differentiation of various tissues. Indeed, the TBC1 protein contains a 200 amino acid domain called the TBC domain that is homologous to regions in the *tre2*-oncogene and in the yeast regulators of mitosis BUB2 and *cdc16*.

The cDNA of the murine *tbc1* gene has been described in US Patent No US 5,700,927 and it encodes a putative protein product of 1141 amino acids. The N-terminus of the murine *tbc1* protein contains stretches of cysteines and histidines which may form zinc finger structures in the mature polypeptides. The N-terminus also comprises short stretches of basic amino acids which may be involved in a nuclear localization signal. The TBC domain of the murine *tbc1* protein contains several tyrosine residues which are conserved in BUB2 and *cdc16*. The C-terminus of the murine *tbc1* protein contains a long stretch of evenly spaced leucine residues which are susceptible to form a leucine zipper motif.

The murine *tbc1* gene has been shown to be highly expressed in testis and kidney. However, lower levels of expression have also been identified in lung, spleen, brain, and heart. Moreover, murine *tbc1* is a nuclear protein which is expressed in a cell- and stage-specific manner.

Studies of murine bone marrow have demonstrated that erythroid cells and megakaryocytes expressed substantial levels of the murine *tbc1* protein, but none was detected in mature neutrophils. Similarly, spermatogonia do not express murine *tbc1*, but primary and secondary spermatocytes express abundant *tbc1*. Later in the differentiation of the germ cells, the *tbc1* levels appear to decrease in spermatids and active sperm. The differentiation program of spermatogonia to spermatocytes therefore involves a significant upregulation of murine *tbc1* expression.

The general distribution of murine *tbc1* is not tissue-specific, but is cell-specific within individual tissues and intimately linked to tissue differentiation. The developmental expression of murine *tbc1*, particularly in hematopoietic and germ cells, suggests that this gene plays a role in the terminal differentiation program of several tissues.

Consequently, an alteration in the expression of the *TBC-1* gene or in the amino acid sequence of the TBC-1 protein leading to an altered biological activity of the latter is likely to cause, directly or indirectly, cell proliferation disorders and thus diseases related to an abnormal cell proliferation such as cancer, particularly prostate cancer.

Genomic Sequence Of *TBC-1*

The present invention concerns the genomic sequence of *TBC-1*. The present invention encompasses the *TBC-1* gene, or *TBC-1* genomic sequences consisting of, consisting essentially of, or comprising a sequence selected from the group consisting of SEQ ID Nos 1 and 2, a sequence
5 complementary thereto, as well as fragments and variants thereof. These polynucleotides may be purified, isolated, or recombinant.

The inventors have sequenced two portions of the *TBC-1* genomic sequence. The first portion of the *TBC-1* gene sequence contains the three first exons of the *TBC-1* gene, designated as Exon 1, Exon 1bis and Exon 2, and the 5' regulatory sequence located upstream of the transcribed
10 sequences. The sequence of the first portion of the genomic sequence is disclosed in SEQ ID No 1. The second portion contains the twelve last exons of the *TBC-1* gene, designated as exons A, B, C, D, E, F, G, H, I, J, K, and L, and the 3' regulatory sequence which is located downstream of the transcribed sequences.

The exon positions in SEQ ID Nos 1 and 2 are detailed below in Table A.

15

Table A

Exon	Position in SEQ ID No 1		Intron	Position in SEQ ID No 1	
	Beginning	End		Beginning	End
1	2001	2077	1	2078	12739
1bis	12292	12373	1bis	12374	12739
2	12740	13249	2	13250	at least 17590
Exon	Position in SEQ ID No 2		Intron	Position in SEQ ID No 2	
	Beginning	End		Beginning	End
A	4661	4789	A	4790	6115
B	6116	6202	B	6203	9918
C	9919	10199	C	10200	14520
D	14521	14660	D	14661	50256
E	50257	50442	E	50443	56255
F	56256	56417	F	56418	63325
G	63326	63484	G	63485	76035
H	76036	76280	H	76281	78363
I	78364	78523	I	78524	85294
J	85295	85464	J	85465	93416
K	93417	93590	K	93591	97475
L	97476	97960			

Intron 1 refers to the nucleotide sequence located between Exon 1 and Exon 2; Intron 1bis refers to the nucleotide sequence located between Exon 1bis and Exon 2; Intron A refers to the nucleotide sequence located between Exon A and Exon B; and so on. The position of the introns is
20 detailed in Table A.

The *TBC-1* introns defined hereinafter for the purpose of the present invention are not exactly what is generally understood as "introns" by the one skilled in the art and will consequently be further defined below.

Generally, an intron is defined as a nucleotide sequence that is present both in the genomic DNA and in the unspliced mRNA molecule, and which is absent from the mRNA molecule which has already gone through splicing events. In the case of the *TBC-1* gene, the inventors have found that at least two different spliced mRNA molecules are produced when this gene is transcribed, as it will be described in detail in a further section of the specification. The first spliced mRNA molecule comprises Exons 1 and 2. Thus, the genomic nucleotide sequence comprised between Exon 1 and Exon 2 is an intronic sequence as regards to this first mRNA molecule, despite the fact that this intronic sequence contains Exon 1bis. In contrast, Exon 1bis is of course an exonic nucleotide sequence as regards to the second *TBC-1* mRNA molecule.

For the purpose of the present invention and in order to make a clear and unambiguous designation of the different nucleic acids encompassed, it has been postulated that the polynucleotides contained both in any of the nucleotide sequences of SEQ ID Nos 1 or 2 and in any of the nucleotide sequences of SEQ ID Nos 3 or 4 are considered as exonic sequences. Conversely, the polynucleotides contained in any of the nucleotide sequences of SEQ ID Nos 1 or 2 but which are absent both from the nucleotide sequence of SEQ ID No 3 and from the nucleotide sequence of SEQ ID No 4 are considered as intronic sequences.

The nucleic acids defining the *TBC-1* introns described above, as well as their fragments and variants, may be used as oligonucleotide primers or probes in order to detect the presence of a copy of the *TBC-1* gene in a test sample, or alternatively in order to amplify a target nucleotide sequence within the *TBC-1* intronic sequences.

Thus, the invention embodies purified, isolated, or recombinant polynucleotides comprising a nucleotide sequence selected from the group consisting of the 15 exons of the *TBC-1* gene which are described in the present invention, or a sequence complementary thereto. The invention also deals with purified, isolated, or recombinant nucleic acids comprising a combination of at least two exons of the *TBC-1* gene, wherein the polynucleotides are arranged within the nucleic acid, from the 5'-end to the 3'-end of said nucleic acid, in the same order as in SEQ ID Nos 1 and 2.

Thus, the invention embodies purified, isolated, or recombinant polynucleotides comprising a nucleotide sequence selected from the group consisting of the introns of the *TBC-1* gene, or a sequence complementary thereto.

The invention also encompasses a purified, isolated, or recombinant polynucleotide comprising a nucleotide sequence having at least 70, 75, 80, 85, 90, or 95% nucleotide identity with a sequence selected from the group consisting of SEQ ID Nos 1 and 2 or a complementary sequence thereto or a fragment thereof. The nucleotide differences as regards to the nucleotide sequence of SEQ ID Nos 1 or 2 may be generally randomly distributed throughout the entire nucleic acid.

Nevertheless, preferred nucleic acids are those wherein the nucleotide differences as regards to the nucleotide sequence of SEQ ID Nos 1 or 2 are predominantly located outside the coding sequences contained in the exons. These nucleic acids, as well as their fragments and variants, may be used as oligonucleotide primers or probes in order to detect the presence of a copy of the *TBC-1* gene in a
5 test sample, or alternatively in order to amplify a target nucleotide sequence within the *TBC-1* sequences.

Another object of the invention consists of a purified, isolated, or recombinant nucleic acid that hybridizes with a sequence selected from the group consisting of SEQ ID Nos 1 and 2 or a complementary sequence thereto or a variant thereof, under the stringent hybridization conditions as
10 defined above.

Particularly preferred nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of a nucleotide sequence selected from the group consisting of SEQ ID Nos 1 and 2, or the complements thereof. Additionally preferred
15 nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 1 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 1: 1-1000, 1001-2000, 2001-3000, 3001-4000, 4001-5000, 5001-6000, 6001-7000, 7001-8000, 8001-9000,
20 9001-10000, 10001-11000, 11001-12000, 12001-13000, 13001-14000, 14001-15000, 15001-16000, 16001-17000, and 17001-17590. Other preferred nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 2 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the
25 following nucleotide positions of SEQ ID No 2: 1-5000, 5001-10000, 10001-15000, 15001-20000, 20001-25000, 25001-30000, 30001-35000, 35001-40000, 40001-45000, 45001-50000, 50001-55000, 55001-60000, 60001-65000, 65001-70000, 70001-75000, 75001-80000, 80001-85000, 85001-90000, 90001-95000, and 95001-99960.

While this section is entitled "Genomic Sequences of *TBC-1*," it should be noted that
30 nucleic acid fragments of any size and sequence may also be comprised by the polynucleotides described in this section, flanking the genomic sequences of *TBC-1* on either side or between two or more such genomic sequences.

***TBC-1* cDNA Sequences**

The inventors have discovered that the expression of the *TBC-1* gene leads to the
35 production of at least two mRNA molecules, respectively a first and a second *TBC-1* transcription

product, as the results of alternative splicing events. They result from two distinct first exons, namely Exon 1 and Exon 1*bis*.

The first transcription product comprises Exons 1, 2, A, B, C, D, E, F, G, H, I, J, K, and L. This cDNA of SEQ ID No 3 includes a 5'-UTR region, spanning the whole Exon 1 and part of Exon 2. This 5'-UTR region starts from the nucleotide at position 1 and ends at the nucleotide at position 170 of the nucleotide sequence of SEQ ID No 3. The cDNA of SEQ ID No 3 includes a 3'-UTR region starting from the nucleotide at position 3726 and ending at the nucleotide at position 3983 of the nucleotide sequence of SEQ ID No 3. This first transcription product harbors a polyadenylation signal located between the nucleotide at position 3942 and the nucleotide at position 3947 of the nucleotide sequence of SEQ ID No 3.

The second *TBC-1* transcription product comprises Exons 1*bis*, 2, A, B, C, D, E, F, G, H, I, J, K, and L. This cDNA of SEQ ID No 4 includes a 5'-UTR region starting from the nucleotide at position 1 and ending at the nucleotide at position 175 of the nucleotide sequence of SEQ ID No 4. This second cDNA also includes a 3'-UTR region starting from the nucleotide at position 3731 and ending at the nucleotide at position 3988 of the nucleotide sequence of SEQ ID No 4. This second transcription product harbors a polyadenylation signal located between the nucleotide at position 3947 and the nucleotide at position 3952 of the nucleotide sequence of SEQ ID No 4.

The 5'-end sequence of this second *TBC-1* mRNA, more particularly the nucleotide sequence comprised between the nucleotide in position 1 and the nucleotide in position 458 of the nucleic acid of SEQ ID No 4 molecule corresponds to the nucleotide sequence of a 5'-EST that has been obtained from a human pancreas cDNA library and characterized following the teachings of the PCT Application No WO 96/34981. This 5'-EST is also part of the invention.

Another object of the invention consists of a purified or isolated nucleic acid comprising a polynucleotide selected from the group consisting of the nucleotide sequences of SEQ ID Nos 3 and 4 and to nucleic acid fragments thereof.

Preferred nucleic acid fragments of the nucleotide sequences of SEQ ID Nos 3 and 4 consist in polynucleotides comprising their respective Open Reading Frames encoding the TBC-1 protein.

Other preferred nucleic acid fragments of the nucleotide sequences of SEQ ID Nos 3 and 4 consist in polynucleotides comprising at least a part of their respective 5'-UTR or 3'-UTR regions.

The invention also pertains to a purified or isolated nucleic acid having at least a 95% of nucleotide identity with any one of the nucleotide sequences of SEQ ID Nos 3 and 4, or a fragment thereof.

Another object of the invention consists of purified, isolated or recombinant nucleic acids comprising a polynucleotide that hybridizes, under the stringent hybridization conditions defined herein, with any one of the nucleotide sequences of SEQ ID Nos 3 and 4, or a sequence complementary thereto or a fragment thereof.

The invention also relates to isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of a nucleotide sequence selected from the group consisting of SEQ ID Nos 3 and 4, or the complements thereof. Particularly preferred nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 3 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 3: 1-500, 501-1000, 1001-1500, 1501-2000, 2001-2500, 2501-3000, 3001-3500, and 3501-3983. Additionally preferred nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 4 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 4: 1-500, 501-1000, 1001-1500, 1501-2000, 2001-2500, 2501-3000, 3001-3500, and 3501-3988. Such a nucleic acid is notably useful as polynucleotide probe or primer specific for the *TBC-1* gene or the *TBC-1* mRNAs and cDNAs.

While this section is entitled "*TBC-1* cDNA Sequences," it should be noted that nucleic acid fragments of any size and sequence may also be comprised by the polynucleotides described in this section, flanking the genomic sequences of *TBC-1* on either side or between two or more such genomic sequences.

20

Coding Regions

The *TBC-1* open reading frame is contained in the two *TBC-1* mRNA molecules of about 4 kilobases isolated by the inventors.

More precisely, the effective *TBC-1* coding sequence is comprised between the nucleotide at position 171 and the nucleotide at position 3725 of SEQ ID No 3, and between the nucleotide at position 176 and the nucleotide at position 3730 of the nucleotide sequence of SEQ ID No 4.

The invention further provides a purified or isolated nucleic acid comprising a polynucleotide selected from the group consisting of a polynucleotide comprising a nucleic acid sequence located between the nucleotide at position 171 and the nucleotide at position 3725 of SEQ ID No 3, and a polynucleotide comprising a nucleic acid sequence located between the nucleotide at position 176 and the nucleotide at position 3730 of SEQ ID No 4 or a variant or fragment thereof or a sequence complementary thereto.

The present invention concerns a purified or isolated nucleic acid encoding a human TBC-1 protein, wherein said TBC-1 protein comprises an amino acid sequence of SEQ ID No 5, a nucleotide sequence complementary thereto, a fragment or a variant thereof. The present invention also embodies isolated, purified, and recombinant polynucleotides which encode a polypeptides comprising a contiguous span of at least 6 amino acids, preferably at least 8 or 10 amino acids,

more preferably at least 12, 15, 20, 25, 30, 40, 50, or 100 amino acids of SEQ ID No 5. In a preferred embodiment, the present invention embodies isolated, purified, and recombinant polynucleotides which encode a polypeptides comprising a contiguous span of at least 6 amino acids, preferably at least 8 or 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, or 100 amino acids of SEQ ID No 5 wherein said contiguous span includes at least 1, 2, 3, 5 or 10 of the following amino acid positions in SEQ ID No 5: 1-300, 301-600, 601-900, and 901-1168.

The above disclosed polynucleotide that contains only coding sequences derived from the *TBC-1* ORF may be expressed in a desired host cell or a desired host organism, when said polynucleotide is placed under the control of suitable expression signals. Such a polynucleotide, when placed under the suitable expression signals, may be inserted in a vector for its expression.

Regulatory Sequences Of *TBC-1*

The invention further deals with a purified or isolated nucleic acid comprising the nucleotide sequence of a regulatory region which is located either upstream of the first exon of the *TBC-1* gene and which is contained in the *TBC-1* genomic sequence of SEQ ID No 1, or downstream of the last exon of the *TBC-1* gene and which is contained in the *TBC-1* genomic sequence of SEQ ID No 2.

The 5'-regulatory sequence of the *TBC-1* gene is localized between the nucleotide in position 1 and the nucleotide in position 2000 of the nucleotide sequence of SEQ ID No 1. The 3'-regulatory sequence of the *TBC-1* gene is localized between nucleotide position 97961 and nucleotide position 99960 of SEQ ID No 2.

Polynucleotides derived from the 5' and 3' regulatory regions are useful in order to detect the presence of at least a copy of a nucleotide sequence of SEQ ID Nos 1 or 2 or a fragment thereof in a test sample.

The promoter activity of the 5' regulatory regions contained in *TBC-1* can be assessed as described below.

Genomic sequences lying upstream of the *TBC-1* Exons are cloned into a suitable promoter reporter vector, such as the pSEAP-Basic, pSEAP-Enhancer, p β gal-Basic, p β gal-Enhancer, or pEGFP-1 Promoter Reporter vectors available from Clontech. Briefly, each of these promoter reporter vectors include multiple cloning sites positioned upstream of a reporter gene encoding a readily assayable protein such as secreted alkaline phosphatase, beta galactosidase, or green fluorescent protein. The sequences upstream of the *TBC-1* coding region are inserted into the cloning sites upstream of the reporter gene in both orientations and introduced into an appropriate host cell. The level of reporter protein is assayed and compared to the level obtained from a vector which lacks an insert in the cloning site. The presence of an elevated expression level in the vector containing the insert with respect to the control vector indicates the presence of a promoter in the insert. If necessary, the upstream sequences can be cloned into vectors which contain an enhancer

for increasing transcription levels from weak promoter sequences. A significant level of expression above that observed with the vector lacking an insert indicates that a promoter sequence is present in the inserted upstream sequence.

Promoter sequences within the upstream genomic DNA may be further defined by
5 constructing nested deletions in the upstream DNA using conventional techniques such as Exonuclease III digestion. The resulting deletion fragments can be inserted into the promoter reporter vector to determine whether the deletion has reduced or obliterated promoter activity. In this way, the boundaries of the promoters may be defined. If desired, potential individual regulatory sites within the promoter may be identified using site directed mutagenesis or linker scanning to
10 obliterate potential transcription factor binding sites within the promoter, individually or in combination. The effects of these mutations on transcription levels may be determined by inserting the mutations into the cloning sites in the promoter reporter vectors.

Thus, the minimal size of the promoter of the *TBC-1* gene can be determined through the measurement of *TBC-1* expression levels. For this assay, an expression vector comprising
15 decreasing sizes from the promoter generally ranging from 2 kb to 100 bp, with a 3' end which is constant, operably linked to *TBC-1* coding sequence or to a reporter gene is used. Cells, which are preferably prostate cells and more preferably prostate cancer cells, are transfected with this vector and the expression level of the gene is assessed.

The strength and the specificity of the promoter of the *TBC-1* gene can be assessed through
20 the expression levels of the gene operably linked to this promoter in different types of cells and tissues. In one embodiment, the efficacy of the promoter of the *TBC-1* gene is assessed in normal and cancer cells. In a preferred embodiment, the efficacy of the promoter of the *TBC-1* gene is assessed in normal prostate cells and in prostate cancer cells which can present different degrees of malignancy.

25 Polynucleotides carrying the regulatory elements located both at the 5' end and at the 3' end of the *TBC-1* cDNAs may be advantageously used to control the transcriptional and translational activity of an heterologous polynucleotide of interest.

Thus, the present invention also concerns a purified or isolated nucleic acid comprising a polynucleotide which is selected from the group consisting of the 5' and 3' regulatory regions, or a
30 sequence complementary thereto or a biologically active fragment or variant thereof. "5' regulatory region" refers to the nucleotide sequence located between positions 1 and 2000 of SEQ ID No 1. "3' regulatory region" refers to the nucleotide sequence located between positions 97961 and 99960 of SEQ ID No 2.

The invention also pertains to a purified or isolated nucleic acid comprising a
35 polynucleotide having at least 95% nucleotide identity with a polynucleotide selected from the group consisting of the 5' and 3' regulatory regions, advantageously 99 % nucleotide identity, preferably 99.5% nucleotide identity and most preferably 99.8% nucleotide identity with a

polynucleotide selected from the group consisting of the 5' and 3' regulatory regions, or a sequence complementary thereto or a variant thereof or a biologically active fragment thereof.

Another object of the invention consists of purified, isolated or recombinant nucleic acids comprising a polynucleotide that hybridizes, under the stringent hybridization conditions defined
5 herein, with a polynucleotide selected from the group consisting of the nucleotide sequences of the 5'- and 3' regulatory regions, or a sequence complementary thereto or a variant thereof or a biologically active fragment thereof.

The 5'UTR and 3'UTR regions of a gene are of particular importance in that they often comprise regulatory elements which can play a role in providing appropriate expression levels,
10 particularly through the control of mRNA stability.

A 5' regulatory polynucleotide of the invention may include the 5'-UTR located between the nucleotide at position 1 and the nucleotide at position 170 of SEQ ID No 3, or a biologically active fragment or variant thereof.

Alternatively, a 5'-regulatory polynucleotide of the invention may include the 5'-UTR
15 located between the nucleotide at position 1 and the nucleotide at position 175 of SEQ ID No 4, or a biologically active fragment or variant thereof.

A 3' regulatory polynucleotide of the invention may include the 3'-UTR located between the nucleotide at position 3726 and the nucleotide at position 3983 of SEQ ID No 4, or a biologically active fragment or variant thereof.

20 Thus, the invention also pertains to a purified or isolated nucleic acid which is selected from the group consisting of :

- a) a nucleic acid comprising the nucleotide sequence of the 5' regulatory region;
- b) a nucleic acid comprising a biologically active fragment or variant of the nucleic acid of the 5' regulatory region.

25 Preferred fragments of the nucleic acid of the 5' regulatory region have a length of about 1000 nucleotides, more particularly of about 400 nucleotides, more preferably of about 200 nucleotides and most preferably about 100 nucleotides. More particularly, the invention further includes specific elements within this regulatory region, these elements preferably including the promoter region.

30 Preferred fragments of the 3' regulatory region are at least 50, 100, 150, 200, 300 or 400 bases in length.

By a "biologically active fragment or variant" of a *TBC-1* regulatory polynucleotide according to the present invention is intended a polynucleotide comprising or alternatively consisting in a fragment of said polynucleotide which is functional as a regulatory region for
35 expressing a recombinant polypeptide or a recombinant polynucleotide in a recombinant cell host.

For the purpose of the invention, a nucleic acid or polynucleotide is "functional" as a regulatory region for expressing a recombinant polypeptide or a recombinant polynucleotide if said

regulatory polynucleotide contains nucleotide sequences which contain transcriptional and translational regulatory information, and if such sequences are "operatively linked" to nucleotide sequences which encode the desired polypeptide or the desired polynucleotide. An operable linkage is a linkage in which the regulatory nucleic acid and the DNA sequence sought to be expressed are
5 linked in such a way as to permit gene expression.

In order, to identify the relevant biologically active polynucleotide derivatives of the 5' or 3' regulatory region, the one skill in the art will refer to the book of Sambrook et al. (Sambrook, 1989) in order to use a recombinant vector carrying a marker gene (i.e. beta galactosidase, chloramphenicol acetyl transferase, etc.) the expression of which will be detected when placed
10 under the control of a biologically active derivative polynucleotide of the 5' or 3' regulatory region.

Regulatory polynucleotides of the invention may be prepared from any of the nucleotide sequences of SEQ ID Nos 1 or 2 by cleavage using the suitable restriction enzymes, the one skill in the art being guided by the book of Sambrook et al. (1989). Regulatory polynucleotides may also be prepared by digestion of any of the nucleotide sequences of SEQ ID Nos 1 or 2 by an exonuclease
15 enzyme, such as Bal31 (Wabiko et al., 1986). These regulatory polynucleotides can also be prepared by chemical synthesis, as described elsewhere in the specification, when the synthesis of oligonucleotide probes or primers is disclosed.

The regulatory polynucleotides according to the invention may be advantageously part of a recombinant expression vector that may be used to express a coding sequence in a desired host cell
20 or host organism. The recombinant expression vectors according to the invention are described elsewhere in the specification.

The invention also encompasses a polynucleotide comprising :

- a) a nucleic acid comprising a regulatory nucleotide sequence of the 5' regulatory region, or a biologically active fragment or variant thereof;
- 25 b) a polynucleotide encoding a desired polypeptide or nucleic acid, operably linked to the nucleic acid comprising a regulatory nucleotide sequence of the 5' regulatory region, or its biologically active fragment or variant.
- c) Optionally, a nucleic acid comprising a 3' regulatory polynucleotide, preferably a 3'regulatory polynucleotide of the invention.

30 The desired polypeptide encoded by the above described nucleic acid may be of various nature or origin, encompassing proteins of prokaryotic or eukaryotic origin. Among the polypeptides expressed under the control of a *TBC-1* regulatory region, it may be cited bacterial, fungal or viral antigens. Are also encompassed eukaryotic proteins such as intracellular proteins, such as "house keeping" proteins, membrane-bound proteins, like receptors, and secreted proteins
35 like the numerous endogenous mediators such as cytokines.

The desired nucleic acid encoded by the above described polynucleotide, usually a RNA molecule, may be complementary to a *TBC-1* coding sequence and thus useful as an antisense polynucleotide.

Such a polynucleotide may be included in a recombinant expression vector in order to
5 express a desired polypeptide or a desired polynucleotide in host cell or in a host organism. Suitable recombinant vectors that contain a polynucleotide such as described hereinbefore are disclosed elsewhere in the specification.

TBC-1 Polypeptide And Peptide Fragments Thereof

It is now easy to produce proteins in high amounts by genetic engineering techniques
10 through expression vectors such as plasmids, phages or phagemids. The polynucleotide that code for one the polypeptides of the present invention is inserted in an appropriate expression vector in order to produce the polypeptide of interest *in vitro*.

Thus, the present invention also concerns a method for producing one of the polypeptides described herein, and especially a polypeptide of SEQ ID No 5 or a fragment or a variant thereof,
15 wherein said method comprises the steps of :

a) culturing, in an appropriate culture medium, a cell host previously transformed or transfected with the recombinant vector comprising a nucleic acid encoding a TBC-1 polypeptide, or a fragment or a variant thereof;

b) harvesting the culture medium thus conditioned or lyse the cell host, for example by
20 sonication or by an osmotic shock;

c) separating or purifying, from the said culture medium, or from the pellet of the resultant host cell lysate the thus produced polypeptide of interest.

d) Optionally characterizing the produced polypeptide of interest.

In a specific embodiment of the above method, step a) is preceded by a step wherein the
25 nucleic acid coding for a TBC-1 polypeptide, or a fragment or a variant thereof, is inserted in an appropriate vector, optionally after an appropriate cleavage of this amplified nucleic acid with one or several restriction endonucleases. The nucleic acid coding for a TBC-1 polypeptide or a fragment or a variant thereof may be the resulting product of an amplification reaction using a pair of primers according to the invention (by SDA, TAS, 3SR NASBA, TMA etc.).

30 The polypeptides according to the invention may be characterized by binding onto an immunoaffinity chromatography column on which polyclonal or monoclonal antibodies directed to a polypeptide of SEQ ID No 5, or a fragment or a variant thereof, have previously been immobilized.

Purification of the recombinant proteins or peptides according to the present invention may
35 be carried out by passage onto a Nickel or Copper affinity chromatography column. The Nickel chromatography column may contain the Ni-NTA resin (Porath et al., 1975).

The polypeptides or peptides thus obtained may be purified, for example by high performance liquid chromatography, such as reverse phase and/or cationic exchange HPLC, as described by Rougeot et al. (1994). The reason to prefer this kind of peptide or protein purification is the lack of byproducts found in the elution samples which renders the resultant purified protein or peptide more suitable for a therapeutic use.

Another object of the present invention consists in a purified or isolated TBC-1 polypeptide or a fragment or a variant thereof.

In a preferred embodiment, the TBC-1 polypeptide comprises an amino acid sequence of SEQ ID No 5 or a fragment or a variant thereof. The present invention also embodies isolated, purified, and recombinant polypeptides comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, 100, 150 or 200 amino acids of SEQ ID No 5. The present invention also embodies isolated, purified, and recombinant polypeptides comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, 100, 150 or 200 amino acids of SEQ ID No 5, wherein said contiguous span includes at least 1, 2, 3, 5 or 10 of the following amino acid positions: 1-200, 201-400, 401-600, 601-800, 801-1000, 1001-1168.

The invention also encompasses a purified, isolated, or recombinant polypeptides comprising an amino acid sequence having at least 90, 95, 98 or 99% amino acid identity with the amino acid sequence of SEQ ID No 5 or a fragment thereof.

The TBC-1 polypeptide of the invention possesses amino acid homologies as regards to the murine TBC-1 protein of 1141 amino acids in length which is described in US Patent No US 5,700,927. The TBC-1 protein of the invention also possesses some homologies with two other proteins : the Pollux drosophila protein (Zhang et al., 1996) and the CDC16 protein from *Caenorhabditis elegans* (Wilson et al., 1994). Figure 1 represents an amino acid alignment of a portion of the amino acid sequence of the TBC-1 protein of SEQ ID No 5 with other proteins sharing amino acid homology with TBC-1. The upper line shows the whole amino acid sequence of the murine tbc-1 protein described in US Patent No US 5,700,927; the second line represents part of the amino acid sequence of the TBC-1 protein of SEQ ID No 5; the third line (Genbank access No : *dmu50542*) depicts the amino acid sequence of the Pollux protein mentioned above; the fourth line (Genbank access No : *celf35h12*) shows the amino acid sequence of the *C. elegans* protein mentioned above; the fifth line presents positions in which consensus amino acids are identified, i.e. amino acids shared by the sequences presented in the four upper lines, when present.

The TBC-1 polypeptide of the amino acid sequence of SEQ ID No 5 has 1168 amino acids in length. The TBC-1 polypeptide includes a "TBC domain" which is spanning from the amino acid in position 786 to the amino acid in position 974 of the amino acid sequence of SEQ ID No 5. This TBC domain is represented in Figure 1 as a grey area spanning from the amino acid numbered 758 to the amino acid numbered 949. This TBC domain is likely to regulate protein-protein interactions.

Moreover, the TBC-1 TBC domain includes the amino acid sequence EVGYCQGL, spanning from the amino acid in position 886 to the amino acid in position 893 of the amino acid sequence of SEQ ID No 5. The EVGYCQGL amino acid sequence spans from the amino acid numbered 861 to the amino acid numbered 868 of Figure 1. This site may interact with a kinase. Based on the structural
5 similarity to *cdc16*, a yeast regulator of mitosis, TBC-1 is likely to regulate mitosis and cytokinesis by interacting with other proteins which also participate with the regulation of mitosis, cytokinesis and septum formation.

Preferred polypeptides of the invention comprise the TBC domain of TBC-1, or alternatively at least the EVGYCQGL amino acid sequence motif.

10 A further object of the present invention concerns a purified or isolated polypeptide which is encoded by a nucleic acid comprising a nucleotide sequence selected from the group consisting of SEQ ID Nos 1, 2, 3, and 4 or fragments or variants thereof.

A single variant molecule of the TBC-1 protein is explicitly excluded from the scope of the present invention, which is a polypeptide having the same amino acid sequence than the murine
15 *tbc1* protein described in the US Patent No 5,700,927.

Amino acid deletions, additions or substitutions in the TBC-1 protein are preferably located outside of the TBC domain as defined above. Most preferably, a mutated TBC-1 protein has an intact "EVGYCQGL" amino acid motif.

Such a mutated TBC-1 protein may be the target of diagnostic tools, such as specific
20 monoclonal or polyclonal antibodies, useful for detecting the mutated TBC-1 protein in a sample.

The invention also encompasses a TBC-1 polypeptide or a fragment or a variant thereof in which at least one peptide bound has been modified as described in the "Definitions" section.

Antibodies That Bind *TBC-1* Polypeptides of the Invention

Any TBC-1 polypeptide or whole protein may be used to generate antibodies capable of
25 specifically binding to an expressed TBC-1 protein or fragments thereof as described.

One antibody composition of the invention is capable of specifically binding or specifically bind to the variant of the TBC-1 protein of SEQ ID No 5. For an antibody composition to specifically bind to TBC-1, it must demonstrate at least a 5%, 10%, 15%, 20%, 25%, 50%, or 100% greater binding affinity for TBC-1 protein than for another protein in an ELISA, RIA, or other
30 antibody-based binding assay.

In a preferred embodiment, the invention concerns antibody compositions, either polyclonal or monoclonal, capable of selectively binding, or selectively bind to an epitope-containing a polypeptide comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, 100, 150 or 200 amino acids of SEQ
35 ID No 5; Optionally said epitope comprises at least 1, 2, 3, 5 or 10 of the following amino acid positions : 1-200, 201-400, 401-600, 601-800, 801-1000, 1001-1168.

The invention also concerns a purified or isolated antibody capable of specifically binding to a mutated TBC-1 protein or to a fragment or variant thereof comprising an epitope of the mutated TBC-1 protein. In another preferred embodiment, the present invention concerns an antibody capable of binding to a polypeptide comprising at least 10 consecutive amino acids of a TBC-1 protein and including at least one of the amino acids which can be encoded by the trait causing mutations.

In a preferred embodiment, the invention concerns the use in the manufacture of antibodies of a polypeptide comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, 100, 150 or 200 amino acids of SEQ ID No 5; Optionally said polypeptide comprises at least 1, 2, 3, 5 or 10 of the following amino acid positions : 1-200, 201-400, 401-600, 601-800, 801-1000, 1001-1168.

The antibodies of the invention may be labeled by any one of the radioactive, fluorescent or enzymatic labels known in the art.

The TBC-1 polypeptide of SEQ ID No 5 or a fragment thereof can be used for the preparation of polyclonal or monoclonal antibodies.

The TBC-1 polypeptide expressed from a DNA sequence comprising at least one of the nucleic acid sequences of SEQ ID Nos 1, 2, 3 and 4 may also be used to generate antibodies capable of specifically binding to the TBC-1 polypeptide of SEQ ID No 5 or a fragment thereof.

Preferred antibodies according to the invention are prepared using TBC-1 peptide fragments that do not comprise the EVGYCQGL amino acid motif.

Other preferred antibodies of the invention are prepared using TBC-1 peptide fragments that do not comprise the TBC domain defined elsewhere in the specification.

The antibodies may be prepared from hybridomas according to the technique described by Kohler and Milstein in 1975. The polyclonal antibodies may be prepared by immunization of a mammal, especially a mouse or a rabbit, with a polypeptide according to the invention that is combined with an adjuvant of immunity, and then by purifying of the specific antibodies contained in the serum of the immunized animal on a affinity chromatography column on which has previously been immobilized the polypeptide that has been used as the antigen.

The present invention also includes, chimeric single chain Fv antibody fragments (Martineau et al., 1998), antibody fragments obtained through phage display libraries (Ridder et al., 1995; Vaughan et al., 1995) and humanized antibodies (Reinmann et al., 1997; Leger et al., 1997).

Antibody preparations prepared according to either protocol are useful in quantitative immunoassays which determine concentrations of antigen-bearing substances in biological samples; they are also used semi-quantitatively or qualitatively to identify the presence of antigen in a biological sample. The antibodies may also be used in therapeutic compositions for killing cells expressing the protein or reducing the levels of the protein in the body.

Consequently, the invention is also directed to a method for detecting specifically the presence of a TBC-1 polypeptide according to the invention in a biological sample, said method comprising the following steps :

- a) bringing into contact the biological sample with a polyclonal or monoclonal antibody that specifically binds a TBC-1 polypeptide comprising an amino acid sequence of SEQ ID No 5, or to a peptide fragment or variant thereof; and
- b) detecting the antigen-antibody complex formed.

The invention also concerns a diagnostic kit for detecting *in vitro* the presence of a TBC-1 polypeptide according to the present invention in a biological sample, wherein said kit comprises:

- a) a polyclonal or monoclonal antibody that specifically binds a TBC-1 polypeptide comprising an amino acid sequence of SEQ ID No 5, or to a peptide fragment or variant thereof, optionally labeled;
- b) a reagent allowing the detection of the antigen-antibody complexes formed, said reagent carrying optionally a label, or being able to be recognized itself by a labeled reagent, more particularly in the case when the above-mentioned monoclonal or polyclonal antibody is not labeled by itself.

***TBC-1*-Related Biallelic Markers**

The inventors have discovered nucleotide polymorphisms located within the genomic DNA containing the *TBC-1* gene, and among them SNP that are also termed biallelic markers. The biallelic markers of the invention can be used for example for the generation of genetic map, the linkage analysis, the association studies.

A- Identification Of *TBC-1*-related Biallelic Markers

There are two preferred methods through which the biallelic markers of the present invention can be generated. In a first method, DNA samples from unrelated individuals are pooled together, following which the genomic DNA of interest is amplified and sequenced. The nucleotide sequences thus obtained are then analyzed to identify significant polymorphisms.

One of the major advantages of this method resides in the fact that the pooling of the DNA samples substantially reduces the number of DNA amplification reactions and sequencing which must be carried out. Moreover, this method is sufficiently sensitive so that a biallelic marker obtained therewith usually shows a sufficient degree of informativeness for conducting association studies.

In a second method for generating biallelic markers, the DNA samples are not pooled and are therefore amplified and sequenced individually. The resulting nucleotide sequences obtained are then also analyzed to identify significant polymorphisms.

It will readily be appreciated that when this second method is used, a substantially higher number of DNA amplification reactions must be carried out. It will further be appreciated that

including such potentially less informative biallelic markers in association studies to identify potential genetic associations with a trait may allow in some cases the direct identification of causal mutations, which may, depending on their penetrance, be rare mutations. This method is usually preferred when biallelic markers need to be identified in order to perform association studies within
5 candidate genes.

In both methods, the genomic DNA samples from which the biallelic markers of the present invention are generated are preferably obtained from unrelated individuals corresponding to a heterogeneous population of known ethnic background, or from familial cases.

The number of individuals from whom DNA samples are obtained can vary substantially,
10 preferably from about 10 to about 1000, preferably from about 50 to about 200 individuals. It is usually preferred to collect DNA samples from at least about 100 individuals in order to have sufficient polymorphic diversity in a given population to generate as many markers as possible and to generate statistically significant results.

As for the source of the genomic DNA to be subjected to analysis, any test sample can be
15 foreseen without any particular limitation. The preferred source of genomic DNA used in the context of the present invention is the peripheral venous blood of each donor.

The techniques of DNA extraction are well-known to the skilled technician. Details of a preferred embodiment are provided in Example 2.

DNA samples can be pooled or unpooled for the amplification step. DNA amplification
20 techniques are well-known to those skilled in the art.

Amplification techniques that can be used in the context of the present invention include, but are not limited to, the ligase chain reaction (LCR) described in EP-A- 320 308, WO 9320227 and EP-A-439 182, the polymerase chain reaction (PCR, RT-PCR) and techniques such as the nucleic acid sequence based amplification (NASBA) described in Guatelli J.C., et al.(1990) and in
25 Compton J.(1991), Q-beta amplification as described in European Patent Application No 4544610, strand displacement amplification as described in Walker et al.(1996) and EP A 684 315 and, target mediated amplification as described in PCT Publication WO 9322461.

LCR and Gap LCR are exponential amplification techniques, both depend on DNA ligase to join adjacent primers annealed to a DNA molecule. In Ligase Chain Reaction (LCR), probe pairs
30 are used which include two primary (first and second) and two secondary (third and fourth) probes, all of which are employed in molar excess to target. The first probe hybridizes to a first segment of the target strand and the second probe hybridizes to a second segment of the target strand, the first and second segments being contiguous so that the primary probes abut one another in 5' phosphate-3'hydroxyl relationship, and so that a ligase can covalently fuse or ligate the two probes into a fused
35 product. In addition, a third (secondary) probe can hybridize to a portion of the first probe and a fourth (secondary) probe can hybridize to a portion of the second probe in a similar abutting fashion. Of course, if the target is initially double stranded, the secondary probes also will

hybridize to the target complement in the first instance. Once the ligated strand of primary probes is separated from the target strand, it will hybridize with the third and fourth probes, which can be ligated to form a complementary, secondary ligated product. It is important to realize that the ligated products are functionally equivalent to either the target or its complement. By repeated
5 cycles of hybridization and ligation, amplification of the target sequence is achieved. A method for multiplex LCR has also been described (WO 9320227). Gap LCR (GLCR) is a version of LCR where the probes are not adjacent but are separated by 2 to 3 bases.

For amplification of mRNAs, it is within the scope of the present invention to reverse transcribe mRNA into cDNA followed by polymerase chain reaction (RT-PCR); or, to use a single
10 enzyme for both steps as described in U.S. Patent No. 5,322,770 or, to use Asymmetric Gap LCR (RT-AGLCR) as described by Marshall et al.(1994). AGLCR is a modification of GLCR that allows the amplification of RNA.

The PCR technology is the preferred amplification technique used in the present invention. A variety of PCR techniques are familiar to those skilled in the art. For a review of PCR
15 technology, see White (1997) and the publication entitled "PCR Methods and Applications" (1991, Cold Spring Harbor Laboratory Press). In each of these PCR procedures, PCR primers on either side of the nucleic acid sequences to be amplified are added to a suitably prepared nucleic acid sample along with dNTPs and a thermostable polymerase such as Taq polymerase, Pfu polymerase, or Vent polymerase. The nucleic acid in the sample is denatured and the PCR primers are
20 specifically hybridized to complementary nucleic acid sequences in the sample. The hybridized primers are extended. Thereafter, another cycle of denaturation, hybridization, and extension is initiated. The cycles are repeated multiple times to produce an amplified fragment containing the nucleic acid sequence between the primer sites. PCR has further been described in several patents including US Patents 4,683,195; 4,683,202; and 4,965,188.

25 The PCR technology is the preferred amplification technique used to identify new biallelic markers. A typical example of a PCR reaction suitable for the purposes of the present invention is provided in Example 3.

One of the aspects of the present invention is a method for the amplification of a *TBC-1* gene, particularly the genomic sequences of SEQ ID Nos 1 and 2 or of the cDNA sequence of SEQ
30 ID Nos 3 or 4 or a fragment or variant thereof in a test sample, preferably using the PCR technology. The method comprises the steps of contacting a test sample suspected of containing the target *TBC-1* sequence or portion thereof with amplification reaction reagents comprising a pair of amplification primers.

Thus, the present invention also relates to a method for the amplification of a *TBC-1* gene
35 sequence, particularly of a fragment of the genomic sequence of SEQ ID No 1 or of the cDNA sequence of SEQ ID No 2 or 3, or a fragment or a variant thereof in a test sample, said method comprising the steps of :

a) contacting a test sample suspected of containing the targeted *TBC-1* gene sequence or portion thereof with amplification reaction reagents comprising a pair of amplification primers located on either side of the *TBC-1* region to be amplified, and

b) optionally, detecting the amplification products.

5 The invention also concerns a kit for the amplification of a *TBC-1* gene sequence, particularly of a portion of the genomic sequence of SEQ ID Nos 1 or 2, or of the cDNA sequence of SEQ ID Nos 3 or 4, or a variant thereof in a test sample, wherein said kit comprises:

a) a pair of oligonucleotide primers located on either side of the *TBC-1* region to be amplified;

10 b) optionally, the reagents necessary for performing the amplification reaction.

In one embodiment of the above amplification method and kit, the amplification product is detected by hybridization with a labeled probe having a sequence which is complementary to the amplified region. In another embodiment of the above amplification method and kit, primers comprise a sequence which is selected from the group consisting of B1 to B15, C1 to C15, D1 to D19, and E1 to E19.

In a first embodiment of the present invention, biallelic markers are identified using genomic sequence information generated by the inventors. Sequenced genomic DNA fragments are used to design primers for the amplification of 500 bp fragments. These 500 bp fragments are amplified from genomic DNA and are scanned for biallelic markers. Primers may be designed using the OSP software (Hillier L. and Green P., 1991). All primers may contain, upstream of the specific target bases, a common oligonucleotide tail that serves as a sequencing primer. Those skilled in the art are familiar with primer extensions, which can be used for these purposes.

Preferred primers, useful for the amplification of genomic sequences encoding the candidate genes, focus on promoters, exons and splice sites of the genes. A biallelic marker presents a higher probability to be an eventual causal mutation if it is located in these functional regions of the gene. Preferred amplification primers of the invention include the nucleotide sequences of B1 to B15 and C1 to C15 further detailed in Example 3.

The amplification products generated as described above with the primers of the invention are then sequenced using methods known and available to the skilled technician. Preferably, the amplified DNA is subjected to automated dideoxy terminator sequencing reactions using a dye-primer cycle sequencing protocol. Following gel image analysis and DNA sequence extraction, sequence data are automatically processed with adequate software to assess sequence quality.

A polymorphism analysis software is used that detects the presence of biallelic sites among individual or pooled amplified fragment sequences. Polymorphism search is based on the presence of superimposed peaks in the electrophoresis pattern. These peaks which present distinct colors correspond to two different nucleotides at the same position on the sequence. The polymorphism has to be detected on both strands for validation.

19 biallelic markers were found in the *TBC-1* gene. They are detailed in the Table 2. They are located in intronic regions.

B- Genotyping Of *TBC-1*-Related Biallelic Markers

The polymorphisms identified above can be further confirmed and their respective frequencies can be determined through various methods using the previously described primers and probes. These methods can also be useful for genotyping either new populations in association studies or linkage analysis or individuals in the context of detection of alleles of biallelic markers which are known to be associated with a given trait. The genotyping of the biallelic markers is also important for the mapping. Those skilled in the art should note that the methods described below can be equally performed on individual or pooled DNA samples.

Once a given polymorphic site has been found and characterized as a biallelic marker as described above, several methods can be used in order to determine the specific allele carried by an individual at the given polymorphic base.

The identification of biallelic markers described previously allows the design of appropriate oligonucleotides, which can be used as probes and primers, to amplify a *TBC-1* gene containing the polymorphic site of interest and for the detection of such polymorphisms.

The biallelic markers according to the present invention may be used in methods for the identification and characterization of an association between alleles for one or several biallelic markers of the sequence of the *TBC-1* gene and a trait.

The identified polymorphisms, and consequently the biallelic markers of the invention, may be used in methods for the detection in an individual of *TBC-1* alleles associated with a trait, more particularly a trait related to a cell differentiation or abnormal cell proliferation disorders, and most particularly a trait related to cancer diseases, specifically prostate cancer.

In one embodiment the invention encompasses methods of genotyping comprising determining the identity of a nucleotide at a *TBC-1*-related biallelic marker or the complement thereof in a biological sample; optionally, wherein said *TBC-1*-related biallelic marker is selected from the group consisting of A1 to A19, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, wherein said biological sample is derived from a single subject; optionally, wherein the identity of the nucleotides at said biallelic marker is determined for both copies of said biallelic marker present in said individual's genome; optionally, wherein said biological sample is derived from multiple subjects; Optionally, the genotyping methods of the invention encompass methods with any further limitation described in this disclosure, or those following, specified alone or in any combination; Optionally, said method is performed *in vitro*; optionally, further comprising amplifying a portion of said sequence comprising the biallelic marker prior to said determining step; Optionally, wherein said amplifying is performed by PCR, LCR, or replication of a recombinant vector comprising an origin of replication and said fragment in a host cell; optionally, wherein said determining is performed by a

hybridization assay, a sequencing assay, a microsequencing assay, or an enzyme-based mismatch detection assay.

Source of Nucleic Acids for genotyping

Any source of nucleic acids, in purified or non-purified form, can be utilized as the starting nucleic acid, provided it contains or is suspected of containing the specific nucleic acid sequence desired. DNA or RNA may be extracted from cells, tissues, body fluids and the like as described above. While nucleic acids for use in the genotyping methods of the invention can be derived from any mammalian source, the test subjects and individuals from which nucleic acid samples are taken are generally understood to be human.

10 Amplification Of DNA Fragments Comprising Biallelic Markers

Methods and polynucleotides are provided to amplify a segment of nucleotides comprising one or more biallelic marker of the present invention. It will be appreciated that amplification of DNA fragments comprising biallelic markers may be used in various methods and for various purposes and is not restricted to genotyping. Nevertheless, many genotyping methods, although not all, require the previous amplification of the DNA region carrying the biallelic marker of interest. Such methods specifically increase the concentration or total number of sequences that span the biallelic marker or include that site and sequences located either distal or proximal to it. Diagnostic assays may also rely on amplification of DNA segments carrying a biallelic marker of the present invention. Amplification of DNA may be achieved by any method known in the art. Amplification techniques are described above in the section entitled, "Identification of *TBC-1*-related biallelic markers."

Some of these amplification methods are particularly suited for the detection of single nucleotide polymorphisms and allow the simultaneous amplification of a target sequence and the identification of the polymorphic nucleotide as it is further described below.

25 The identification of biallelic markers as described above allows the design of appropriate oligonucleotides, which can be used as primers to amplify DNA fragments comprising the biallelic markers of the present invention. Amplification can be performed using the primers initially used to discover new biallelic markers which are described herein or any set of primers allowing the amplification of a DNA fragment comprising a biallelic marker of the present invention.

30 In some embodiments the present invention provides primers for amplifying a DNA fragment containing one or more biallelic markers of the present invention. Preferred amplification primers are listed in Example 2. It will be appreciated that the primers listed are merely exemplary and that any other set of primers which produce amplification products containing one or more biallelic markers of the present invention are also of use.

35 The spacing of the primers determines the length of the segment to be amplified. In the context of the present invention, amplified segments carrying biallelic markers can range in size from at least about 25 bp to 35 kbp. Amplification fragments from 25-3000 bp are typical,

fragments from 50-1000 bp are preferred and fragments from 100-600 bp are highly preferred. It will be appreciated that amplification primers for the biallelic markers may be any sequence which allow the specific amplification of any DNA fragment carrying the markers. Amplification primers may be labeled or immobilized on a solid support as described in "Oligonucleotide probes and
5 primers".

Methods of Genotyping DNA samples for Biallelic Markers

Any method known in the art can be used to identify the nucleotide present at a biallelic marker site. Since the biallelic marker allele to be detected has been identified and specified in the present invention, detection will prove simple for one of ordinary skill in the art by employing any
10 of a number of techniques. Many genotyping methods require the previous amplification of the DNA region carrying the biallelic marker of interest. While the amplification of target or signal is often preferred at present, ultrasensitive detection methods which do not require amplification are also encompassed by the present genotyping methods. Methods well-known to those skilled in the art that can be used to detect biallelic polymorphisms include methods such as, conventional dot
15 blot analyzes, single strand conformational polymorphism analysis (SSCP) described by Orita et al.(1989), denaturing gradient gel electrophoresis (DGGE), heteroduplex analysis, mismatch cleavage detection, and other conventional techniques as described in Sheffield et al.(1991), White et al.(1992), Grompe et al.(1989 and 1993). Another method for determining the identity of the nucleotide present at a particular polymorphic site employs a specialized exonuclease-resistant
20 nucleotide derivative as described in US patent 4,656,127.

Preferred methods involve directly determining the identity of the nucleotide present at a biallelic marker site by sequencing assay, enzyme-based mismatch detection assay, or hybridization assay. The following is a description of some preferred methods. A highly preferred method is the microsequencing technique. The term "sequencing" is generally used herein to refer to polymerase
25 extension of duplex primer/template complexes and includes both traditional sequencing and microsequencing.

1) Sequencing Assays

The nucleotide present at a polymorphic site can be determined by sequencing methods. In a preferred embodiment, DNA samples are subjected to PCR amplification before sequencing as
30 described above. DNA sequencing methods are described in "Sequencing Of Amplified Genomic DNA And Identification Of Single Nucleotide Polymorphisms".

Preferably, the amplified DNA is subjected to automated dideoxy terminator sequencing reactions using a dye-primer cycle sequencing protocol. Sequence analysis allows the identification of the base present at the biallelic marker site.

2) Microsequencing Assays

In microsequencing methods, the nucleotide at a polymorphic site in a target DNA is detected by a single nucleotide primer extension reaction. This method involves appropriate microsequencing primers which, hybridize just upstream of the polymorphic base of interest in the target nucleic acid. A polymerase is used to specifically extend the 3' end of the primer with one single ddNTP (chain terminator) complementary to the nucleotide at the polymorphic site. Next the identity of the incorporated nucleotide is determined in any suitable way.

Typically, microsequencing reactions are carried out using fluorescent ddNTPs and the extended microsequencing primers are analyzed by electrophoresis on ABI 377 sequencing machines to determine the identity of the incorporated nucleotide as described in EP 412 883, the disclosure of which is incorporated herein by reference in its entirety. Alternatively capillary electrophoresis can be used in order to process a higher number of assays simultaneously. An example of a typical microsequencing procedure that can be used in the context of the present invention is provided in Example 4.

Different approaches can be used for the labeling and detection of ddNTPs. A homogeneous phase detection method based on fluorescence resonance energy transfer has been described by Chen and Kwok (1997) and Chen et al.(1997). In this method, amplified genomic DNA fragments containing polymorphic sites are incubated with a 5'-fluorescein-labeled primer in the presence of allelic dye-labeled dideoxyribonucleoside triphosphates and a modified Taq polymerase. The dye-labeled primer is extended one base by the dye-terminator specific for the allele present on the template. At the end of the genotyping reaction, the fluorescence intensities of the two dyes in the reaction mixture are analyzed directly without separation or purification. All these steps can be performed in the same tube and the fluorescence changes can be monitored in real time. Alternatively, the extended primer may be analyzed by MALDI-TOF Mass Spectrometry. The base at the polymorphic site is identified by the mass added onto the microsequencing primer (see Haff and Smirnov, 1997).

Microsequencing may be achieved by the established microsequencing method or by developments or derivatives thereof. Alternative methods include several solid-phase microsequencing techniques. The basic microsequencing protocol is the same as described previously, except that the method is conducted as a heterogeneous phase assay, in which the primer or the target molecule is immobilized or captured onto a solid support. To simplify the primer separation and the terminal nucleotide addition analysis, oligonucleotides are attached to solid supports or are modified in such ways that permit affinity separation as well as polymerase extension. The 5' ends and internal nucleotides of synthetic oligonucleotides can be modified in a number of different ways to permit different affinity separation approaches, e.g., biotinylation. If a single affinity group is used on the oligonucleotides, the oligonucleotides can be separated from the incorporated terminator reagent. This eliminates the need of physical or size separation. More than one oligonucleotide can be separated from the terminator reagent and analyzed simultaneously if

more than one affinity group is used. This permits the analysis of several nucleic acid species or more nucleic acid sequence information per extension reaction. The affinity group need not be on the priming oligonucleotide but could alternatively be present on the template. For example, immobilization can be carried out via an interaction between biotinylated DNA and streptavidin-coated microtitration wells or avidin-coated polystyrene particles. In the same manner, oligonucleotides or templates may be attached to a solid support in a high-density format. In such solid phase microsequencing reactions, incorporated ddNTPs can be radiolabeled (Syvänen, 1994) or linked to fluorescein (Livak and Hainer, 1994). The detection of radiolabeled ddNTPs can be achieved through scintillation-based techniques. The detection of fluorescein-linked ddNTPs can be based on the binding of anti fluorescein antibody conjugated with alkaline phosphatase, followed by incubation with a chromogenic substrate (such as *p*-nitrophenyl phosphate). Other possible reporter-detection pairs include: ddNTP linked to dinitrophenyl (DNP) and anti-DNP alkaline phosphatase conjugate (Harju et al., 1993) or biotinylated ddNTP and horseradish peroxidase-conjugated streptavidin with *o*-phenylenediamine as a substrate (WO 92/15712). As yet another alternative solid-phase microsequencing procedure, Nyren et al. (1993) described a method relying on the detection of DNA polymerase activity by an enzymatic luminometric inorganic pyrophosphate detection assay (ELIDA).

Pastinen et al. (1997) describe a method for multiplex detection of single nucleotide polymorphism in which the solid phase minisequencing principle is applied to an oligonucleotide array format. High-density arrays of DNA probes attached to a solid support (DNA chips) are further described below.

In one aspect the present invention provides polynucleotides and methods to genotype one or more biallelic markers of the present invention by performing a microsequencing assay. Preferred microsequencing primers include the nucleotide sequences D1 to D15 and E1 to E15. It will be appreciated that the microsequencing primers listed in Example 5 are merely exemplary and that, any primer having a 3' end immediately adjacent to the polymorphic nucleotide may be used. Similarly, it will be appreciated that microsequencing analysis may be performed for any biallelic marker or any combination of biallelic markers of the present invention. One aspect of the present invention is a solid support which includes one or more microsequencing primers listed in Example 5, or fragments comprising at least 8, 12, 15, 20, 25, 30, 40, or 50 consecutive nucleotides thereof, to the extent that such lengths are consistent with the primer described, and having a 3' terminus immediately upstream of the corresponding biallelic marker, for determining the identity of a nucleotide at a biallelic marker site.

3) Mismatch detection assays based on polymerases and ligases

In one aspect the present invention provides polynucleotides and methods to determine the allele of one or more biallelic markers of the present invention in a biological sample, by mismatch

detection assays based on polymerases and/or ligases. These assays are based on the specificity of polymerases and ligases. Polymerization reactions places particularly stringent requirements on correct base pairing of the 3' end of the amplification primer and the joining of two oligonucleotides hybridized to a target DNA sequence is quite sensitive to mismatches close to the ligation site, especially at the 3' end. Methods, primers and various parameters to amplify DNA fragments comprising biallelic markers of the present invention are further described above in "Amplification Of DNA Fragments Comprising Biallelic Markers".

Allele Specific Amplification Primers

Discrimination between the two alleles of a biallelic marker can also be achieved by allele specific amplification, a selective strategy, whereby one of the alleles is amplified without amplification of the other allele. For allele specific amplification, at least one member of the pair of primers is sufficiently complementary with a region of a *TBC-1* gene comprising the polymorphic base of a biallelic marker of the present invention to hybridize therewith and to initiate the amplification. Such primers are able to discriminate between the two alleles of a biallelic marker.

This is accomplished by placing the polymorphic base at the 3' end of one of the amplification primers. Because the extension forms from the 3' end of the primer, a mismatch at or near this position has an inhibitory effect on amplification. Therefore, under appropriate amplification conditions, these primers only direct amplification on their complementary allele. Determining the precise location of the mismatch and the corresponding assay conditions are well within the ordinary skill in the art.

Ligation/Amplification Based Methods

The "Oligonucleotide Ligation Assay" (OLA) uses two oligonucleotides which are designed to be capable of hybridizing to abutting sequences of a single strand of a target molecules. One of the oligonucleotides is biotinylated, and the other is detectably labeled. If the precise complementary sequence is found in a target molecule, the oligonucleotides will hybridize such that their termini abut, and create a ligation substrate that can be captured and detected. OLA is capable of detecting single nucleotide polymorphisms and may be advantageously combined with PCR as described by Nickerson et al.(1990). In this method, PCR is used to achieve the exponential amplification of target DNA, which is then detected using OLA.

Other amplification methods which are particularly suited for the detection of single nucleotide polymorphism include LCR (ligase chain reaction), Gap LCR (GLCR) which are described above in "DNA Amplification". LCR uses two pairs of probes to exponentially amplify a specific target. The sequences of each pair of oligonucleotides, is selected to permit the pair to hybridize to abutting sequences of the same strand of the target. Such hybridization forms a substrate for a template-dependant ligase. In accordance with the present invention, LCR can be performed with oligonucleotides having the proximal and distal sequences of the same strand of a biallelic marker site. In one embodiment, either oligonucleotide will be designed to include the

biallelic marker site. In such an embodiment, the reaction conditions are selected such that the oligonucleotides can be ligated together only if the target molecule either contains or lacks the specific nucleotide that is complementary to the biallelic marker on the oligonucleotide. In an alternative embodiment, the oligonucleotides will not include the biallelic marker, such that when they hybridize to the target molecule, a "gap" is created as described in WO 90/01069. This gap is then "filled" with complementary dNTPs (as mediated by DNA polymerase), or by an additional pair of oligonucleotides. Thus at the end of each cycle, each single strand has a complement capable of serving as a target during the next cycle and exponential allele-specific amplification of the desired sequence is obtained.

10 Ligase/Polymerase-mediated Genetic Bit AnalysisTM is another method for determining the identity of a nucleotide at a preselected site in a nucleic acid molecule (WO 95/21271). This method involves the incorporation of a nucleoside triphosphate that is complementary to the nucleotide present at the preselected site onto the terminus of a primer molecule, and their subsequent ligation to a second oligonucleotide. The reaction is monitored by detecting a specific
15 label attached to the reaction's solid phase or by detection in solution.

4) Hybridization Assay Methods

A preferred method of determining the identity of the nucleotide present at a biallelic marker site involves nucleic acid hybridization. The hybridization probes, which can be conveniently used in such reactions, preferably include the probes defined herein. Any
20 hybridization assay may be used including Southern hybridization, Northern hybridization, dot blot hybridization and solid-phase hybridization (see Sambrook et al., 1989).

Hybridization refers to the formation of a duplex structure by two single stranded nucleic acids due to complementary base pairing. Hybridization can occur between exactly complementary nucleic acid strands or between nucleic acid strands that contain minor regions of mismatch.
25 Specific probes can be designed that hybridize to one form of a biallelic marker and not to the other and therefore are able to discriminate between different allelic forms. Allele-specific probes are often used in pairs, one member of a pair showing perfect match to a target sequence containing the original allele and the other showing a perfect match to the target sequence containing the alternative allele. Hybridization conditions should be sufficiently stringent that there is a significant
30 difference in hybridization intensity between alleles, and preferably an essentially binary response, whereby a probe hybridizes to only one of the alleles. Stringent, sequence specific hybridization conditions, under which a probe will hybridize only to the exactly complementary target sequence are well known in the art (Sambrook et al., 1989). Stringent conditions are sequence dependent and will be different in different circumstances. Generally, stringent conditions are selected to be about
35 5°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. Although such hybridization can be performed in solution, it is preferred to employ a

solid-phase hybridization assay. The target DNA comprising a biallelic marker of the present invention may be amplified prior to the hybridization reaction. The presence of a specific allele in the sample is determined by detecting the presence or the absence of stable hybrid duplexes formed between the probe and the target DNA. The detection of hybrid duplexes can be carried out by a number of methods. Various detection assay formats are well known which utilize detectable labels bound to either the target or the probe to enable detection of the hybrid duplexes. Typically, hybridization duplexes are separated from unhybridized nucleic acids and the labels bound to the duplexes are then detected. Those skilled in the art will recognize that wash steps may be employed to wash away excess target DNA or probe as well as unbound conjugate. Further, standard heterogeneous assay formats are suitable for detecting the hybrids using the labels present on the primers and probes.

Two recently developed assays allow hybridization-based allele discrimination with no need for separations or washes (see Landegren U. et al., 1998). The TaqMan assay takes advantage of the 5' nuclease activity of Taq DNA polymerase to digest a DNA probe annealed specifically to the accumulating amplification product. TaqMan probes are labeled with a donor-acceptor dye pair that interacts via fluorescence energy transfer. Cleavage of the TaqMan probe by the advancing polymerase during amplification dissociates the donor dye from the quenching acceptor dye, greatly increasing the donor fluorescence. All reagents necessary to detect two allelic variants can be assembled at the beginning of the reaction and the results are monitored in real time (see Livak et al., 1995). In an alternative homogeneous hybridization based procedure, molecular beacons are used for allele discriminations. Molecular beacons are hairpin-shaped oligonucleotide probes that report the presence of specific nucleic acids in homogeneous solutions. When they bind to their targets they undergo a conformational reorganization that restores the fluorescence of an internally quenched fluorophore (Tyagi et al., 1998).

The polynucleotides provided herein can be used to produce probes which can be used in hybridization assays for the detection of biallelic marker alleles in biological samples. These probes are characterized in that they preferably comprise between 8 and 50 nucleotides, and in that they are sufficiently complementary to a sequence comprising a biallelic marker of the present invention to hybridize thereto and preferably sufficiently specific to be able to discriminate the targeted sequence for only one nucleotide variation. A particularly preferred probe is 25 nucleotides in length. Preferably the biallelic marker is within 4 nucleotides of the center of the polynucleotide probe. In particularly preferred probes, the biallelic marker is at the center of said polynucleotide. Preferred probes comprise a nucleotide sequence selected from the group consisting of amplicons listed in Table 1 and the sequences complementary thereto, or a fragment thereof, said fragment comprising at least about 8 consecutive nucleotides, preferably 10, 15, 20, more preferably 25, 30, 40, 47, or 50 consecutive nucleotides and containing a polymorphic base. Preferred probes comprise a nucleotide sequence selected from the group consisting of P1 to P7, P9 to P13, P15 to

P19 and the sequences complementary thereto. In preferred embodiments the polymorphic base(s) are within 5, 4, 3, 2, 1, nucleotides of the center of the said polynucleotide, more preferably at the center of said polynucleotide.

Preferably the probes of the present invention are labeled or immobilized on a solid support.

- 5 Labels and solid supports are further described in "Oligonucleotide Probes and Primers". The probes can be non-extendable as described in "Oligonucleotide Probes and Primers".

By assaying the hybridization to an allele specific probe, one can detect the presence or absence of a biallelic marker allele in a given sample. High-Throughput parallel hybridization in array format is specifically encompassed within "hybridization assays" and are described below.

10 **5) Hybridization To Addressable Arrays Of Oligonucleotides**

Hybridization assays based on oligonucleotide arrays rely on the differences in hybridization stability of short oligonucleotides to perfectly matched and mismatched target sequence variants. Efficient access to polymorphism information is obtained through a basic structure comprising high-density arrays of oligonucleotide probes attached to a solid support (e.g.,
15 the chip) at selected positions. Each DNA chip can contain thousands to millions of individual synthetic DNA probes arranged in a grid-like pattern and miniaturized to the size of a dime.

The chip technology has already been applied with success in numerous cases. For example, the screening of mutations has been undertaken in the BRCA1 gene, in *S. cerevisiae* mutant strains, and in the protease gene of HIV-1 virus (Hacia et al., 1996; Shoemaker et al., 1996;
20 Kozal et al., 1996). Chips of various formats for use in detecting biallelic polymorphisms can be produced on a customized basis by Affymetrix (GeneChip™), Hyseq (HyChip and HyGnostics), and Protogene Laboratories.

In general, these methods employ arrays of oligonucleotide probes that are complementary to target nucleic acid sequence segments from an individual which, target sequences include a
25 polymorphic marker. EP 785280 describes a tiling strategy for the detection of single nucleotide polymorphisms. Briefly, arrays may generally be "tiled" for a large number of specific polymorphisms. By "tiling" is generally meant the synthesis of a defined set of oligonucleotide probes which is made up of a sequence complementary to the target sequence of interest, as well as preselected variations of that sequence, e.g., substitution of one or more given positions with one or
30 more members of the basis set of nucleotides. Tiling strategies are further described in PCT application No. WO 95/11995. In a particular aspect, arrays are tiled for a number of specific, identified biallelic marker sequences. In particular, the array is tiled to include a number of detection blocks, each detection block being specific for a specific biallelic marker or a set of biallelic markers. For example, a detection block may be tiled to include a number of probes, which
35 span the sequence segment that includes a specific polymorphism. To ensure probes that are complementary to each allele, the probes are synthesized in pairs differing at the biallelic marker.

In addition to the probes differing at the polymorphic base, monosubstituted probes are also generally tiled within the detection block. These monosubstituted probes have bases at and up to a certain number of bases in either direction from the polymorphism, substituted with the remaining nucleotides (selected from A, T, G, C and U). Typically the probes in a tiled detection block will include substitutions of the sequence positions up to and including those that are 5 bases away from the biallelic marker. The monosubstituted probes provide internal controls for the tiled array, to distinguish actual hybridization from artefactual cross-hybridization. Upon completion of hybridization with the target sequence and washing of the array, the array is scanned to determine the position on the array to which the target sequence hybridizes. The hybridization data from the scanned array is then analyzed to identify which allele or alleles of the biallelic marker are present in the sample. Hybridization and scanning may be carried out as described in PCT application No. WO 92/10092 and WO 95/11995 and US patent No. 5,424,186.

Thus, in some embodiments, the chips may comprise an array of nucleic acid sequences of fragments of about 15 nucleotides in length. In further embodiments, the chip may comprise an array including at least one of the sequences selected from the group consisting of amplicons listed in table 1 and the sequences complementary thereto, or a fragment thereof, said fragment comprising at least about 8 consecutive nucleotides, preferably 10, 15, 20, more preferably 25, 30, 40, 47, or 50 consecutive nucleotides and containing a polymorphic base. In preferred embodiments the polymorphic base is within 5, 4, 3, 2, 1, nucleotides of the center of the said polynucleotide, more preferably at the center of said polynucleotide. In some embodiments, the chip may comprise an array of at least 2, 3, 4, 5, 6, 7, 8 or more of these polynucleotides of the invention. Solid supports and polynucleotides of the present invention attached to solid supports are further described in "Oligonucleotide Probes And Primers".

6) Integrated Systems

Another technique, which may be used to analyze polymorphisms, includes multicomponent integrated systems, which miniaturize and compartmentalize processes such as PCR and capillary electrophoresis reactions in a single functional device. An example of such technique is disclosed in US patent 5,589,136, which describes the integration of PCR amplification and capillary electrophoresis in chips.

Integrated systems can be envisaged mainly when microfluidic systems are used. These systems comprise a pattern of microchannels designed onto a glass, silicon, quartz, or plastic wafer included on a microchip. The movements of the samples are controlled by electric, electroosmotic or hydrostatic forces applied across different areas of the microchip to create functional microscopic valves and pumps with no moving parts.

For genotyping biallelic markers, the microfluidic system may integrate nucleic acid amplification, microsequencing, capillary electrophoresis and a detection method such as laser-induced fluorescence detection.

Association Studies With The Biallelic Markers Of The *TBC-1* Gene

5 The identification of genes involved in suspected heterogeneous, polygenic and multifactorial traits such as cancer can be carried out through two main strategies currently used for genetic mapping: linkage analysis and association studies. Association studies examine the frequency of marker alleles in unrelated trait positive (T+) individuals compared with trait negative (T-) controls, and are generally employed in the detection of polygenic inheritance. Association
10 studies as a method of mapping genetic traits rely on the phenomenon of linkage disequilibrium.

If two genetic loci lie on the same chromosome, then sets of alleles of these loci on the same chromosomal segment (called haplotypes) tend to be transmitted as a block from generation to generation. When not broken up by recombination, haplotypes can be tracked not only through pedigrees but also through populations. The resulting phenomenon at the population level is that the
15 occurrence of pairs of specific alleles at different loci on the same chromosome is not random, and the deviation from random is called linkage disequilibrium (LD).

If a specific allele in a given gene is directly involved in causing a particular trait T, its frequency will be statistically increased in a trait positive population when compared to the frequency in a trait negative population. As a consequence of the existence of linkage
20 disequilibrium, the frequency of all other alleles present in the haplotype carrying the trait-causing allele (TCA) will also be increased in trait positive individuals compared to trait negative individuals. Therefore, association between the trait and any allele in linkage disequilibrium with the trait-causing allele will suffice to suggest the presence of a trait-related gene in that particular allele's region. Linkage disequilibrium allows the relative frequencies in trait positive and trait
25 negative populations of a limited number of genetic polymorphisms (specifically biallelic markers) to be analyzed as an alternative to screening all possible functional polymorphisms in order to find trait-causing alleles.

The general strategy to perform association studies using biallelic markers derived from a candidate region is to scan two groups of individuals (trait positive and trait negative control
30 individuals which are characterized by a well defined phenotype as described below) in order to measure and statistically compare the allele frequencies of such biallelic markers in both groups.

If a statistically significant association with a trait is identified for at least one or more of the analyzed biallelic markers, one can assume that : either the associated allele is directly responsible for causing the trait (associated allele is the trait-causing allele), or the associated allele
35 is in linkage disequilibrium with the trait-causing allele. If the evidence indicates that the associated allele within the candidate region is most probably not the trait-causing allele but is in linkage

disequilibrium with the real trait-causing allele, then the trait-causing allele, and by consequence the gene carrying the trait-causing allele, can be found by sequencing the vicinity of the associated marker.

Collection of DNA samples from trait positive (trait +) and trait negative (trait -) individuals

5 (inclusion criteria)

In order to perform efficient and significant association studies such as those described herein, the trait under study should preferably follow a bimodal distribution in the population under study, presenting two clear non-overlapping phenotypes, trait positive and trait negative.

Nevertheless, even in the absence of such a bimodal distribution (as may in fact be the case for more complex genetic traits), any genetic trait may still be analyzed by the association method proposed here by carefully selecting the individuals to be included in the trait positive and trait negative phenotypic groups. The selection procedure involves to select individuals at opposite ends of the non-bimodal phenotype spectra of the trait under study, so as to include in these trait positive and trait negative populations individuals which clearly represent extreme, preferably non-overlapping phenotypes.

The definition of the inclusion criteria for the trait positive and trait negative populations is an important aspect of the present invention. The selection of drastically different but relatively uniform phenotypes enables efficient comparisons in association studies and the possible detection of marked differences at the genetic level, provided that the sample sizes of the populations under study are significant enough.

Generally, trait positive and trait negative populations to be included in association studies such as proposed in the present invention consist of phenotypically homogenous populations of individuals each representing 100% of the corresponding trait if the trait distribution is bimodal.

A first group of between 50 and 300 trait positive individuals, preferably about 100 individuals, can be recruited according to clinical inclusion criteria.

In each case, a similar number of trait negative individuals, preferably more than 100 individuals, are included in such studies who are preferably both ethnically- and age-matched to the trait positive cases. They are checked for the absence of the clinical criteria defined above. Both trait positive and trait negative individuals should correspond to unrelated cases.

30 Genotyping of trait positive and trait negative individuals

Allelic frequencies of the biallelic markers in each of the above described population can be determined using one of the methods described above under the heading "Methods of Genotyping DNA samples for biallelic markers". Analyses are preferably performed on amplified fragments obtained by genomic PCR performed on the DNA samples from each individual in similar conditions as those described above for the generation of biallelic markers.

In a preferred embodiment, amplified DNA samples are subjected to automated microsequencing reactions using fluorescent ddNTPs (specific fluorescence for each ddNTP) and

the appropriate microsequencing oligonucleotides which hybridize just upstream of the polymorphic base.

Genotyping is further described in Example 5.

Associations studies can be carried out by the skilled technician using the biallelic markers of the invention defined above, with different trait positive and trait negative populations. Suitable examples of association studies using biallelic markers of the *TBC-1* gene, including the biallelic markers A1 to A19, involve studies on the following populations:

- a trait positive population suffering from a cancer, preferably prostate cancer and a healthy unaffected population; or
- 10 - a trait positive population suffering from prostate cancer treated with agents acting against prostate cancer and suffering from side-effects resulting from this treatment and an trait negative population suffering from prostate cancer treated with same agents without any substantial side-effects, or
- a trait positive population suffering from prostate cancer treated with agents acting against prostate cancer showing a beneficial response and a trait negative population suffering from prostate cancer treated with same agents without any beneficial response, or
- 15 - a trait positive population suffering from prostate cancer presenting highly aggressive prostate cancer tumors and a trait negative population suffering from prostate cancer with prostate cancer tumors devoid of aggressiveness.
- 20 It is another object of the present invention to provide a method for the identification and characterization of an association between an allele of one or more biallelic markers of a *TBC-1* gene and a trait. The method comprises the steps of :
 - genotyping a marker or a group of biallelic markers according to the invention in trait positive;
 - 25 - genotyping a marker or a group of biallelic markers according to the invention in and trait negative individuals; and
 - establishing a statistically significant association between one allele of at least one marker and the trait.

Preferably, the trait positive and trait negative individuals are selected from non-overlapping phenotypes as regards to the trait under study. In one embodiment, the biallelic marker are selected from the group consisting of the biallelic markers A1 to A19.

In a preferred embodiment, the trait is cancer, prostate cancer, an early onset of prostate cancer, a susceptibility to prostate cancer, the level of aggressiveness of prostate cancer tumors, a modified expression of the *TBC-1* gene, a modified production of the TBC-1 protein, or the production of a modified TBC-1 protein.

In a further embodiment, the trait negative population can be replaced in the association studies by a random control population.

The step of testing for and detecting the presence of DNA comprising specific alleles of a biallelic marker or a group of biallelic markers of the present invention can be carried out as described further below.

Oligonucleotide Probes And Primers

- 5 The invention relates also to oligonucleotide molecules useful as probes or primers, wherein said oligonucleotide molecules hybridize specifically with a nucleotide sequence comprised in the *TBC-1* gene, particularly the *TBC-1* genomic sequence of SEQ ID Nos 1 and 2 or the *TBC-1* cDNAs sequences of SEQ ID Nos 3 and 4. More particularly, the present invention also concerns oligonucleotides for the detection of alleles of biallelic markers of the *TBC-1* gene. These
- 10 oligonucleotides are useful either as primers for use in various processes such as DNA amplification and microsequencing or as probes for DNA recognition in hybridization analyses. Polynucleotides derived from the *TBC-1* gene are useful in order to detect the presence of at least a copy of a nucleotide sequence of SEQ ID Nos 1-4, or a fragment, complement, or variant thereof in a test sample.
- 15 Particularly preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of a nucleotide sequence selected from the group consisting of SEQ ID Nos 1 and 2, or the complements thereof. Additionally preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides
- 20 comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 1 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 1: 1-1000, 1001-2000, 2001-3000, 3001-4000, 4001-5000, 5001-6000, 6001-7000, 7001-8000, 8001-9000, 9001-10000, 10001-11000, 11001-12000, 12001-13000, 13001-14000, 14001-15000,
- 25 15001-16000, 16001-17000, and 17001-17590. Other preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 2 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 2: 1-5000, 5001-10000, 10001-15000, 15001-
- 30 20000, 20001-25000, 25001-30000, 30001-35000, 35001-40000, 40001-45000, 45001-50000, 50001-55000, 55001-60000, 60001-65000, 65001-70000, 70001-75000, 75001-80000, 80001-85000, 85001-90000, 90001-95000, and 95001-99960.

Moreover, preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40,

35 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of a nucleotide sequence selected from the group consisting of SEQ ID Nos 3 and 4, or the complements thereof.. Particularly preferred

probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 3 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID

5 No 3: 1-500, 501-1000, 1001-1500, 1501-2000, 2001-2500, 2501-3000, 3001-3500, and 3501-3983.

Additional preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 4 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide

10 positions of SEQ ID No 4: 1-500, 501-1000, 1001-1500, 1501-2000, 2001-2500, 2501-3000, 3001-3500, and 3501-3988.

Thus, the invention also relates to nucleic acid probes characterized in that they hybridize specifically, under the stringent hybridization conditions defined above, with a nucleic acid selected from the group consisting of the nucleotide sequences of SEQ ID Nos 1-4 or a variant thereof or a

15 sequence complementary thereto.

In one embodiment the invention encompasses isolated, purified, and recombinant polynucleotides consisting of, or consisting essentially of a contiguous span of 8 to 50 nucleotides of any one of SEQ ID Nos 1 and 2 and the complement thereof, wherein said span includes a *TBC-1*-related biallelic marker in said sequence; optionally, wherein said *TBC-1*-related biallelic marker

20 is selected from the group consisting of A1 to A19, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, wherein said contiguous span is 18 to 35 nucleotides in length and said biallelic marker is within 4 nucleotides of the center of said polynucleotide; optionally, wherein said polynucleotide consists of said contiguous span and said contiguous span is 25 nucleotides in length and said biallelic marker is at the center of said

25 polynucleotide; optionally, wherein the 3' end of said contiguous span is present at the 3' end of said polynucleotide; and optionally, wherein the 3' end of said contiguous span is located at the 3' end of said polynucleotide and said biallelic marker is present at the 3' end of said polynucleotide.

In a preferred embodiment, said probes comprises, consists of, or consists essentially of a sequence selected from the following sequences: P1 to P7, P9 to P13, P15 to P19 and the complementary

30 sequences thereto.

In another embodiment the invention encompasses isolated, purified and recombinant polynucleotides comprising, consisting of, or consisting essentially of a contiguous span of 8 to 50 nucleotides of SEQ ID Nos 1 and 2, or the complements thereof, wherein the 3' end of said contiguous span is located at the 3' end of said polynucleotide, and wherein the 3' end of said

35 polynucleotide is located within 20 nucleotides upstream of a *TBC-1*-related biallelic marker in said sequence; optionally, wherein said *TBC-1*-related biallelic marker is selected from the group consisting of A1 to A19, and the complements thereof, or optionally the biallelic markers in linkage

disequilibrium therewith; optionally, wherein the 3' end of said polynucleotide is located 1 nucleotide upstream of said *TBC-1*-related biallelic marker in said sequence; and optionally, wherein said polynucleotide consists essentially of a sequence selected from the following sequences: D1 to D19 and E1 to E19.

- 5 In a further embodiment, the invention encompasses isolated, purified, or recombinant polynucleotides comprising, consisting of, or consisting essentially of a sequence selected from the following sequences: B1 to B15 and C1 to C15.

- In an additional embodiment, the invention encompasses polynucleotides for use in hybridization assays, sequencing assays, and enzyme-based mismatch detection assays for
10 determining the identity of the nucleotide at a *TBC-1*-related biallelic marker in SEQ ID Nos 1 and 2, or the complements thereof, as well as polynucleotides for use in amplifying segments of nucleotides comprising a *TBC-1*-related biallelic marker in SEQ ID Nos 1 and 2, or the complements thereof; optionally, wherein said *TBC-1*-related biallelic marker is selected from the group consisting of A1 to A19, and the complements thereof, or optionally the biallelic markers in
15 linkage disequilibrium therewith.

- A probe or a primer according to the invention has between 8 and 1000 nucleotides in length, or is specified to be at least 12, 15, 18, 20, 25, 35, 40, 50, 60, 70, 80, 100, 250, 500 or 1000 nucleotides in length. More particularly, the length of these probes and primers can range from 8, 10, 15, 20, or 30 to 100 nucleotides, preferably from 10 to 50, more preferably from 15 to 30
20 nucleotides. Shorter probes and primers tend to lack specificity for a target nucleic acid sequence and generally require cooler temperatures to form sufficiently stable hybrid complexes with the template. Longer probes and primers are expensive to produce and can sometimes self-hybridize to form hairpin structures. The appropriate length for primers and probes under a particular set of assay conditions may be empirically determined by one of skill in the art. A preferred probe or
25 primer consists of a nucleic acid comprising a polynucleotide selected from the group of the nucleotide sequences of P1 to P7, P9 to P13, P15 to P19 and the complementary sequence thereto, B1 to B15, C1 to C15, D1 to D19, E1 to E19, for which the respective locations in the sequence listing are provided in Tables 2, 3 and 4.

- The formation of stable hybrids depends on the melting temperature (T_m) of the DNA. The
30 T_m depends on the length of the primer or probe, the ionic strength of the solution and the G+C content. The higher the G+C content of the primer or probe, the higher is the melting temperature because G:C pairs are held by three H bonds whereas A:T pairs have only two. The GC content in the probes of the invention usually ranges between 10 and 75 %, preferably between 35 and 60 %, and more preferably between 40 and 55 %.

- 35 The primers and probes can be prepared by any suitable method, including, for example, cloning and restriction of appropriate sequences and direct chemical synthesis by a method such as the phosphodiester method of Narang et al.(1979), the phosphodiester method of Brown et

al.(1979), the diethylphosphoramidite method of Beaucage et al.(1981) and the solid support method described in EP 0 707 592.

Detection probes are generally nucleic acid sequences or uncharged nucleic acid analogs such as, for example peptide nucleic acids which are disclosed in International Patent Application WO 92/20702, morpholino analogs which are described in U.S. Patents Numbered 5,185,444; 5,034,506 and 5,142,047. The probe may have to be rendered "non-extendable" in that additional dNTPs cannot be added to the probe. In and of themselves analogs usually are non-extendable and nucleic acid probes can be rendered non-extendable by modifying the 3' end of the probe such that the hydroxyl group is no longer capable of participating in elongation. For example, the 3' end of the probe can be functionalized with the capture or detection label to thereby consume or otherwise block the hydroxyl group. Alternatively, the 3' hydroxyl group simply can be cleaved, replaced or modified, U.S. Patent Application Serial No. 07/049,061 filed April 19, 1993 describes modifications, which can be used to render a probe non-extendable.

Any of the polynucleotides of the present invention can be labeled, if desired, by incorporating any label known in the art to be detectable by spectroscopic, photochemical, biochemical, immunochemical, or chemical means. For example, useful labels include radioactive substances (including, ^{32}P , ^{35}S , ^3H , ^{125}I), fluorescent dyes (including, 5-bromodesoxyuridin, fluorescein, acetylaminofluorene, digoxigenin) or biotin. Preferably, polynucleotides are labeled at their 3' and 5' ends. Examples of non-radioactive labeling of nucleic acid fragments are described in the French patent No. FR-7810975 or by Urdea et al (1988) or Sanchez-Pescador et al (1988). In addition, the probes according to the present invention may have structural characteristics such that they allow the signal amplification, such structural characteristics being, for example, branched DNA probes as those described by Urdea et al. in 1991 or in the European patent No. EP 0 225 807 (Chiron).

A label can also be used to capture the primer, so as to facilitate the immobilization of either the primer or a primer extension product, such as amplified DNA, on a solid support. A capture label is attached to the primers or probes and can be a specific binding member which forms a binding pair with the solid's phase reagent's specific binding member (e.g. biotin and streptavidin). Therefore depending upon the type of label carried by a polynucleotide or a probe, it may be employed to capture or to detect the target DNA. Further, it will be understood that the polynucleotides, primers or probes provided herein, may, themselves, serve as the capture label. For example, in the case where a solid phase reagent's binding member is a nucleic acid sequence, it may be selected such that it binds a complementary portion of a primer or probe to thereby immobilize the primer or probe to the solid phase. In cases where a polynucleotide probe itself serves as the binding member, those skilled in the art will recognize that the probe will contain a sequence or "tail" that is not complementary to the target. In the case where a polynucleotide

primer itself serves as the capture label, at least a portion of the primer will be free to hybridize with a nucleic acid on a solid phase. DNA Labeling techniques are well known to the skilled technician.

The probes of the present invention are useful for a number of purposes. They can be notably used in Southern hybridization to genomic DNA. The probes can also be used to detect
5 PCR amplification products. They may also be used to detect mismatches in the *TBC-1* gene or mRNA using other techniques.

Any of the polynucleotides, primers and probes of the present invention can be conveniently immobilized on a solid support. Solid supports are known to those skilled in the art and include the walls of wells of a reaction tray, test tubes, polystyrene beads, magnetic beads,
10 nitrocellulose strips, membranes, microparticles such as latex particles, sheep (or other animal) red blood cells, duracytes and others. The solid support is not critical and can be selected by one skilled in the art. Thus, latex particles, microparticles, magnetic or non-magnetic beads, membranes, plastic tubes, walls of microtiter wells, glass or silicon chips, sheep (or other suitable animal's) red blood cells and duracytes are all suitable examples. Suitable methods for immobilizing nucleic
15 acids on solid phases include ionic, hydrophobic, covalent interactions and the like. A solid support, as used herein, refers to any material which is insoluble, or can be made insoluble by a subsequent reaction. The solid support can be chosen for its intrinsic ability to attract and immobilize the capture reagent. Alternatively, the solid phase can retain an additional receptor which has the ability to attract and immobilize the capture reagent. The additional receptor can
20 include a charged substance that is oppositely charged with respect to the capture reagent itself or to a charged substance conjugated to the capture reagent. As yet another alternative, the receptor molecule can be any specific binding member which is immobilized upon (attached to) the solid support and which has the ability to immobilize the capture reagent through a specific binding reaction. The receptor molecule enables the indirect binding of the capture reagent to a solid
25 support material before the performance of the assay or during the performance of the assay. The solid phase thus can be a plastic, derivatized plastic, magnetic or non-magnetic metal, glass or silicon surface of a test tube, microtiter well, sheet, bead, microparticle, chip, sheep (or other suitable animal's) red blood cells, duracytes® and other configurations known to those of ordinary skill in the art. The polynucleotides of the invention can be attached to or immobilized on a solid
30 support individually or in groups of at least 2, 5, 8, 10, 12, 15, 20, or 25 distinct polynucleotides of the invention to a single solid support. In addition, polynucleotides other than those of the invention may be attached to the same solid support as one or more polynucleotides of the invention.

Consequently, the invention also deals with a method for detecting the presence of a nucleic
35 acid comprising a nucleotide sequence selected from a group consisting of SEQ ID Nos 1-4, a fragment or a variant thereof and a complementary sequence thereto in a sample, said method comprising the following steps of:

a) bringing into contact a nucleic acid probe or a plurality of nucleic acid probes which can hybridize with a nucleotide sequence included in a nucleic acid selected from the group consisting of the nucleotide sequences of SEQ ID Nos 1-4, a fragment or a variant thereof and a complementary sequence thereto and the sample to be assayed; and

5 b) detecting the hybrid complex formed between the probe and a nucleic acid in the sample.

The invention further concerns a kit for detecting the presence of a nucleic acid comprising a nucleotide sequence selected from a group consisting of SEQ ID Nos 1-4, a fragment or a variant thereof and a complementary sequence thereto in a sample, said kit comprising:

a) a nucleic acid probe or a plurality of nucleic acid probes which can hybridize with a
10 nucleotide sequence included in a nucleic acid selected from the group consisting of the nucleotide sequences of SEQ ID Nos 1-4, a fragment or a variant thereof and a complementary sequence thereto; and

b) optionally, the reagents necessary for performing the hybridization reaction.

In a first preferred embodiment of this detection method and kit, said nucleic acid probe or
15 the plurality of nucleic acid probes are labeled with a detectable molecule. In a second preferred embodiment of said method and kit, said nucleic acid probe or the plurality of nucleic acid probes has been immobilized on a substrate. In a third preferred embodiment, the nucleic acid probe or the plurality of nucleic acid probes comprise either a sequence which is selected from the group consisting of the nucleotide sequences of P1 to P7, P9 to P13, P15 to P19 and the complementary
20 sequence thereto, B1 to B15, C1 to C15, D1 to D19, E1 to E19 or a biallelic marker selected from the group consisting of A1 to A19 and the complements thereto.

Oligonucleotide Arrays

A substrate comprising a plurality of oligonucleotide primers or probes of the invention may be used either for detecting or amplifying targeted sequences in the *TBC-1* gene and may also
25 be used for detecting mutations in the coding or in the non-coding sequences of the *TBC-1* gene.

Any polynucleotide provided herein may be attached in overlapping areas or at random locations on the solid support. Alternatively the polynucleotides of the invention may be attached in an ordered array wherein each polynucleotide is attached to a distinct region of the solid support which does not overlap with the attachment site of any other polynucleotide. Preferably, such an
30 ordered array of polynucleotides is designed to be "addressable" where the distinct locations are recorded and can be accessed as part of an assay procedure. Addressable polynucleotide arrays typically comprise a plurality of different oligonucleotide probes that are coupled to a surface of a substrate in different known locations. The knowledge of the precise location of each polynucleotides location makes these "addressable" arrays particularly useful in hybridization
35 assays. Any addressable array technology known in the art can be employed with the polynucleotides of the invention. One particular embodiment of these polynucleotide arrays is known as the Genechips™, and has been generally described in US Patent 5,143,854; PCT

publications WO 90/15070 and 92/10092. These arrays may generally be produced using mechanical synthesis methods or light directed synthesis methods which incorporate a combination of photolithographic methods and solid phase oligonucleotide synthesis (Fodor et al., 1991). The immobilization of arrays of oligonucleotides on solid supports has been rendered possible by the development of a technology generally identified as "Very Large Scale Immobilized Polymer Synthesis" (VLSIPSTTM) in which, typically, probes are immobilized in a high density array on a solid surface of a chip. Examples of VLSIPSTTM technologies are provided in US Patents 5,143,854; and 5,412,087 and in PCT Publications WO 90/15070, WO 92/10092 and WO 95/11995, which describe methods for forming oligonucleotide arrays through techniques such as light-directed synthesis techniques. In designing strategies aimed at providing arrays of nucleotides immobilized on solid supports, further presentation strategies were developed to order and display the oligonucleotide arrays on the chips in an attempt to maximize hybridization patterns and sequence information. Examples of such presentation strategies are disclosed in PCT Publications WO 94/12305, WO 94/11530, WO 97/29212 and WO 97/31256.

In another embodiment of the oligonucleotide arrays of the invention, an oligonucleotide probe matrix may advantageously be used to detect mutations occurring in the *TBC-1* gene and preferably in its regulatory region. For this particular purpose, probes are specifically designed to have a nucleotide sequence allowing their hybridization to the genes that carry known mutations (either by deletion, insertion or substitution of one or several nucleotides). By known mutations, it is meant, mutations on the *TBC-1* gene that have been identified according, for example to the technique used by Huang et al.(1996) or Samson et al.(1996).

Another technique that is used to detect mutations in the *TBC-1* gene is the use of a high-density DNA array. Each oligonucleotide probe constituting a unit element of the high density DNA array is designed to match a specific subsequence of the *TBC-1* genomic DNA or cDNA. Thus, an array consisting of oligonucleotides complementary to subsequences of the target gene sequence is used to determine the identity of the target sequence with the wild gene sequence, measure its amount, and detect differences between the target sequence and the reference wild gene sequence of the *TBC-1* gene. In one such design, termed 4L tiled array, is implemented a set of four probes (A, C, G, T), preferably 15-nucleotide oligomers. In each set of four probes, the perfect complement will hybridize more strongly than mismatched probes. Consequently, a nucleic acid target of length L is scanned for mutations with a tiled array containing 4L probes, the whole probe set containing all the possible mutations in the known wild reference sequence. The hybridization signals of the 15-mer probe set tiled array are perturbed by a single base change in the target sequence. As a consequence, there is a characteristic loss of signal or a "footprint" for the probes flanking a mutation position. This technique was described by Chee et al. in 1996.

Consequently, the invention concerns an array of nucleic acid molecules comprising at least one polynucleotide described above as probes and primers. Preferably, the invention concerns an

array of nucleic acid comprising at least two polynucleotides described above as probes and primers.

A further object of the invention consists of an array of nucleic acid sequences comprising either at least one of the sequences selected from the group consisting of P1 to P7, P9 to P13, P15 to P19, B1 to B15, C1 to C15, D1 to D19, E1 to E19, the sequences complementary thereto, a fragment thereof of at least 8, 10, 12, 15, 18, 20, 25, 30, or 40 consecutive nucleotides thereof, and at least one sequence comprising a biallelic marker selected from the group consisting of A1 to A19 and the complements thereto.

The invention also pertains to an array of nucleic acid sequences comprising either at least two of the sequences selected from the group consisting of P1 to P7, P9 to P13, P15 to P19, B1 to B15, C1 to C15, D1 to D19, E1 to E19, the sequences complementary thereto, a fragment thereof of at least 8 consecutive nucleotides thereof, and at least two sequences comprising a biallelic marker selected from the group consisting of A1 to A19 and the complements thereof.

Vectors For The Expression Of A Regulatory Or A Coding Polynucleotide Of *TBC-1*.

Any of the regulatory polynucleotides or the coding polynucleotides of the invention may be inserted into recombinant vectors for expression in a recombinant host cell or a recombinant host organism.

Thus, the present invention also encompasses a family of recombinant vectors that contains either a regulatory polynucleotide selected from the group consisting of any one of the regulatory polynucleotides derived from the *TBC-1* genomic sequences of SEQ ID Nos 1 and 2, or a polynucleotide comprising the *TBC-1* coding sequence, or both.

In a first preferred embodiment, a recombinant vector of the invention is used as an expression vector : (a) the *TBC-1* regulatory sequence comprised therein drives the expression of a coding polynucleotide operably linked thereto; (b) the *TBC-1* coding sequence is operably linked to regulation sequences allowing its expression in a suitable cell host and/or host organism.

In a second preferred embodiment, a recombinant vector of the invention is used to amplify the inserted polynucleotide derived from the *TBC-1* genomic sequences of SEQ ID Nos 1 and 2 or *TBC-1* cDNAs in a suitable cell host , this polynucleotide being amplified at every time that the recombinant vector replicates.

More particularly, the present invention relates to expression vectors which include nucleic acids encoding a TBC-1 protein, preferably the TBC-1 protein of the amino acid sequence of SEQ ID No 5 described therein, under the control of a regulatory sequence selected among the *TBC-1* regulatory polynucleotides, or alternatively under the control of an exogenous regulatory sequence.

A recombinant expression vector comprising a nucleic acid selected from the group consisting of 5' and 3' regulatory regions, or biologically active fragments or variants thereof, is also part of the present invention.

The invention also encompasses a recombinant expression vector comprising :

- a) a nucleic acid comprising the 5' regulatory polynucleotide of the nucleotide sequence SEQ ID No 1, or a biologically active fragment or variant thereof;
- b) a polynucleotide encoding a polypeptide or a polynucleotide of interest operably linked
5 with said nucleic acid.

- c) optionally, a nucleic acid comprising a 3'-regulatory polynucleotide, preferably a 3'-regulatory polynucleotide of the invention, or a biologically active fragment or variant thereof.

The nucleic acid comprising the 5' regulatory polynucleotide or a biologically active fragment or variant thereof may also comprises the 5'-UTR sequence from any of the two cDNA of
10 the invention or a biologically active fragment or variant thereof.

The invention also pertains to a recombinant expression vector useful for the expression of the *TBC-1* coding sequence, wherein said vector comprises a nucleic acid selected from the group consisting of SEQ ID Nos 3 and 4 or a nucleic acid having at least 95% nucleotide identity with a polynucleotide selected from the group consisting of the nucleotide sequences of SEQ ID Nos 3 and
15 4.

Another recombinant expression vector of the invention consists in a recombinant vector comprising a nucleic acid comprising the nucleotide sequence beginning at the nucleotide in position 176 and ending in position 3730 of the polynucleotide of SEQ ID No 4.

Generally, a recombinant vector of the invention may comprise any of the polynucleotides
20 described herein, including regulatory sequences, and coding sequences, as well as any *TBC-1* primer or probe as defined above. More particularly, the recombinant vectors of the present invention can comprise any of the polynucleotides described in the "*TBC-1* cDNA Sequences" section, the "Coding Regions" section, "Genomic sequence of *TBC-1*" section and the "Oligonucleotide Probes And Primers" section.

25 Some of the elements which can be found in the vectors of the present invention are described in further detail in the following sections.

a) Vectors

A recombinant vector according to the invention comprises, but is not limited to, a YAC (Yeast Artificial Chromosome), a BAC (Bacterial Artificial Chromosome), a phage, a phagemid, a
30 cosmid, a plasmid or even a linear DNA molecule which may consist of a chromosomal, non-chromosomal and synthetic DNA. Such a recombinant vector can comprise a transcriptional unit comprising an assembly of :

- (1) a genetic element or elements having a regulatory role in gene expression, for example promoters or enhancers. Enhancers are cis-acting elements of DNA, usually from about 10 to 300
35 bp in length that act on the promoter to increase the transcription.

- (2) a structural or coding sequence which is transcribed into mRNA and eventually translated into a polypeptide, and

(3) appropriate transcription initiation and termination sequences. Structural units intended for use in yeast or eukaryotic expression systems preferably include a leader sequence enabling extracellular secretion of translated protein by a host cell. Alternatively, where a recombinant protein is expressed without a leader or transport sequence, it may include an N-terminal residue.

- 5 This residue may or may not be subsequently cleaved from the expressed recombinant protein to provide a final product.

Generally, recombinant expression vectors will include origins of replication, selectable markers permitting transformation of the host cell, and a promoter derived from a highly expressed gene to direct transcription of a downstream structural sequence. The heterologous structural
10 sequence is assembled in appropriate phase with translation initiation and termination sequences, and preferably a leader sequence capable of directing secretion of the translated protein into the periplasmic space or the extracellular medium.

The selectable marker genes for selection of transformed host cells are preferably dihydrofolate reductase or neomycin resistance for eukaryotic cell culture, TRP1 for *S. cerevisiae* or
15 tetracycline, rifampicin or ampicillin resistance in *E. coli*, or levan saccharase for mycobacteria.

As a representative but non-limiting example, useful expression vectors for bacterial use can comprise a selectable marker and a bacterial origin of replication derived from commercially available plasmids comprising genetic elements of pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Pharmacia, Uppsala, Sweden), and GEM1 (Promega
20 Biotec, Madison, WI, USA).

Large numbers of suitable vectors and promoters are known to those of skill in the art, and commercially available, such as bacterial vectors : pQE70, pQE60, pQE-9 (Qiagen), pbs, pD10, phagescript, psiX174, pbluescript SK, pbsks, pNH8A, pNH16A, pNH18A, pNH46A (Stratagene); p25 pSV2CAT, pOG44, pXT1, pSG (Stratagene); pSVK3, pBPV, pMSG, pSVL (Pharmacia); baculovirus transfer vector pVL1392/1393 (Pharminogen); pQE-30 (QIAexpress).

A suitable vector for the expression of the TBC-1 polypeptide of SEQ ID No 5 is a baculovirus vector that can be propagated in insect cells and in insect cell lines. A specific suitable host vector system is the pVL1392/1393 baculovirus transfer vector (Pharminogen) that is used to
30 transfect the SF9 cell line (ATCC N^oCRL 1711) which is derived from *Spodoptera frugiperda*.

Other suitable vectors for the expression of the TBC-1 polypeptide of SEQ ID No 5 in a baculovirus expression system include those described by Chai et al. (1993), Vlasak et al. (1983) and Lenhard et al. (1996).

Mammalian expression vectors will comprise an origin of replication, a suitable promoter
35 and enhancer, and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 viral genome, for example SV40 origin, early promoter,

enhancer, splice and polyadenylation sites may be used to provide the required nontranscribed genetic elements.

b) Promoters

The suitable promoter regions used in the expression vectors according to the present invention are chosen taking into account the cell host in which the heterologous gene has to be expressed.

A suitable promoter may be heterologous with respect to the nucleic acid for which it controls the expression or alternatively can be endogenous to the native polynucleotide containing the coding sequence to be expressed. Additionally, the promoter is generally heterologous with respect to the recombinant vector sequences within which the construct promoter/coding sequence has been inserted.

Preferred bacterial promoters are the LacI, LacZ, the T3 or T7 bacteriophage RNA polymerase promoters, the polyhedrin promoter, or the p10 protein promoter from baculovirus (Kit Novagen) (Smith et al., 1983; O'Reilly et al., 1992), the lambda P_R promoter or also the trc promoter.

Promoter regions can be selected from any desired gene using, for example, CAT (chloramphenicol transferase) vectors and more preferably pKK232-8 and pCM7 vectors. Particularly preferred bacterial promoters include lacI, lacZ, T3, T7, gpt, lambda PR, PL and trp. Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein-L. Selection of a convenient vector and promoter is well within the level of ordinary skill in the art.

The choice of a promoter is well within the ability of a person skilled in the field of genetic engineering. For example, one may refer to the book of Sambrook et al. (1989) or also to the procedures described by Fuller et al. (1996).

The vector containing the appropriate DNA sequence as described above, more preferably a *TBC-1* gene regulatory polynucleotide, a polynucleotide encoding the TBC-1 polypeptide of SEQ ID No 5 or both of them, can be utilized to transform an appropriate host to allow the expression of the desired polypeptide or polynucleotide.

c) Other types of vectors

The *in vivo* expression of a TBC-1 polypeptide of SEQ ID No 5 may be useful in order to correct a genetic defect related to the expression of the native gene in a host organism or to the production of a biologically inactive TBC-1 protein.

Consequently, the present invention also deals with recombinant expression vectors mainly designed for the *in vivo* production of the TBC-1 polypeptide of SEQ ID No 5 by the introduction of the appropriate genetic material in the organism of the patient to be treated. This genetic material may be introduced *in vitro* in a cell that has been previously extracted from the organism, the

modified cell being subsequently reintroduced in the said organism, directly *in vivo* into the appropriate tissue.

By « vector » according to this specific embodiment of the invention is intended either a circular or a linear DNA molecule.

5 One specific embodiment for a method for delivering a protein or peptide to the interior of a cell of a vertebrate *in vivo* comprises the step of introducing a preparation comprising a physiologically acceptable carrier and a naked polynucleotide operatively coding for the polypeptide of interest into the interstitial space of a tissue comprising the cell, whereby the naked polynucleotide is taken up into the interior of the cell and has a physiological effect.

10 In a specific embodiment, the invention provides a composition for the *in vivo* production of the TBC-1 protein or polypeptide described herein. It comprises a naked polynucleotide operatively coding for this polypeptide, in solution in a physiologically acceptable carrier, and suitable for introduction into a tissue to cause cells of the tissue to express the said protein or polypeptide.

Compositions comprising a polynucleotide are described in PCT application N° WO
15 90/11092 (Vical Inc.) and also in PCT application N° WO 95/11307 (Institut Pasteur, INSERM, Université d'Ottawa) as well as in the articles of Tacson et al. (1996) and of Huygen et al. (1996).

The amount of vector to be injected to the desired host organism varies according to the site of injection. As an indicative dose, it will be injected between 0,1 and 100 µg of the vector in an animal body, preferably a mammal body, for example a mouse body.

20 In another embodiment of the vector according to the invention, it may be introduced *in vitro* in a host cell, preferably in a host cell previously harvested from the animal to be treated and more preferably a somatic cell such as a muscle cell. In a subsequent step, the cell that has been transformed with the vector coding for the desired TBC-1 polypeptide or the desired fragment thereof is reintroduced into the animal body in order to deliver the recombinant protein within the
25 body either locally or systemically.

In one specific embodiment, the vector is derived from an adenovirus. Preferred adenovirus vectors according to the invention are those described by Feldman and Steg (1996) or Ohno et al. (1994). Another preferred recombinant adenovirus according to this specific embodiment of the present invention is the human adenovirus type 2 or 5 (Ad 2 or Ad 5) or an adenovirus of animal
30 origin (French patent application N° FR-93.05954).

Retrovirus vectors and adeno-associated virus vectors are generally understood to be the recombinant gene delivery systems of choice for the transfer of exogenous polynucleotides *in vivo* , particularly to mammals, including humans. These vectors provide efficient delivery of genes into cells, and the transferred nucleic acids are stably integrated into the chromosomal DNA of the host
35 Particularly preferred retroviruses for the preparation or construction of retroviral *in vitro* or *in vitro* gene delivery vehicles of the present invention include retroviruses selected from the group consisting of Mink-Cell Focus Inducing Virus, Murine Sarcoma Virus, Reticuloendotheliosis virus

and Rous Sarcoma virus. Particularly preferred Murine Leukemia Viruses include the 4070A and the 1504A viruses, Abelson (ATCC No VR-999), Friend (ATCC No VR-245), Gross (ATCC No VR-590), Rauscher (ATCC No VR-998) and Moloney Murine Leukemia Virus (ATCC No VR-190; PCT Application No WO 94/24298). Particularly preferred Rous Sarcoma Viruses include
5 Bryan high titer (ATCC Nos VR-334, VR-657, VR-726, VR-659 and VR-728). Other preferred retroviral vectors are those described in Roth et al. (Roth J.A. et al., 1996), PCT Application No WO 93/25234, PCT Application No WO 94/ 06920, Roux et al., 1989, Julan et al., 1992 and Neda et al., 1991.

Yet another viral vector system that is contemplated by the invention consists in the adeno-
10 associated virus (AAV). The adeno-associated virus is a naturally occurring defective virus that requires another virus, such as an adenovirus or a herpes virus, as a helper virus for efficient replication and a productive life cycle (Muzyczka et al., 1992). It is also one of the few viruses that may integrate its DNA into non-dividing cells, and exhibits a high frequency of stable integration (Flotte et al., 1992; Samulski et al., 1989; McLaughlin et al., 1989). One advantageous feature of
15 AAV derives from its reduced efficacy for transducing primary cells relative to transformed cells.

Other compositions containing a vector of the invention advantageously comprise an oligonucleotide fragment of a nucleic sequence selected from the group consisting of SEQ ID Nos 3 or 4 as an antisense tool that inhibits the expression of the corresponding *TBC-1* gene. Preferred methods using antisense polynucleotide according to the present invention are the procedures
20 described by Sczakiel et al. (1995) or those described in PCT Application No WO 95/24223.

Host cells

Another object of the invention consists in host cell that have been transformed or transfected with one of the polynucleotides described therein, and more precisely a polynucleotide either comprising a *TBC-1* regulatory polynucleotide or the coding sequence of the *TBC-1*
25 polypeptide having the amino acid sequence of SEQ ID No 5. Are included host cells that are transformed (prokaryotic cells) or that are transfected (eukaryotic cells) with a recombinant vector such as one of those described above.

A recombinant host cell of the invention comprises any one of the polynucleotides or the recombinant vectors described therein. More particularly, the cell hosts of the present invention can
30 comprise any of the polynucleotides described in "*TBC-1* cDNA Sequences" section, the "Coding Regions" section, "Genomic sequence of *TBC-1*" section and the "Oligonucleotide Probes And Primers" section.

Another preferred recombinant cell host according to the present invention is characterized in that its genome or genetic background (including chromosome, plasmids) is modified by the
35 nucleic acid coding for the *TBC-1* polypeptide of SEQ ID No 5.

Preferred host cells used as recipients for the expression vectors of the invention are the following :

- a) Prokaryotic host cells : *Escherichia coli* strains (I.E. DH5- α strain) or *Bacillus subtilis*.
- b) Eukaryotic host cells : HeLa cells (ATCC N^oCCL2; N^oCCL2.1; N^oCCL2.2), Cv 1 cells
- 5 (ATCC N^oCCL70), COS cells (ATCC N^oCRL1650; N^oCRL1651), Sf-9 cells (ATCC N^oCRL1711).

The constructs in the host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence.

Following transformation of a suitable host and growth of the host to an appropriate cell density, the selected promoter is induced by appropriate means, such as temperature shift or

10 chemical induction, and cells are cultivated for an additional period.

Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification.

Microbial cells employed in the expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing

15 agents. Such methods are well known by the skill artisan.

Transgenic animals

The terms "transgenic animals" or "host animals" are used herein to designate animals that have their genome genetically and artificially manipulated so as to include one of the nucleic acids according to the invention. Preferred animals are non-human mammals and include those belonging

20 to a genus selected from *Mus* (e.g. mice), *Rattus* (e.g. rats) and *Oryctogalus* (e.g. rabbits) which have their genome artificially and genetically altered by the insertion of a nucleic acid according to the invention.

The transgenic animals of the invention all include within a plurality of their cells a cloned recombinant or synthetic DNA sequence, more specifically one of the purified or isolated nucleic acids comprising a *TBC-1* coding sequence, a *TBC-1* regulatory polynucleotide or a DNA sequence

25 encoding an antisense polynucleotide such as described in the present specification.

More particularly, transgenic animals according to the invention contain in their somatic cells and/or in their germ line cells any of the polynucleotides described in "*TBC-1* cDNA Sequences" section, the "Coding Regions" section, "Genomic sequence of *TBC-1*" section, the

30 "Oligonucleotide Probes And Primers" section and the "Vectors for the expression of a regulatory or coding polynucleotide of *TBC-1*" section.

The transgenic animals of the invention thus contain specific sequences of exogenous genetic material such as the nucleotide sequences described above in detail.

In a first preferred embodiment, these transgenic animals may be good experimental models

35 in order to study the diverse pathologies related to cell differentiation, in particular concerning the

transgenic animals within the genome of which has been inserted one or several copies of a polynucleotide encoding a native TBC-1 protein, or alternatively a mutant TBC-1 protein.

In a second preferred embodiment, these transgenic animals may express a desired polypeptide of interest under the control of the regulatory polynucleotides of the *TBC-1* gene, leading to good yields in the synthesis of this protein of interest, and eventually a tissue specific expression of this protein of interest.

Since it is possible to produce transgenic animals of the invention using a variety of different sequences, a general description will be given of the production of transgenic animals by referring generally to exogenous genetic material. This general description can be adapted by those skilled in the art in order to incorporate the DNA sequences into animals. For more details regarding the production of transgenic animals, and specifically transgenic mice, it may be referred to Sandou et al. (1994) and also to US Patents Nos 4,873,191, issued Oct.10, 1989, 5,968,766, issued Dec. 16, 1997 and 5,387,742, issued Feb. 28, 1995, these documents being herein incorporated by reference to disclose methods for producing transgenic mice.

Transgenic animals of the present invention are produced by the application of procedures which result in an animal with a genome that incorporates exogenous genetic material which is integrated into the genome. The procedure involves obtaining the genetic material, or a portion thereof, which encodes either a *TBC-1* coding sequence, a *TBC-1* regulatory polynucleotide or a DNA sequence encoding an antisense polynucleotide such as described in the present specification.

A recombinant polynucleotide of the invention is inserted into an embryonic or ES stem cell line. The insertion is made using electroporation. The cells subjected to electroporation are screened (e.g. Southern blot analysis) to find positive cells which have integrated the exogenous recombinant polynucleotide into their genome. An illustrative positive-negative selection procedure that may be used according to the invention is described by Mansour et al. (1988). Then, the positive cells are isolated, cloned and injected into 3.5 days old blastocysts from mice. The blastocysts are then inserted into a female host animal and allowed to grow to term. The offsprings of the female host are tested to determine which animals are transgenic e.g. include the inserted exogenous DNA sequence and which are wild-type.

Screening Of Agents Interacting With TBC-1

In a further embodiment, the present invention also concerns a method for the screening of new agents, or candidate substances interacting with TBC-1. These new agents could be useful against cancer.

In a preferred embodiment, the invention relates to a method for the screening of candidate substances comprising the following steps:

- providing a cell line, an organ, or a mammal expressing a *TBC-1* gene or a fragment thereof, preferably the regulatory region or the promoter region of the *TBC-1* gene.

- obtaining a candidate substance preferably a candidate substance capable of inhibiting the binding of a transcription factor to the *TBC-1* regulatory region,
- testing the ability of the candidate substance to decrease the symptoms of prostate cancer and/or to modulate the expression levels of *TBC-1*.

5 In some embodiments, the cell line, organ or mammal expresses a heterologous protein, the coding sequence of which is operably linked to the *TBC-1* regulatory or promoter sequence. In other embodiments, they express a *TBC-1* gene comprising alleles of one or more *TBC-1*-related biallelic markers.

A candidate substance is a substance which can interact with or modulate, by binding or
10 other intramolecular interactions, expression, stability, and function of *TBC-1*. Such substances may be potentially interesting for patients who are not responsive to existing drugs or develop side effects to them. Screening may be effected using either *in vitro* methods or *in vivo* methods.

Such methods can be carried out in numerous ways such as on transformed cells which express the considered alleles of the *TBC-1* gene, on tumors induced by said transformed cells, for
15 example in mice, or on a *TBC-1* protein encoded by the considered allelic variant of *TBC-1*.

Screening assays of the present invention generally involve determining the ability of a candidate substance to present a cytotoxic effect, to change the characteristics of transformed cells such as proliferative and invasive capacity, to affect the tumor growth, or to modify the expression level of *TBC-1*.

20 Typically, this method includes preparing transformed cells with different forms of *TBC-1* sequences containing particular alleles of one or more biallelic markers and/or trait causing mutations described above. This is followed by testing the cells expressing the *TBC-1* with a candidate substance to determine the ability of the substance to present cytotoxic effect, to affect the characteristics of transformed cells, the tumor growth, or to modify the expression level of *TBC-1*.

25 Typical examples of such drug screening assays are provided below. It is to be understood that the parameters set forth in these examples can be modified by the skilled person without undue experimentation.

Methods for screening substances interacting with a *TBC-1* polypeptide

A method for the screening of a candidate substance according to the invention comprises
30 the following steps :

a) providing a polypeptide comprising the amino acid sequence SEQ ID No 5, or a peptide fragment or a variant thereof;

b) obtaining a candidate substance;

c) bringing into contact said polypeptide with said candidate substance;

35 d) detecting the complexes formed between said polypeptide and said candidate substance.

For the purpose of the present invention, a ligand means a molecule, such as a protein, a peptide, an antibody or any synthetic chemical compound capable of binding to the *TBC-1* protein

or one of its fragments or variants or to modulate the expression of the polynucleotide coding for TBC-1 or a fragment or variant thereof.

In the ligand screening method according to the present invention, a biological sample or a defined molecule to be tested as a putative ligand of the TBC-1 protein is brought into contact with
5 a purified TBC-1 protein, for example a purified recombinant TBC-1 protein produced by a recombinant cell host as described hereinbefore, in order to form a complex between the TBC-1 protein and the putative ligand molecule to be tested.

A. Candidate ligands obtained from random peptide libraries

In a particular embodiment of the screening method, the putative ligand is the expression
10 product of a DNA insert contained in a phage vector (Parmley and Smith, 1988). Specifically, random peptide phages libraries are used. The random DNA inserts encode peptides of 8 to 20 aminoacids in length (Oldenburg K.R. et al., 1992,.; Valadon P., et al., 1996; Lucas A.H., 1994; Westerink M.A.J., 1995; Castagnoli L. et al., 1991). According to this particular embodiment, the recombinant phages expressing a protein that binds to the immobilized TBC-1 protein are retained
15 and the complex formed between the TBC-1 protein and the recombinant phage may be subsequently immunoprecipitated by a polyclonal or a monoclonal antibody directed against the TBC-1 protein.

Once the ligand library in recombinant phages has been constructed, the phage population is brought into contact with the immobilized TBC-1 protein. Then the preparation of complexes is
20 washed in order to remove the non-specifically bound recombinant phages. The phages that bind specifically to the TBC-1 protein are then eluted by a buffer (acid pH) or immunoprecipitated by the anti-TBC-1 monoclonal antibody produced by a hybridoma, and this phage population is subsequently amplified by an over-infection of bacteria (for example *E. coli*). The selection step may be repeated several times, preferably 2-4 times, in order to select the more specific
25 recombinant phage clones. The last step consists in characterizing the peptide produced by the selected recombinant phage clones either by expression in infected bacteria and isolation, expressing the phage insert in another host-vector system, or sequencing the insert contained in the selected recombinant phages.

B. Candidate ligands obtained through a two-hybrid screening assay.

30 The yeast two-hybrid system is designed to study protein-protein interactions *in vivo* (Fields and Song, 1989), and relies upon the fusion of a bait protein to the DNA binding domain of the yeast Gal4 protein. This technique is also described in US Patent N° US 5,667,973 and US Patent N° 5,283,173 (Fields et al.) the technical teachings of both patents being herein incorporated by reference.

35 The general procedure of library screening by the two-hybrid assay may be performed as described by Harper et al. (Harper JW et al., 1993) or as described by Cho et al. (1998) or also Fromont-Racine et al. (1997).

The bait protein or polypeptide consists of a TBC-1 polypeptide or a fragment or variant thereof.

More precisely, the nucleotide sequence encoding the TBC-1 polypeptide or a fragment or variant thereof is fused to a polynucleotide encoding the DNA binding domain of the GAL4 protein, the fused nucleotide sequence being inserted in a suitable expression vector, for example pAS2 or pM3.

Then, a human cDNA library is constructed in a specially designed vector, such that the human cDNA insert is fused to a nucleotide sequence in the vector that encodes the transcriptional domain of the GAL4 protein. Preferably, the vector used is the pACT vector. The polypeptides encoded by the nucleotide inserts of the human cDNA library are termed "pray" polypeptides.

A third vector contains a detectable marker gene, such as beta galactosidase gene or CAT gene that is placed under the control of a regulation sequence that is responsive to the binding of a complete Gal4 protein containing both the transcriptional activation domain and the DNA binding domain. For example, the vector pG5EC may be used.

Two different yeast strains are also used. As an illustrative but non limiting example the two different yeast strains may be the following :

- Y190, the phenotype of which is (*MATa, Leu2-3, 112 ura3-12, trp1-901, his3-D200, ade2-101, gal4Dgal180D URA3 GAL-LacZ, LYS GAL-HIS3, cyh'*);
- Y187, the phenotype of which is (*MATa gal4 gal80 his3 trp1-901 ade2-101 ura3-52 leu2-3, -112 URA3 GAL-lacZmet'*), which is the opposite mating type of Y190.

Briefly, 20 µg of pAS2/TBC-1 and 20 µg of pACT-cDNA library are co-transformed into yeast strain Y190. The transformants are selected for growth on minimal media lacking histidine, leucine and tryptophan, but containing the histidine synthesis inhibitor 3-AT (50 mM). Positive colonies are screened for beta galactosidase by filter lift assay. The double positive colonies (*His*⁺, *beta-gal*⁺) are then grown on plates lacking histidine, leucine, but containing tryptophan and cycloheximide (10 mg/ml) to select for loss of pAS2/TBC-1 plasmids but retention of pACT-cDNA library plasmids. The resulting Y190 strains are mated with Y187 strains expressing TBC-1 or non-related control proteins; such as cyclophilin B, lamin, or SNF1, as *Gal4* fusions as described by Harper et al. (1993) and by Bram et al. (1993), and screened for beta galactosidase by filter lift assay. Yeast clones that are *beta gal-* after mating with the control *Gal4* fusions are considered false positives.

In another embodiment of the two-hybrid method according to the invention, the interaction between TBC-1 or a fragment or variant thereof with cellular proteins may be assessed using the Matchmaker Two Hybrid System 2 (Catalog No. K1604-1, Clontech).). As described in the manual accompanying the Matchmaker Two Hybrid System 2 (Catalog No. K1604-1, Clontech), the disclosure of which is incorporated herein by reference, nucleic acids encoding the TBC-1 protein or a portion thereof, are inserted into an expression vector such that they are in frame with DNA encoding the DNA

binding domain of the yeast transcriptional activator GAL4. A desired cDNA, preferably human cDNA, is inserted into a second expression vector such that they are in frame with DNA encoding the activation domain of GAL4. The two expression plasmids are transformed into the yeast cells and the yeast cells are plated on selection medium which selects for expression of selectable markers on each of
5 the expression vectors as well as GAL4 dependent expression of the HIS3 gene. Transformants capable of growing on medium lacking histidine are screened for GAL4 dependent lacZ expression. Those cells which are positive in both the histidine selection and the lacZ assay are those in which an interaction between TBC-1 and the protein or peptide encoded by the initially selected cDNA insert has taken place.

10 **Method for screening ligands that modulate the expression of the *TBC-1* gene.**

Another subject of the present invention is a method for screening molecules that modulate the expression of the TBC-1 protein. Such a screening method comprises the steps of :

- a) cultivating a prokaryotic or an eukaryotic cell that has been transfected with a nucleotide sequence encoding the TBC-1 protein, operably linked to a *TBC-1* 5'-regulatory sequence;
- 15 b) bringing into contact the cultivated cell with a molecule to be tested;
- c) quantifying the expression of the TBC-1 protein.

Using DNA recombination techniques well known by the one skill in the art, the TBC-1 protein encoding DNA sequence is inserted into an expression vector, downstream from a *TBC-1* 5'-regulatory sequence that contains a *TBC-1* promoter sequence.

- 20 The quantification of the expression of the TBC-1 protein may be realized either at the mRNA level or at the protein level. In the latter case, polyclonal or monoclonal antibodies may be used to quantify the amounts of the TBC-1 protein that have been produced, for example in an ELISA or a RIA assay.

In a preferred embodiment, the quantification of the *TBC-1* mRNAs is realized by a
25 quantitative PCR amplification of the cDNAs obtained by a reverse transcription of the total mRNA of the cultivated *TBC-1*-transfected host cell, using a pair of primers specific for *TBC-1*.

Expression levels and patterns of *TBC-1* may be analyzed by solution hybridization with long probes as described in International Patent Application No. WO 97/05277, the entire contents of which are incorporated herein by reference. Briefly, the *TBC-1* cDNA or the *TBC-1* genomic DNA described
30 above, or fragments thereof, is inserted at a cloning site immediately downstream of a bacteriophage (T3, T7 or SP6) RNA polymerase promoter to produce antisense RNA. Preferably, the *TBC-1* insert comprises at least 100 or more consecutive nucleotides of the genomic DNA sequence or the cDNA sequences, particularly those comprising one of the nucleotide sequences of SEQ ID Nos 3, 4 and 6-8 or those encoding a mutated TBC-1. The plasmid is linearized and transcribed in the presence of
35 ribonucleotides comprising modified ribonucleotides (i.e. biotin-UTP and DIG-UTP). An excess of this doubly labeled RNA is hybridized in solution with mRNA isolated from cells or tissues of interest. The hybridizations are performed under standard stringent conditions (40-50°C for 16 hours in an 80%

formamide, 0.4 M NaCl buffer, pH 7-8). The unhybridized probe is removed by digestion with ribonucleases specific for single-stranded RNA (i.e. RNases CL3, T1, Phy M, U2 or A). The presence of the biotin-UTP modification enables capture of the hybrid on a microtitration plate coated with streptavidin. The presence of the DIG modification enables the hybrid to be detected and quantified by

5 ELISA using an anti-DIG antibody coupled to alkaline phosphatase.

Quantitative analysis of *TBC-1* gene expression may also be performed using arrays. As used herein, the term array means a one dimensional, two dimensional, or multidimensional arrangement of a plurality of nucleic acids of sufficient length to permit specific detection of expression of mRNAs capable of hybridizing thereto. For example, the arrays may contain a
10 plurality of nucleic acids derived from genes whose expression levels are to be assessed. The arrays may include the *TBC-1* genomic DNA, the *TBC-1* cDNA sequences or the sequences complementary thereto or fragments thereof, particularly those comprising at least one of the biallelic markers according to the present invention. Preferably, the fragments are at least 15 nucleotides in length. In other embodiments, the fragments are at least 25 nucleotides in length. In
15 some embodiments, the fragments are at least 50 nucleotides in length. More preferably, the fragments are at least 100 nucleotides in length. In another preferred embodiment, the fragments are more than 100 nucleotides in length. In some embodiments the fragments may be more than 500 nucleotides in length.

For example, quantitative analysis of *TBC-1* gene expression may be performed with a
20 complementary DNA microarray as described by Schena et al. (1995). Full length *TBC-1* cDNAs or fragments thereof are amplified by PCR and arrayed from a 96-well microtiter plate onto silylated microscope slides using high-speed robotics. Printed arrays are incubated in a humid chamber to allow rehydration of the array elements and rinsed, once in 0.2% SDS for 1 min, twice in water for 1 min and once for 5 min in sodium borohydride solution. The arrays are submerged in water for 2
25 min at 95°C, transferred into 0.2% SDS for 1 min, rinsed twice with water, air dried and stored in the dark at 25°C.

Cell or tissue mRNA is isolated or commercially obtained and probes are prepared by a single round of reverse transcription. Probes are hybridized to 1 cm² microarrays under a 14 x 14 mm glass coverslip for 6-12 hours at 60°C. Arrays are washed for 5 min at 25°C in low stringency
30 wash buffer (1 x SSC/0.2% SDS), then for 10 min at room temperature in high stringency wash buffer (0.1 x SSC/0.2% SDS). Arrays are scanned in 0.1 x SSC using a fluorescence laser scanning device fitted with a custom filter set. Accurate differential expression measurements are obtained by taking the average of the ratios of two independent hybridizations.

Quantitative analysis of *TBC-1* gene expression may also be performed with full length
35 *TBC-1* cDNAs or fragments thereof in complementary DNA arrays as described by Pietu et al. (1996). The full length *TBC-1* cDNA or fragments thereof is PCR amplified and spotted on membranes. Then, mRNAs originating from various tissues or cells are labeled with radioactive

nucleotides. After hybridization and washing in controlled conditions, the hybridized mRNAs are detected by phospho-imaging or autoradiography. Duplicate experiments are performed and a quantitative analysis of differentially expressed mRNAs is then performed.

Alternatively, expression analysis using the *TBC-1* genomic DNA, the *TBC-1* cDNAs, or
5 fragments thereof can be done through high density nucleotide arrays or chips as described by Lockhart et al. (1996) and Sosnowsky et al. (1997). Oligonucleotides of 15-50 nucleotides from the sequences of the *TBC-1* genomic DNA, the *TBC-1* cDNA sequences particularly those comprising at least one of biallelic markers according the present invention, preferably at least one of SEQ ID No 7-8 or those comprising the trait causing mutation, or the sequences complementary thereto, are
10 synthesized directly on the chip (Lockhart et al., supra) or synthesized and then addressed to the chip (Sosnowski et al., supra). Preferably, the oligonucleotides are about 20 nucleotides in length.

TBC-1 cDNA probes labeled with an appropriate compound, such as biotin, digoxigenin or fluorescent dye, are synthesized from the appropriate mRNA population and then randomly fragmented to an average size of 50 to 100 nucleotides. The said probes are then hybridized to the
15 chip. After washing as described in Lockhart et al., supra and application of different electric fields (Sosnowsky et al., 1997), the dyes or labeling compounds are detected and quantified. Duplicate hybridizations are performed. Comparative analysis of the intensity of the signal originating from cDNA probes on the same target oligonucleotide in different cDNA samples indicates a differential expression of *TBC-1* mRNAs.

20 Thus, is also part of the present invention a method for screening of a candidate substance or molecule that modulates the expression of the *TBC-1* gene according to the invention, wherein this method comprises the following steps :

- a) providing a recombinant cell host containing a nucleic acid, wherein said nucleic acid comprises the 5' regulatory region sequence or a biologically active fragment or variant thereof, the
25 5' regulatory region or its biologically active fragment or variant being operably linked to a polynucleotide encoding a detectable protein;
- b) obtaining a candidate substance, and
- c) determining the ability of the candidate substance to modulate the expression levels of the polynucleotide encoding the detectable protein.

30 In a preferred embodiment of the above screening method, the nucleic acid comprising the 5' regulatory region sequence or a biologically active fragment or variant thereof also includes a 5'UTR region of one of the *TBC-1* cDNAs of SEQ ID Nos 3 and 4, or one of their biologically active fragments or variants thereof.

A second method for the screening of a candidate substance or molecule that modulates the
35 expression of the *TBC-1* gene comprises the following steps :

- a) providing a recombinant cell host containing a nucleic acid, wherein said nucleic acid comprises a 5'UTR sequence of one of the *TBC-1* cDNAs of SEQ ID Nos 3 and 4, or one of their

biologically active fragments or variants, the 5'UTR sequence or its biologically active fragment or variant being operably linked to a polynucleotide encoding a detectable protein;

b) obtaining a candidate substance, and

c) determining the ability of the candidate substance to modulate the expression levels of the polynucleotide encoding the detectable protein.

In a preferred embodiment of the screening method described above, the nucleic acid that comprises a nucleotide sequence selected from the group consisting of the 5'UTR sequence of one of the *TBC-1* cDNAs of SEQ ID Nos 3 and 4 or one of their biologically active fragments or variants, includes a promoter sequence, wherein said promoter sequence can be either endogenous, or in contrast exogenous with respect to the *TBC-1* 5'UTR sequences defined therein.

Among the preferred polynucleotides encoding a detectable protein, there may be cited polynucleotides encoding beta galactosidase, green fluorescent protein (GFP) and chloramphenicol acetyl transferase (CAT).

For the design of suitable recombinant vectors useful for performing the screening methods described above, it will be referred to the section of the present specification wherein the preferred recombinant vectors of the invention are detailed.

Screening using transgenic animals

In vivo methods can utilize transgenic animals for drug screening. Nucleic acids including at least one of the biallelic polymorphisms of interest can be used to generate genetically modified non-human animals or to generate site specific gene modifications in cell lines. The term "transgenic" is intended to encompass genetically modified animals having a deletion or other knock-out of *TBC-1* gene activity, having an exogenous *TBC-1* gene that is stably transmitted in the host cells, or having an exogenous *TBC-1* promoter operably linked to a reporter gene. Transgenic animals may be made through homologous recombination, where the *TBC-1* locus is altered. Alternatively, a nucleic acid construct is randomly integrated into the genome. Vectors for stable integration include for example plasmids, retroviruses and other animal viruses, and YACs. Of interest are transgenic mammals e.g. cows, pigs, goats, horses, and particularly rodents such as rats and mice. Transgenic animals allow to study both efficacy and toxicity of the candidate drug.

Methods for inhibiting the expression of a *TBC-1* gene

Other therapeutic compositions according to the present invention comprise advantageously an oligonucleotide fragment of the nucleic sequence of *TBC-1* as an antisense tool that inhibits the expression of the corresponding *TBC-1* gene. Preferred methods using antisense polynucleotide according to the present invention are the procedures described by Sczakiel et al. (1995).

Preferably, the antisense tools are chosen among the polynucleotides (15-200 bp long) that are complementary to the 5' end of the *TBC-1* mRNA. In another embodiment, a combination of different antisense polynucleotides complementary to different parts of the desired targetted gene are used.

Preferred antisense polynucleotides according to the present invention are complementary to a sequence of the mRNAs of *TBC-1* that contains the translation initiation codon ATG.

The antisense nucleic acid molecules to be used in gene therapy may be either DNA or RNA sequences. They comprise a nucleotide sequence complementary to the targeted sequence of the PTCA-1 genomic DNA, the sequence of which can be determined using one of the detection methods of the present invention. The targeted DNA or RNA sequence preferably comprises at least one of the biallelic markers according to the present invention. The antisense nucleic acids should have a length and melting temperature sufficient to permit formation of an intracellular duplex having sufficient stability to inhibit the expression of the *TBC-1* mRNA in the duplex. Strategies for designing antisense nucleic acids suitable for use in gene therapy are disclosed in Green et al., (1986) and Izant and Weintraub, (1984), the disclosures of which are incorporated herein by reference.

In some strategies, antisense molecules are obtained by reversing the orientation of the *TBC-1* coding region with respect to a promoter so as to transcribe the opposite strand from that which is normally transcribed in the cell. The antisense molecules may be transcribed using in vitro transcription systems such as those which employ T7 or SP6 polymerase to generate the transcript. Another approach involves transcription of *TBC-1* antisense nucleic acids in vivo by operably linking DNA containing the antisense sequence to a promoter in a suitable expression vector.

Alternatively, suitable antisense strategies are those described by Rossi et al. (1991), in the International Applications Nos. WO 94/23026, WO 95/04141, WO 92/18522 and in the European Patent Application No. EP 0 572 287 A2

An alternative to the antisense technology that is used according to the present invention consists in using ribozymes that will bind to a target sequence via their complementary polynucleotide tail and that will cleave the corresponding RNA by hydrolyzing its target site (namely « hammerhead ribozymes »). Briefly, the simplified cycle of a hammerhead ribozyme consists of (1) sequence specific binding to the target RNA via complementary antisense sequences; (2) site-specific hydrolysis of the cleavable motif of the target strand; and (3) release of cleavage products, which gives rise to another catalytic cycle. Indeed, the use of long-chain antisense polynucleotide (at least 30 bases long) or ribozymes with long antisense arms are advantageous. A preferred delivery system for antisense ribozyme is achieved by covalently linking these antisense ribozymes to lipophilic groups or to use liposomes as a convenient vector. Preferred antisense ribozymes according to the present invention are prepared as described by Sczakiel et al. (1995), the specific preparation procedures being referred to in said article being herein incorporated by reference.

35

Throughout this application, various publications, patents and published patent applications are cited. The disclosures of these publications, patents and published patent specification

referenced in this application are hereby incorporated by reference into the present disclosure to more fully describe the state of the art to which this invention pertains.

EXAMPLES

EXAMPLE 1 :

5 **Analysis of the first mRNA encoding a TBC-1 polypeptide synthesized by the cells.**

TBC-1 cDNA was obtained as follows : 4 µl of ethanol suspension containing 1 mg of human prostate total RNA (Clontech laboratories, Inc., Palo Alto, USA; Catalogue N. 64038-1) was centrifuged, and the resulting pellet was air dried for 30 minutes at room temperature.

First strand cDNA synthesis was performed using the Advantage™ RT-for-PCR kit
10 (Clontech laboratories Inc., catalogue N. K1402-1). 1 µl of 20 mM solution of a specific oligo dT primer was added to 12.5 µl of RNA solution in water, heated at 74°C for 2.5 min and rapidly quenched in an ice bath. 10 µl of 5 x RT buffer (50 mM Tris-HCl, pH 8.3, 75 mM KCl, 3 mM MgCl₂), 2.5 µl of dNTP mix (10 mM each), 1.25 µl of human recombinant placental RNA inhibitor were mixed with 1 ml of MMLV reverse transcriptase (200 units). 6.5 µl of this solution were
15 added to RNA-primer mix and incubated at 42°C for one hour. 80 µl of water were added and the solution was incubated at 94°C for 5 minutes.

5 µl of the resulting solution were used in a Long Range PCR reaction with hot start, in 50 µl final volume, using 2 units of rtTHXL, 20 pmol/µl of each of 5'-TGACCACCATGCCCCATGCT-3' (271-289 in SEQ ID No 3) and 5'-
20 GCATTATTCACGTCCACGCC-3' (3929-3949 in SEQ ID No 3) primers with 35 cycles of elongation for 6 minutes at 67°C in thermocycler.

The amplification products corresponding to both cDNA strands were partially sequenced in order to ensure the specificity of the amplification reaction.

Results of Northern blot analysis of prostate mRNAs supported the existence of the first
25 *TBC-1* cDNA having about 4 kb in length, which is the nucleotide sequence of SEQ ID No 3.

Example 2 :

Detection of *TBC-1* biallelic markers: DNA extraction

Donors were unrelated and healthy. They presented a sufficient diversity for being representative of a French heterogeneous population. The DNA from 100 individuals was extracted
30 and tested for the detection of the biallelic markers.

30 ml of peripheral venous blood were taken from each donor in the presence of EDTA. Cells (pellet) were collected after centrifugation for 10 minutes at 2000 rpm. Red cells were lysed by a lysis solution (50 ml final volume : 10 mM Tris pH7.6; 5 mM MgCl₂; 10 mM NaCl). The

solution was centrifuged (10 minutes, 2000 rpm) as many times as necessary to eliminate the residual red cells present in the supernatant, after resuspension of the pellet in the lysis solution.

The pellet of white cells was lysed overnight at 42°C with 3.7 ml of lysis solution composed of:

- 5 - 3 ml TE 10-2 (Tris-HCl 10 mM, EDTA 2 mM) / NaCl 0.4 M
- 200 µl SDS 10%
- 500 µl K-proteinase (2 mg K-proteinase in TE 10-2 / NaCl 0.4 M).

For the extraction of proteins, 1 ml saturated NaCl (6M) (1/3.5 v/v) was added. After vigorous agitation, the solution was centrifuged for 20 minutes at 10000 rpm.

- 10 For the precipitation of DNA, 2 to 3 volumes of 100% ethanol were added to the previous supernatant, and the solution was centrifuged for 30 minutes at 2000 rpm. The DNA solution was rinsed three times with 70% ethanol to eliminate salts, and centrifuged for 20 minutes at 2000 rpm. The pellet was dried at 37°C, and resuspended in 1 ml TE 10-1 or 1 ml water. The DNA concentration was evaluated by measuring the OD at 260 nm (1 unit OD = 50 µg/ml DNA).

- 15 To determine the presence of proteins in the DNA solution, the OD 260 / OD 280 ratio was determined. Only DNA preparations having a OD 260 / OD 280 ratio between 1.8 and 2 were used in the subsequent examples described below.

The pool was constituted by mixing equivalent quantities of DNA from each individual.

Example 3 :

20 **Detection of the biallelic markers: amplification of genomic DNA by PCR**

The amplification of specific genomic sequences of the DNA samples of example 2 was carried out on the pool of DNA obtained previously. In addition, 50 individual samples were similarly amplified.

PCR assays were performed using the following protocol:

25	Final volume	25 µl
	DNA	2 ng/µl
	MgCl ₂	2 mM
	dNTP (each)	200 µM
	primer (each)	2.9 ng/µl
30	Ampli Taq Gold DNA polymerase	0.05 unit/µl
	PCR buffer (10x = 0.1 M TrisHCl pH8.3 0.5M KCl	1x

- Each pair of first primers was designed using the sequence information of the *TBC-1* gene disclosed herein and the OSP software (Hillier & Green, 1991). This first pair of primers was about 20 nucleotides in length and had the sequences disclosed in Table 1 in the columns labeled PU and
- 35 RP.

Table 1

Amplicon	Position range of the amplicon in SEQ ID 1		Primer name	Position range of amplification primer in SEQ ID No 1		Primer name	Complementary position range of amplification primer in SEQ ID No 1	
99-430	9391	9845	B1	9391	9408	C1	9828	9845
Amplicon	Position range of the amplicon in SEQ ID 2		Primer name	Position range of amplification primer in SEQ ID No 2		Primer name	Complementary position range of amplification primer in SEQ ID No 2	
99-20508	988	1529	B2	988	1006	C2	1509	1529
99-20469	5039	5554	B3	5039	5056	C3	5534	5554
5-254	5997	6350	B4	5997	6015	C4	6332	6350
5-257	14371	14817	B5	14371	14390	C5	14798	14817
99-20511	18751	19217	B6	18751	18771	C6	19198	19217
99-20510	19605	20005	B7	19605	19625	C7	19986	20005
99-20504	29529	30061	B8	29529	29547	C8	30041	30061
99-20493	42268	42752	B9	42268	42287	C9	42732	42752
99-20499	69026	69543	B10	69026	69046	C10	69525	69543
99-20473	76323	76790	B11	76323	76343	C11	76771	76790
5-249	78292	78721	B12	78292	78309	C12	78704	78721
99-20485	81893	82372	B13	81893	81912	C13	82353	82372
99-20481	84392	84929	B14	84392	84412	C14	84909	84929
99-20480	89746	90198	B15	89746	89765	C15	90179	90198

Preferably, the primers contained a common oligonucleotide tail upstream of the specific bases targeted for amplification which was useful for sequencing.

5 Primers PU contain the following additional PU 5' sequence :

TGTAAAACGACGGCCAGT (SEQ ID No 6); primers RP contain the following RP 5' sequence : CAGGAAACAGCTATGACC (SEQ ID No 7).

The synthesis of these primers was performed following the phosphoramidite method, on a GENSET UFPS 24.1 synthesizer.

10 DNA amplification was performed on a Genius II thermocycler. After heating at 95°C for 10 min, 40 cycles were performed. Each cycle comprised: 30 sec at 95°C, 54°C for 1 min, and 30 sec at 72°C. For final elongation, 10 min at 72°C ended the amplification. The quantities of the amplification products obtained were determined on 96-well microtiter plates, using a fluorometer and Picogreen as intercalant agent (Molecular Probes).

15

Example 4 :

Detection of the biallelic markers: sequencing of amplified genomic DNA and identification of polymorphisms.

The sequencing of the amplified DNA obtained in example 3 was carried out on ABI 377 sequencers. The sequences of the amplification products were determined using automated dideoxy
20 terminator sequencing reactions with a dye terminator cycle sequencing protocol. The products of

the sequencing reactions were run on sequencing gels and the sequences were determined using gel image analysis [ABI Prism DNA Sequencing Analysis software (2.1.2 version)].

The sequence data were further evaluated to detect the presence of biallelic markers among the pooled amplified fragments. The polymorphism search was based on the presence of
5 superimposed peaks in the electrophoresis pattern resulting from different bases occurring at the same position as described previously.

15 fragments of amplification was analyzed. In this segment, 19 biallelic markers were detected. The localization of the biallelic marker is as shown in Table 2.

Table 2

Amplicon	BM	Marker Name	Localization in <i>TBC-1</i> gene	Polymorphism		BM position in SEQ ID No 1
				Allele 1	allele 2	
99-430	A1	99-430-352	Intron 1	A	G	9494
Amplicon	BM	Marker Name	Localization in <i>TBC-1</i> gene	Polymorphism		BM position in SEQ ID No 2
				allele 1	allele 2	
99-20508	A2	99-20508-456	Intron upstream to Exon A	C	T	1443
99-20469	A3	99-20469-213	Intron A	C	T	5247
5-254	A4	5-254-227	Intron B	A	G	6223
5-257	A5	5-257-353	Intron D	C	T	14723
99-20511	A6	99-20511-32	Intron D	C	T	19186
99-20511	A7	99-20511-221	Intron D	A	G	18997
99-20510	A8	99-20510-115	Intron D	deletion of TCT		19891
99-20504	A9	99-20504-90	Intron D	A	G	29617
99-20493	A10	99-20493-238	Intron D	A	C	42519
99-20499	A11	99-20499-221	Intron G	A	G	69324
99-20499	A12	99-20499-364	Intron G	A	T	69181
99-20499	A13	99-20499-399	Intron G	A	G	69146
99-20473	A14	99-20473-138	Intron H	deletion of TAACA		76458
5-249	A15	5-249-304	Intron I	A	G	78595
99-20485	A16	99-20485-269	Intron I	A	G	82159
99-20481	A17	99-20481-131	Intron I	G	C	84522
99-20481	A18	99-20481-419	Intron I	A	T	84810
99-20480	A19	99-20480-233	Intron J	A	G	89967

10 BM refers to "biallelic marker". All1 and all2 refer respectively to allele 1 and allele 2 of the biallelic marker.

Table 3

BM	Marker Name	Position range of probes in SEQ ID No 1		Probes
A1	99-430-352	9482	9506	P1

BM	Marker Name	Position range of probes in SEQ ID No 2		Probes
A2	99-20508-456	1431	1455	P2
A3	99-20469-213	5235	5259	P3
A4	5-254-227	6211	6235	P4
A5	5-257-353	14711	14735	P5
A6	99-20511-32	19174	19198	P6
A7	99-20511-221	18985	19009	P7
A9	99-20504-90	29605	29629	P9
A10	99-20493-238	42507	42531	P10
A11	99-20499-221	69312	69336	P11
A12	99-20499-364	69169	69193	P12
A13	99-20499-399	69134	69158	P13
A15	5-249-304	78583	78607	P15
A16	99-20485-269	82147	82171	P16
A17	99-20481-131	84510	84534	P17
A18	99-20481-419	84798	84822	P18
A19	99-20480-233	89955	89979	P19

Example 5 :**Validation of the polymorphisms through microsequencing**

The biallelic markers identified in example 4 were further confirmed and their respective
5 frequencies were determined through microsequencing. Microsequencing was carried out for each individual DNA sample described in Example 2.

Amplification from genomic DNA of individuals was performed by PCR as described
above for the detection of the biallelic markers with the same set of PCR primers (Table 1).

The preferred primers used in microsequencing were about 19 nucleotides in length and
10 hybridized just upstream of the considered polymorphic base. According to the invention, the primers used in microsequencing are detailed in Table 4.

Table 4

Marker Name	Biallelic Marker	Mis. 1	Position range of microsequencing primer mis 1 in SEQ ID No 1		Mis. 2	Complementary position range of microsequencing primer mis. 2 in SEQ ID No 1	
99-430-352	A1	D1	9475	9493	E1	9495	9513
Marker Name	Biallelic Marker	Mis. 1	Position range of microsequencing primer mis 1 in SEQ ID No 2		Mis. 2	Complementary position range of microsequencing primer mis. 2 in SEQ ID No 2	
99-20508-456	A2	D2	1424	1442	E2	1444	1462
99-20469-213	A3	D3	5228	5246	E3	5248	5266
5-254-227	A4	D4	6204	6222	E4	6224	6242
5-257-353	A5	D5	14704	14722	E5	14724	14742
99-20511-32	A6	D6	19167	19185	E6	19187	19205
99-20511-221	A7	D7	18978	18996	E7	18998	19016

99-20510-115	A8	D8	19872	19890	E8	19892	19910
99-20504-90	A9	D9	29598	29616	E9	29618	29636
99-20493-238	A10	D10	42500	42518	E10	42520	42538
99-20499-221	A11	D11	69305	69323	E11	69325	69343
99-20499-364	A12	D12	69162	69180	E12	69182	69200
99-20499-399	A13	D13	69127	69145	E13	69147	69165
99-20473-138	A14	D14	76439	76457	E14	76459	76477
5-249-304	A15	D15	78576	78594	E15	78596	78614
99-20485-269	A16	D16	82140	82158	E16	82160	82178
99-20481-131	A17	D17	84503	84521	E17	84523	84541
99-20481-419	A18	D18	84791	84809	E18	84811	84829
99-20480-233	A19	D19	89948	89966	E19	89968	89986

The microsequencing reaction was performed as follows :

- After purification of the amplification products, the microsequencing reaction mixture was prepared by adding, in a 20µl final volume: 10 pmol microsequencing oligonucleotide, 1 U Thermosequenase (Amersham E79000G), 1.25 µl Thermosequenase buffer (260 mM Tris HCl pH 9.5, 65 mM MgCl₂), and the two appropriate fluorescent ddNTPs (Perkin Elmer, Dye Terminator Set 401095) complementary to the nucleotides at the polymorphic site of each biallelic marker tested, following the manufacturer's recommendations. After 4 minutes at 94°C, 20 PCR cycles of 15 sec at 55°C, 5 sec at 72°C, and 10 sec at 94°C were carried out in a Tetrad PTC-225 thermocycler (MJ Research). The unincorporated dye terminators were then removed by ethanol precipitation. Samples were finally resuspended in formamide-EDTA loading buffer and heated for 2 min at 95°C before being loaded on a polyacrylamide sequencing gel. The data were collected by an ABI PRISM 377 DNA sequencer and processed using the GENESCAN software (Perkin Elmer).

- Following gel analysis, data were automatically processed with software that allows the determination of the alleles of biallelic markers present in each amplified fragment.

- The software evaluates such factors as whether the intensities of the signals resulting from the above microsequencing procedures are weak, normal, or saturated, or whether the signals are ambiguous. In addition, the software identifies significant peaks (according to shape and height criteria). Among the significant peaks, peaks corresponding to the targeted site are identified based on their position. When two significant peaks are detected for the same position, each sample is categorized classification as homozygous or heterozygous type based on the height ratio.

References

- Altschul et al., 1990, J. Mol. Biol. 215(3):403-410 / Altschul et al., 1993, Nature Genetics 3:266-272 / Altschul et al., 1997, Nuc. Acids Res. 25:3389-3402 / Ausubel et al. (1989) Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y. / Beaucage et al., *Tetrahedron Lett* 1981, 22: 1859-1862 / Bram RJ et al., 1993, Mol. Cell Biol., 13 : 4760-4769. / Brown EL, Belagaje R, Ryan MJ, Khorana HG, *Methods Enzymol* 1979;68:109-151 / Castagnoli L. et al. (Felici F.), 1991, J. Mol. Biol., 222:301-310. /

- Chai H. et al., 1993, *Biotechnol. Appl. Biochem.*, **18**:259-273 / Chee et al. (1996) *Science*. 274:610-614. / Chen and Kwok *Nucleic Acids Research* 25:347-353 1997 / Chen et al. *Proc. Natl. Acad. Sci. USA* 94/20 10756-10761, 1997 / Cho RJ et al., 1998, *Proc. Natl. Acad. Sci. USA*, **95**(7) : 3752-3757. / Chumakov I. et al., 1995, *Nature*, **377**(6547 Suppl): 175-297. / Compton J. 5 (1991) *Nature*. 350(6313):91-92. / Dib et al., 1996, *Nature*, **380**: III-V. / Ellis NA, 1997 *Curr. Op. Genet. Dev.*, **7** : 354-363 / Feldman and Steg, 1996, *Medecine/Sciences, synthese*, **12**:47-55 / Fields and Song, 1989, *Nature*, Vol. 340 : 245-246. / Fishel R & Wilson T. 1997, *Curr. Op. Genet. Dev.* **7**: 105-113 / Flotte et al., 1992, *Am. J. Respir. Cell Mol. Biol.*, **7** : 349-356. / Fodor et al. (1991) *Science* 251:767-777. / Fromont-Racine M. et al., 1997, *Nature Genetics*, 10 **16**(3) : 277-282. / Fuller S.A. et al., 1996, *Immunology in Current Protocols in Molecular Biology*, Ausubel et al. Eds, John Wiley & Sons, Inc., USA / Geysen H. Mario et al. 1984. *Proc. Natl. Acad. Sci. U.S.A.* **81**:3998-4002 / Gonnet et al., 1992, *Science* 256:1443-1445 / Green et al., *Ann. Rev. Biochem.* **55**:569-597 (1986) / Grompe, M. et al., *Proc. Natl. Acad. Sci. U.S.A* 1989; **86**:5855-5892 / Grompe, M. *Nature Genetics* 1993; **5**:111-117 / Guatelli J C et al. *Proc. Natl. Acad. Sci. USA*. **35**:273-286. / Haber D & Harlow E, 1997, *Nature Genet.* **16**:320-322. / Hacia 15 JG, Brody LC, Chee MS, Fodor SP, Collins FS, *Nat Genet* 1996;**14**(4):441-447 / Haff L. A. and Smirnov I. P. (1997) *Genome Research*, **7**:378-388. / Hames B.D. and Higgins S.J. (1985) *Nucleic Acid Hybridization: A Practical Approach*. Hames and Higgins Ed., IRL Press, Oxford. / Harju L, et al., *Clin Chem* 1993;**39**(11Pt 1):2282-2287 / Harper JW et al., 1993, *Cell*, Vol. **75** : 20 805-816. / Harris H et al., 1969, *Nature* 223:363-368. / Henikoff and Henikoff, 1993, *Proteins* **17**:49-61 / Higgins et al., 1996, *Methods Enzymol.* **266**:383-402 / Hillier L. and Green P. *Methods Appl.*, 1991, **1**: 124-8. / Huang L. et al. (1996) *Cancer Res* **56**(5):1137-1141. / Huygen et al., 1996, *Nature Medicine*, **2**(8):893-898 / Izant and Weintraub, *Cell* **36**:1007-1015 (1984) / Julian et al., 1992, *J. Gen. Virol.*, **73** : 3251 – 3255. / Karlin and Altschul, 1990, *Proc. Natl. Acad. Sci. USA* **87**:2267-2268 / Koch Y., 1977, *Biochem. Biophys. Res. Commun.*, **74**:488-491 / 25 Kohler G. and Milstein C., 1975, *Nature*, **256** : 495. / Kozal MJ, et al., *Nat Med* 1996;**2**(7):753-759 / Landegren U. et al. (1998) *Genome Research*, **8**:769-776. / Leger OJ, et al., 1997, *Hum Antibodies*, **8**(1): 3-16 / Lenhard T. et al., 1996, *Gene*, **169**:187-190 / Livak et al., *Nature Genetics*, **9**:341-342, 1995 / Livak KJ, and Hainer JW., 1994, *Hum Mutat.*, **3**(4): 379-385. / 30 Lockhart et al. *Nature Biotechnology* **14**: 1675-1680, 1996 / Lucas A.H., 1994, In : *Development and Clinical Uses of Haemophilus b Conjugate*. / Mansour SL et al., 1988, *Nature*, **336** : 348-352. / Marshall R. L. et al. (1994) *PCR Methods and Applications*. **4**:80-84. / Martineau P, Jones P, Winter G, 1998, *J Mol Biol*, **280**(1):117-127 / Mc Whorter W.P., et al. A screening study of prostate cancer in high risk families. *J Urol* 1992;**148**:826-828. / McLaughlin et al., 1989, *J. Virol.*, 35 **62** : 1963 – 1973. / Muzyczka et al., 1992, *Curr. Topics in Micro. and Immunol.*, **158** : 97-129. / Narang SA, Hsiung HM, Brousseau R, *Methods Enzymol* 1979;**68**:90-98 / Neda et al., 1991, *J. Biol. Chem.*, **266** : 14143 – 14146. / Nickerson D.A. et al. (1990) *Proc. Natl. Acad. Sci. U.S.A.*

- 87:8923-8927. / Nyren P, Pettersson B, Uhlen M, *Anal Biochem* 1993;**208**(1):171-175 / O'Reilly et al., 1992, *Baculovirus expression vectors : a Laboratory Manual*. W.H. Freeman and Co., New York / Ohno et al., 1994, *Sciences*, **265**:781-784 / Oldenburg K.R. et al., 1992, *Proc. Natl. Acad. Sci.*, **89**:5393-5397. / Orita et al., *Proc. Natl. Acad. Sci. U.S.A.* 1989;**86**: 2776-2770 /
- 5 Parmley and Smith, *Gene*, 1988, **73**:305-318. / Pastinen et al., *Genome Research* 1997; **7**:606-614 / *PCR Methods and Applications*", 1991, Cold Spring Harbor Laboratory Press. / Pearson and Lipman, 1988, *Proc. Natl. Acad. Sci. USA* **85**(8):2444-2448 / Pietu et al. *Genome Research* **6**:492-503, 1996 / Porath J et al., 1975, *Nature*, **258**(5536) : 598-599. / Reimann KA, et al., 1997, *AIDS Res Hum Retroviruses*. **13**(11): 933-943 / Ridder R, et al., 1995, *Biotechnology (N Y)*,
- 10 **13**(3):255-260 / Rossi et al., *Pharmacol. Ther.* **50**:245-254, (1991) / Roth J.A. et al., 1996, *Nature Medicine*, **2**(9):985-991 / Rougeot, C. et al., *Eur. J. Biochem.* **219** (3): 765-773, 1994 / Roux et al., 1989, *Proc. Natl Acad. Sci. USA*, **86** : 9079 – 9083. / Sambrook, et al. 1989. *Molecular cloning: a laboratory manual*. 2ed. Cold Spring Harbor Laboratory, Cold spring Harbor, New York. / Samson M, et al. (1996) *Nature*, **382**(6593):722-725. / Samulski et al., 1989, *J. Virol.*, **63** :
- 15 3822-3828. / Sanchez-Pescador R., 1988, *J. Clin. Microbiol.*, **26**(10):1934-1938 / Sandou et al., 1994, *Science*, **265** : 1875-1878. / Schena et al. *Science* **270**:467-470, 1995 / Schwartz and Dayhoff, eds., 1978, *Matrices for Detecting Distance Relationships: Atlas of Protein Sequence and Structure*, Washington: National Biomedical Research Foundation / Sczakiel G. et al., 1995, *Trends Microbiol.*, 1995, **3**(6):213-217 / Sheffield, V.C. et al, *Proc. Natl. Acad. Sci. U.S.A* 1991;
- 20 **49**:699-706 / Shoemaker DD, et al., *Nat Genet* 1996;**14**(4):450-456 / Smith et al., 1983, *Mol. Cell. Biol.*, **3**:2156-2165. / Sosnowski RG, et al., *Proc Natl Acad Sci U S A* 1997;**94**:1119-1123 / Steinberg G.D., et al. *Family history and the risk of prostate cancer, The prostate* 1990;**17**,337-347. / Stryer, L., *Biochemistry*, 4th edition, 1995 / Syvanen AC, et al., 1994, *Hum Mutat.*, **3**(3): 172-179. / Tacson et al., 1996, *Nature Medicine*, **2**(8):888-892. / Thompson et al., 1994, *Nucleic*
- 25 *Acids Res.* **22**(2):4673-4680 / Tyagi et al. (1998) *Nature Biotechnology*. **16**:49-53. / Urdea M.S., 1988, *Nucleic Acids Research*, **11**: 4937-4957 / Urdea MS et al., 1991, *Nucleic Acids Symp Ser.*, **24**: 197-200. / Valadon P., et al., 1996, *J. Mol. Biol.*, Vol. **261**:11-22. / Vaughan TJ, et al., 1996, *Nat Biotechnol.* **14**(3): 309-314 / Vlasak R. et al., 1983, *Eur. J. Biochem.*, **135**:123-126 / Wabiko et al., 1986, *DNA*, **5**(4):305-314. / Walker et al. (1996) *Clin. Chem.* **42**:9-13. /
- 30 Westerink M.A.J., 1995, *Proc. Natl. Acad. Sci.*, **92**:4021-4025. / White, M.B. et al. (1992) *Genomics*. **12**:301-306. / White, M.B. et al. (1997) *Genomics*. **12**:301-306. / Wilson R. et al., 1994, *Nature*, **368**(6466) : 32-38. / Zhang SD et al., 1996, *Genes and development*, **10** : 1108-1119.

SEQUENCE LISTING FREE TEXT

- 35 The following free text appears in the accompanying Sequence Listing :
5' regulatory region

polymorphic base

complement

3' regulatory region

deletion of

5

or

probe

homology with Genset 5' EST in ref

sequencing oligonucleotide PrimerPU

sequencing oligonucleotide PrimerRP

What is claimed is :

1. An isolated, purified, or recombinant polynucleotide comprising a contiguous span of at least 12 nucleotides of SEQ ID No 1 or the complements thereof.

5

2. An isolated, purified, or recombinant polynucleotide comprising a contiguous span of at least 12 nucleotides of SEQ ID No 2 or the complements thereof.

3. An isolated, purified, or recombinant polynucleotide comprising a contiguous span of at least 12 nucleotides of SEQ ID No 3 or the complements thereof.

10

4. An isolated, purified, or recombinant polynucleotide comprising a contiguous span of at least 12 nucleotides of SEQ ID No 4 or the complements thereof.

15

5. An isolated, purified, or recombinant polynucleotide consisting essentially of a contiguous span of 8 to 50 nucleotides of anyone of SEQ ID Nos 1 and 2 or the complement thereof, wherein said span includes a *TBC-1*-related biallelic marker in said sequence.

6. A polynucleotide according to claim 5, wherein said *TBC-1*-related biallelic marker is selected from the group consisting of A1 to A19.

20

7. A polynucleotide according to any one of claims 5 or 6, wherein said contiguous span is 18 to 35 nucleotides in length and said biallelic marker is within 4 nucleotides of the center of said polynucleotide.

25

8. A polynucleotide according to claim 7, wherein said polynucleotide consists of said contiguous span and said contiguous span is 25 nucleotides in length and said biallelic marker is at the center of said polynucleotide.

9. A polynucleotide according to claim 8, wherein said polynucleotide consists essentially of a sequence selected from the following sequences: P1 to P7, P9 to P13, P15 to P19, and the complementary sequences thereto.

30

10. A polynucleotide according to any one of claims 1 to 6, wherein the 3' end of said contiguous span is present at the 3' end of said polynucleotide.

35

11. A polynucleotide according to any one of claims 5 or 6, wherein the 3' end of said contiguous span is located at the 3' end of said polynucleotide and said biallelic marker is present at the 3' end of said polynucleotide.

5 12. An isolated, purified, or recombinant polynucleotide consisting essentially of a contiguous span of 8 to 50 nucleotides of anyone of SEQ ID Nos 1 and 2 or the complement thereof, wherein the 3' end of said contiguous span is located at the 3' end of said polynucleotide, and wherein the 3' end of said polynucleotide is located within 20 nucleotides upstream of a *TBC-1*-related biallelic marker in said sequence.

10 13. A polynucleotide according to claim 12, wherein the 3' end of said polynucleotide is located 1 nucleotide upstream of said *TBC-1*-related biallelic marker in said sequence.

15 14. A polynucleotide according to claim 13, wherein said polynucleotide consists essentially of a sequence selected from the following sequences: D1 to D19, and E1 to E19.

15. An isolated, purified, or recombinant polynucleotide consisting essentially of a sequence selected from the following sequences: B1 to B15 and C1 to C15.

20 16. An isolated, purified, or recombinant polynucleotide which encodes a polypeptide comprising a contiguous span of at least 6 amino acids of SEQ ID No 5.

17. A polynucleotide for use in a genotyping assay for determining the identity of the nucleotide at a *TBC-1*-related biallelic marker or the complement thereof.

25 18. A polynucleotide according to claim 17, wherein the polynucleotide is used in a hybridization assay.

30 19. A polynucleotide according to claim 17, wherein the polynucleotide is used in a sequencing assay.

20. A polynucleotide according to claim 17, wherein the polynucleotide is used in an enzyme-based mismatch detection assay.

35 21. A polynucleotide according to claim 17, wherein the polynucleotide is used in amplifying a segment of nucleotides comprising said biallelic marker.

22. A polynucleotide according to any one of claims 1 to 21 attached to a solid support.
23. An array of polynucleotides comprising at least one polynucleotide according to claim 22.
24. An array according to claim 23, wherein said array is addressable.
25. A polynucleotide according to any one of claims 1 to 21 further comprising a label.
26. A recombinant vector comprising a polynucleotide according to any one of claims 1 to 4 and 16.
27. A host cell comprising a recombinant vector according to claim 26.
28. A non-human host animal or mammal comprising a recombinant vector according to claim 26.
29. A method of genotyping comprising determining the identity of a nucleotide at a *TBC-I*-related biallelic marker or the complement thereof in a biological sample.
30. A method according to claim 29, wherein said biological sample is derived from a single subject.
31. A method according to claim 30, wherein the identity of the nucleotides at said biallelic marker is determined for both copies of said biallelic marker present in said individual's genome.
32. A method according to claim 29, wherein said biological sample is derived from multiple subjects.
33. A method according to claim 29, further comprising amplifying a portion of said sequence comprising the biallelic marker prior to said determining step.
34. A method according to claim 33, wherein said amplifying is performed by PCR.
35. A method according to claim 29, wherein said determining is performed by a hybridization assay.

36. A method according to claim 29, wherein said determining is performed by a sequencing assay.

37. A method according to claim 29, wherein said determining is performed by a
5 microsequencing assay.

38. A method according to claim 29, wherein said determining is performed by an enzyme-based mismatch detection assay.

10 39. A method according to any one of claims 29 to 38 wherein said *TBC-1*-related biallelic marker is selected from the group consisting of A1 to A19 and the complements thereof.

1/5
Figure 1

	1				50
Mur. tbc1	MPMLPWVVAE	VRRLSGQCSK	KEPRTKQVRL	WVSPSGLRCE	PDLEKSQPWD
TBC-1	mpmlpwvvae	vrrlsrqstr	kepvtkqvrl	cvspsglrce	pepgrsqgwd
dmu50542	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
celf35h12	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
Consensus	-----	-----	-----	-----	-----
	51				100
Mur. tbc1	PLICSSIFEC	KPQRVHKLIH	NSHDPSYFAC	LIKEDAAHRQ	SLCYVFKADD
TBC-1	plyssifec	kpqrvhklh	nshdpsyfac	likedavhrq	sicyvfkadd
dmu50542	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
celf35h12	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
Consensus	-----	-----	-----	-----	-----
	101				150
Mur. tbc1	QTKVPEIISS	IRQAGKIARQ	EELRCPSEFD	DTFAKKFEVL	FCGRVTVAHK
TBC-1	qtkvpeiiss	irqagkiarq	eelhcpsefd	dtfskkfevl	fcgrvtvahk
dmu50542	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
celf35h12	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
Consensus	-----	-----	-----	-----	-----
	151				200
Mur. tbc1	KAPPALIDEC	IEKFNHVSCG	RRTDWEAPTG	Q.....PSA	PGPRPMRKSF
TBC-1	kappalidec	iekfnhvsgt	ggprapaptr	pmprpqwsqe	pvrprmrksf
dmu50542	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
celf35h12	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
Consensus	-----	-----	-----	-----	-----
	201				250
Mur. tbc1	SQPGLRSLAF	RKEFQDASLR	SS.TFSSF.D	NDIENHLIGG	HNVVQPTDME
TBC-1	sqpglrslaf	rkeldqgglr	ssgffssfee	sdienhligg	hnivqptdie
dmu50542	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
celf35h12	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
Consensus	-----	-----	-----	-----	-----
	251				300
Mur. tbc1	ENRTMLFTIG	PSEVYLISPD	TKKIALEKNF	KEISFCSQGI	RHVDHFGFIC
TBC-1	enrtmlftig	qsevylistp	tkkialekknf	keisfcsqgi	rhvdhfgfic
dmu50542	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
celf35h12	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
Consensus	-----	-----	-----	-----	-----
	301				350
Mur. tbc1	RECSGGGSGG	FHFVCYVFQC	TNEALVDEIM	MTLKQAFTVA	AVQQTAKA.P
TBC-1	ressgg..gg	fhfvcyvfqc	tnealvdeim	mtlkqaftva	avqqataka.p
dmu50542	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
celf35h12	MEDFKDFTEV	TQFTNVQYLG	CSQLVNNDND	NEMKALMKVL	DEQKGAQTIN
Consensus	-E-----	--F-----	-----	---K---V-	--Q--A----

2/5

Figure 1 (Continued I)

	351				400
Mur. tbc1	AQLCEGCPLQ	GLHKLCEIE	GMNSSKTKLE	LQKHLTTLTN	QEQTIFEEV
TBC-1	aqlcegcplq	slhklcerie	gmsssktkle	lqkhlttltn	qeqatifeev
dmu50542	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
celf35h12	VTLVVPHNIS	GTVKLIDAQG	KVLSSFSLVN	IRFCIRGESE	TSQNNCF.GI
Consensus	--L-----	---KL-----	---SS-----	-----	--Q---F---
	401				450
Mur. tbc1	QKLRPRNEQR	ENELIISFLR	CLYEEKQKEH	SHTGAPKQTL	QVAAENIGSD
TBC-1	qklrprneqr	eneliisflr	clyeekqkeh	ihigemkqts	qmaeenigse
dmu50542	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
celf35h12	SFTHKISVGE	HNSSDILHQC	HVFRTSKAET	AAKALYSFSY	AFSNKNVSSE
Consensus	-----	-N---I----	-----E-	-----	-----N--S-
	451				500
Mur. tbc1	LPPSASRFRL	DSLKNRAKRS	LTESLESILS	RGNKARGLQD	HSASVDLDSS
TBC-1	lppsatrfrl	dmlknkakrs	lteslesils	rgnkarrlqe	hsisvldss
dmu50542	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
celf35h12	SNRLEFQFES	ILEVKENDGT	VEKPSWKLCP	QHNGVFKVRR	DREKKIVVQL
Consensus	-----F--	-----	-----	--N-----	-----
	501				550
Mur. tbc1	TSSTLSNTSK	ELSMGDKEAF	PVSETSFKLL	GSSDDLSSDS	EGHIAEESAL
TBC-1	lsstlsntsk	epsvcekeaf	pisessfkll	gssedlssds	eshlpeepap
dmu50542	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
celf35h12	RQVARKKTID	GFLLNKKCF	GMLLAAGRNL	RHSDLQLLEM	DRNATGTDSA
Consensus	-----T--	-----K--F	-----L	--S-----	-----
	551				600
Mur. tbc1	LSPQQAFFFF	ANTLSHFPE	CPAPPEPAQS	SPGVSQRKLM	RYHSVSTETP
TBC-1	lspqqaffff	antlshfpie	cqepqparg	spgvsqrklm	ryhsvstetp
dmu50542	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
celf35h12	VFVIEA..NW	DPRVHMFVLE	NTETPRDTRV	FMTVAIDVIV	SEISEPIRFS
Consensus	-----A----	-----F---	-----P-----	---V-----	---S-----
	601				650
Mur. tbc1	HERKDFESKA	NHLGDTDGTP	VKTRRHWSRQ	QIFLRVATPQ	KACDSPSRYE
TBC-1	herkdfeska	nhlgdsggtp	vktrshswrq	qiflrvatpq	kacdssrye
dmu50542	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
celf35h12	MEAMSRVFHE	HERFYKTPQT	VVSEEFVLVL	EVRIIRTEKLL	ETHGNMLLKS
Consensus	-E-----	-----V-	-----R-----	-----	-----
	651				700
Mur. tbc1	DYSELGELPP	RSPLEPVCE	GPFGQYRKKR	GRRHASFESC	GKRPSCSRSC
TBC-1	dyselgelpp	rsplepvced	gpfpqhqrkr	kghlvssesc	akrlffnryc
dmu50542	~~~~~	~~~~~	~MRKPAKRG	KRDAAELREL	WRTAIRQTIM
celf35h12	VPIDFAWQLE	GVYFLPTPSK	SCDQSDPNDR	KLTFISLESD	SDRKRSKQNL
Consensus	-----	-----P----	-----R	K----S-ES-	--R-----

3/5

Figure 1 (Continued II)

	701		750
Mur. tbc1	...LVRMEKE NQKLQASEND LLNKRLKLDY .EEITPCLKE VTTVWEKMLS		
TBC-1	...clgmeke nqklqasend llnkrlklrf mkeitpclke vttvwekmls		
dmu50542	...LNRMETE NAMLQARQNE NELKRIKLDY .EEIVPCDKQ LIERWEQIIE		
celf35h12	GKSPSRMPTQ LLHPTGDDES DCDEPLLSGS GKVSQECKEE HLEMWDQLIE		
Consensus	-----RME-E N--LQA--N- ---KRLKL-- --EI-PC-KE ----WE----		
	751		800
Mur. tbc1	.TPGRSKIKF DMEKVHSAVG QGVPRHHRGE IWKFLAEQFH LK.HPFP SKQ		
TBC-1	.tpgrskikf dmekmh savg qgvprhhrge iwkflaeqfh lk.hqfp skq		
dmu50542	RNSTQIGNKK DPKVLGHAIR TGVPRSKRGD VWTFLAEQHS MNTAPVDTKR		
celf35h12	NWDQQSD... RPQKISELVL DGIPDKLRGR VWQLLSNAI.....D		
Consensus	-----S--K- D--K---AV- -GVPR--RG- -W-FLAEQ-- -----K-		
	801		850
Mur. tbc1	QPK.DVPYKE LLKCLTSQOH AILIDLGRTE PTHPYFSAQL GAGQLSLYNI		
TBC-1	qpk.dvpyke llkqltsqgh ailidlgtrtf pthpyfsaql gagqlslyni		
dmu50542	FPNFNTPYHM LLKHLTEHQH AIFIDLGRTE PNHQFYKDPL GLGQLSLFNL		
celf35h12	QPDLEKYHI FLSQPCPSEQ VIMRDIHRTF PAHDYFKESQ GKGOOSLYKI		
Consensus	QP----PY-- LLK-LT--QH AI-IDLGRTF P-H-YF---L G-GQLSLYNI		
	851		900
Mur. tbc1	LKAYS.LLDQ EVGYCQGLSF VAGILLLHMS EEEAFKMLKF LMFDMGLRKO		
TBC-1	lkaycllldq evgycqglse vagilrlhms eeeackmlkf livdmglwvq		
dmu50542	LKAYS.IIDP ELGYCQGLGF ICGVLLLLHCD EANSFQLLKH LMFRRNMRTK		
celf35h12	SKVYS.LYDE EVSYCQGLSF LAASLLLLHMP EEQAFCTLVK IMFNYGLRDL		
Consensus	LKAYS-LLD- EVGYCQGLSF -AG-LLLHM- EE-AF--LK- LMF--GLR--		
	901		950
Mur. tbc1	YRPDMITLQI QMYQLSRLLH DYHRDLNHL EEHETGPPTY AAPWFLTVEA		
TBC-1	yrpdmilqi qmyqlsrllh dyhrdlnnhl eeheigpsly aapwfltmfa		
dmu50542	YLPDMKKFOL QLYQLSRLLV DHPDLVWL DQNDVSPTLY AAPWILTVES		
celf35h12	EKLGFEDNLHL RFFQLTALLK DYTPDLSHHL EHIGIETHMY ASQWFLTLET		
Consensus	Y-PDM--LQ- Q-YQLSRLL- DY--DL--HL E-----P--Y AAPWFLT-F-		
	951		1000
Mur. tbc1	SQFPLGFVAR VFDMIFLQGS EVIFKVALSL LGSHKPLILQ HENLETIVDF		
TBC-1	sqfplgfvar vfdmiflqgt evifkvavsl lgshkplilq henletivdf		
dmu50542	SQFPLGFVAR VFDLLFLESS DVIFKFAIAL LSVHKQQLLA KDNFEEIMDY		
celf35h12	AKFPLQMVFF ILDLFLSQGM NTIFHISLAL LDDAKTDLLQ LD.FEGTLKY		
Consensus	SQFPLGFVAR VFD--FLQG- -VIFK-A--L L--HK---LQ --N-E-I-D-		
	1001		1050
Mur. tbc1	IKNTLPNLGL VQMEKTISQV FEMDIKQLQ AYEVEYHVQ EELIESSPLS		
TBC-1	ikstlpnlgl vqmektinqv femdiakqlq ayevehhvlq eelidsspls		
dmu50542	LKTVVPKMEH TCMEQIMKLV FSDMIGKQLA EYNVEYNVLQ EEI.....TT		
celf35h12	FRVSLPRKYR T..EASTKCL IHKAVKFRLN HSKLEVYENE YKRIKELERE		
Consensus	-K--LP---- --ME-----V F-MDI-KQL- -Y-VE--V-Q EE-I-----		

4/5

Figure 1 (Continued III)

	1051				1100
Mur. tbc1	DNQRMEKLEK	TNSTLRKQNL	DLL.EQLQVA	NARIQSLEAT	VEKLLTSESK
TBC-1	dnqrmdklek	tnsslrkqnl	dll.eqlqva	ngriqsleat	iekllssesk
dmu50542	TNHHLEMLNR	E...KTQNQ	HLE.QQLQFA	QSSIAQLETT	RSSQQAQITT
celf35h12	NEDPVLMEK	EIGRHQANTL	RLERENDDLA	HELVTSKIEL	RRKLDVAEDQ
Consensus	-N-----LEK	-----QNL	-L--EQLQ-A	---I-SLE-T	--KL---E--
	1101				1150
Mur. tbc1	LKQRALTLEV	ERRPAADGGG	AAEAKRPAQH	SR.ARLHPAG	AHRRRLTAAR.
TBC-1	lkqamltlel	ersallgtve	elrrrsaeps	drepectqpe	ptgd~~~~~~
dmu50542	LQSQVQSLEL	TIQTLGRYVG	QLVEHNP...	DLELPNEVRR	MLQQLDDLDLR
celf35h12	IETSANAIER	LTRQNMDILE	E..NKNLMRE	YEQIKEMYRR	DVLRLEENGs
Consensus	L-----LE-	-----	-----	-----	---RL-----
	1151				1200
Mur. tbc1	RDCAPLTLsk	P~~~~~	~~~~~	~~~~~	~~~~~
TBC-1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
dmu50542	QRRKPIFTER	KIGKSVSVNS	HLGFPLKVLE	ELTERDELGS	PQKQKKEKTP
celf35h12	RAEKLLAEYK	KLFSERSKRA	ENEREHFEVQ	KKAIARISD	CDKCWPAVCE
Consensus	R-----	-----	-----	-----	-----
	1201				1250
Mur. tbc1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
TBC-1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
dmu50542	FFEQLRQQQQ	QHRLNGGGQS	SNVGESGSPT	PPSRPNRLLD	NASARTVMQV
celf35h12	.WEKNRSPVH	SASTPTGPD	LTKLEEREDH	IKNLEIDLAQ	TKLSLVEAEC
Consensus	-----	-----	-----	-----	-----
	1251				1300
Mur. tbc1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
TBC-1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
dmu50542	KLDELKLPEH	VDKFVANIks	PLEVDsgvgT	PLSPPSTASN	SSGGSIFSRM
celf35h12	RNQDLTHQLM	AQSESDGKKW	FKKTITQLKE	VGSSLKHHER	SNSSVTPHFS
Consensus	-----	-----	-----	-----	-----
	1301				1350
Mur. tbc1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
TBC-1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
dmu50542	GYRTTPPALS	PLAQRQSYGV	AITTAPCPQH	MEEVAPATTM	AVMPQEDVEE
celf35h12	STFQLQMDHT	ETTTsNNIGY	NSSSESFAVR	FMQTPSAVLK	ITNGEMTEDN
Consensus	-----	-----	-----	-----	-----
	1351				1400
Mur. tbc1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
TBC-1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
dmu50542	PQPMHPLSMV	GGDVNVRFKG	TTQLKSIRPV	HHMRAIPLGG	VQHPSSTEPa
celf35h12	NNMLHGINGV	DLLDLQSTDN	DDQYSNssSL	ESRNSLTNHQ	GKAEDSTMVT
Consensus	-----	-----	-----	-----	-----

5/5

Figure 1 (Continued IV)

	1401					1450
Mur. tbc1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
TBC-1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
dmu50542	VRVAPVPVEL	APPAATATTG	RS~~~~~	~~~~~	~~~~~	~~~~~
celf35h12	VNLDQLPARR	TLLKCLVLVV	FGKSYTKVIF	QMLRFSGFLM	RRGVYPGTLY	
Consensus	-----	-----	-----	-----	-----	-----
	1451	1465				
Mur. tbc1	~~~~~	~~~~~				
TBC-1	~~~~~	~~~~~				
dmu50542	~~~~~	~~~~~				
celf35h12	FRRCSKILLK	KYDRI				
Consensus	-----	-----				

<110> Genset SA

<120> Nucleic acids encoding human TBC-1 protein and polymorphic markers thereof.

<130> D.18363

<150> US 60/095,653

<151> 1998-08-07

<160> 7

<170> Patent.pm

<210> 1

<211> 17590

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> 1..2000

<223> 5' regulatory region

<220>

<221> exon

<222> 2001..2077

<223> exon 1

<220>

<221> exon

<222> 12292..12373

<223> exon 1b

<220>

<221> exon

<222> 12740..13249

<223> exon 2

<220>

<221> allele

<222> 9494

<223> 99-430-352 : polymorphic base A or G

<220>

<221> primer_bind

<222> 9391..9408

<223> 99-430.rp

<220>

<221> primer_bind

<222> 9828..9845

<223> 99-430.pu complement

<220>

<221> primer_bind

<222> 9475..9493

<223> 99-430-352.mis

<220>

<221> primer_bind

<222> 9495..9513

<223> 99-430-352.mis complement

<220>

<221> primer_bind

<222> 9482..9506

<223> 99-430-352.probe

<220>

<221> misc_feature

<222> 3953,4056,4167,4739,6217,6245,6860,9998..9999,10006,10012,10104

10477,10822,10825,11095,11256,11273,11857..11858,11895..11896

14057,15912..15913,16217..16218,16329..16330,17504

<223> n=a, g, c or t

<400> 1

aggacagtat ctagcacaat accccaaatc gactaaactcc tccgtaaaga atagctacca	60
ctattgtgag agttttaagt caagctgtga ataaaactct tgggtccact taaaaatacc	120
tcccctggat gtaagcatcc agggaaatca gggaatgcc taagacagcc ctaatctaaa	180
agcctacaag aagctcagtg ggcttcaagg aagacactgc tcttggtacg atgaggaaac	240
ctggccctct atttgccctc tgggccacag taatattgat aatagctgct gcttttagtt	300
gaggaccatg tacgtctgtg tcaactgcact ggccacttta cttacacttt cctgctttgt	360
cctcacaag atcctgtaag gtgtgtattg gtcccattta gcaggtaaga caatgaagac	420
cagaggtcca gcaccttgcc taaaccacac ctgctgggat ttggattcaa gtccaaccgt	480

acagctcaaa cgctcagcca cttccctaata gtccaccccc agctacatta agtaaaaaaa 540
tccagaaaga tgccacctgg gggctctggaa ctgcctcctc cgagcaccgc gctctcccct 600
ccctgcggac tcttctctgg agaggatgtg atgcttctta cttttctcag atccctctcc 660
ccacctgcgc agtgacgttg cgcctctgtg cctgggtggga tagggatctg ggagcttcgc 720
ctgttttttg cactctgcca tcccctagtc ttagggagcg agctctgtcc cgcttttcac 780
atctccgcgt ctttccttgc actctacatc accgctggga atgtccccag acctgatcgg 840
ggcatgcaca ctgggggtgtg cgtgtgcgtg tgggtgtgtg tcctgcgcgt gtgccgggct 900
cgcgggggcag gaaaaagcgc ctaatccagg ctctgcgtca ctcccgcaat tggtagaaa 960
tggagtttcc tgggtgttaa tcccgggagg gcacttcgcc ttcgttggtt cccagagtcc 1020
ctgattttcc tgcctcgcgt gccagcgccc catagggcat ccgtgcctca gttcacctct 1080
tgccatcctc caaggacggg gagaaggggt aaggcgggg agagcaagggt ggcttggtcg 1140
cccccgccc ccgccccca tgttgtgtgc agtttccacc acgtctgttt cggagggaga 1200
agaggagggt gcagatgagg cgaggcgctc tccggagcgc ggagagcggg caggcagtgc 1260
cacctgctga gagccactca ggccgagcaa gcgcggggca gtgccacctg ctataaatag 1320
gccgccaagg acagggtgtg cgactgtaca tcccgccacg agggcctgca tcacgcgcgg 1380
ggccccgcgc ccccggtctc ccagggaac gctgtgcca gatcctgcgc aggggtctgg 1440
atggggcggc ggcccgagta cttccccct attccccca cagacactgg ctgaggatgg 1500
cccgcgggct tggggcggg ggggtggcaag gaggggaggg aggcgcggc ggaccgcag 1560
tgcagcagct gttgctcgc tgtgactgc ccgtccgggc cgtgctgccc aggcacagtc 1620
acacggcgca gtggggagga ggaggacacc gagtccccct cccagctccc cggggaccga 1680
gtggggagat cccggctcct gtcttcccc cgcctccagc gcgctcgccc aggtgggag 1740
gaggaaacca gagccgcgc cagacacctc ctcttctctc tctcttctt cctcctctc 1800
ctcctctctc tctcctctt cggctgctgc tctgtgtgcc gccaccgtcc gccgggtgct 1860
gttgcctgcc ccgcccggg acctgctgtg tctcagctg ggtggagaag aggcgggcgc 1920
cgagccgagg ggagccccct ccccgctccc ccgcgggcggg aagagcgcag ccagccgggt 1980
gcgatggact ccccgccgc ccaggccgtc cccaggatgc cccaagcac ctgcgcgtcc 2040
cgccccggcc ccgggtctct agcgcgcgc ggcacaggta aggcgcttcc tggggcttcg 2100
tctggccac cctgctggct cctctcgggg cgtcgcggcc gccccctccc gcagcacgcc 2160
cctgccccgc ctggccgcgc aggggaaggc atctggccgc ccacggacgc gaggccaggg 2220
tctctcgggg gaggaagtct attgccatct cgttgcccc cttaccccc ccccccgcc 2280
gcccttgga gaaagcgaaa ccttaatgtt gctagcgacc cgagagctcc gccggcttct 2340
cccccaacc ccgccagctc actggtccgc gcattctctc cctccccct cccgccaatt 2400
atcctagcgt gtttgcaagg cgaccagatt ggaaagagt tggtcagagt gacccaagc 2460
cacgctttaa aagttcaggg tactttgcag tagtaacttt ggcagctcca ccagtgcgcg 2520
caacatttct ttctatgggt acatcctgta ccagtcattt tgaaacctg cttcattgtt 2580
tctagccgct tctgatggc tctgtgatta tgagaccccc ctcaaacttc accaggcatt 2640
aaggttttgt ttttgctttt ttttcagaga ggtatcattt cgtttgaaat ccacctagat 2700
gtggcttttc ctgttttgat tttacttaac atagcttatt ctctggaagt tgctttaaaa 2760
agaaattgaa agtgatgggt gttccttcca ccaaacagtt taattttcag ggtgcctcat 2820
attaatggat atgttttccc ttcataagatt tctcattgtt tcccttatga tgggatgatt 2880
tcatttatta ataaaatcag actttgaaag agcattttaa aatgacctgg tttaaatagg 2940
tcacacccaa gaaactcagc tatctgtaca agttcaaact tctaaacttt ttcaatgagc 3000

taggggtggt ggcacccacc tgtagtccca gctacttggg aggctgaggc aggaggatca 3060
cttgagccca ggagttcgag gccatagtga gctatgactg tgccacctca ctggagcctg 3120
ggtgacaaaag tgagatccca tctcttaaaa aaaaagagtt taggggacat tttctgaagt 3180
gaacacaagt agagcattct aacactattg agtgcaagga gacctggaag ggactaagtg 3240
gttcaaagca ggaaataaaa tcatcagggtg ataattaaaa taatttcttt cctgtggatt 3300
tgtccagcca tttgcaaacc aggagaatag gaaaaaaaaat cactagtgtg gttataaatt 3360
attacattac gttttcaaag gaaaattttg caaatgcgtc tcttgtcat agtctattgt 3420
tatctacccc actgagagtg ctggggcttc ccttttcac cacgacagca tttctggtg 3480
ggtggcagtc atgcagtgtt gacctggtgt ccataaggc acagtttgtc aaaacactag 3540
tgggtattag gaggaacgt gcaactctga agcaacagag cttgcccctt ctctctcatt 3600
atccagctgg tgataatccc tgtccccac ttccctagaa gacagctttg accaggaagg 3660
ctgcaatgac aatgagatgt acccctatgc agagccagat gtgggcgggt ggcttttttg 3720
tggtccagat cttctaggat cttctaggat gtaaccctgg caagcagtgg ggagcctgaa 3780
tcaagcagca tggtgtttac ctcttctgtg ttacagcag catcttcagt tgtcttgggtg 3840
cctggagcag gcaccacagc tgctgtctct gttggccacc agctttctag agtagatggt 3900
agggaggaga gcaaggggct caagaggatt ctgtctttga acatgctttt aantttgatc 3960
tgacagaatg gcagctccct gaagtccttc ctactctctc cacagcattt ctctgtaggt 4020
ccccagtttt tgctcttttc agattcccag aggacntgaa aatgtatcac ggcccatttg 4080
gggacttcct gtatatgtgt ggggtgcctca ggatcatttg ttttgccctt ttccagtcta 4140
ccgtgtgtgc cttctcaagt ttaatgnacc acgttagttt caatatttta tatatttctc 4200
agcagttttc atctcttgggt cattaaactt gagaagtaaa atctgtctcat taaaatgact 4260
gagtcctatgg ccaggcatgg tggtcatgc ctgtaatccc agcacttttg gagtccaagg 4320
cgggtggatc acttgaggtc aggagtctga gaccagcctg gccagcatgg caaaaccctg 4380
tctctacaaa aatatagatc tacaaaaact agccaggcat ggtggcatgt gcctgtagtc 4440
ccagctattt gggaggctga gacaggagaa tcgcttgaag ccaggaggcg gaggttgcag 4500
tgaaacatga tcgtgccact gagtccattc agcagcagag tagtgttggg gtttgtatcc 4560
ctgtagtgat gacgaaggat ttaggttttc agtcagaact gttaccttac aatttccttc 4620
actgactttt ctccctttcc aacaccacat tccaataaaa aatatcttta gaccagattc 4680
ttcacgaaag acatgaagggt tttcatgctt caagggtttt gacttttttt ttttttttna 4740
aaggagtctt gctgtgtcac ccaggctgga gtgcagtggc gtgatctcag ctactgcaa 4800
cctccgcctc ctgggttcaa gtgattctcc tgctcagcc tcccaagtag ctgggactac 4860
aggcgtgtc taccacggcc ggctaaattt tgtgttttta gtagaggcga ggtttcacca 4920
tcttggccag gctggtcttg aactcccgac cttgtgatcc acccgcttg gcctcccaaa 4980
gtgctgggat tacaggtgtg agccacggcg cccgaccagt ttttgacatt tctaagccaa 5040
aagttccatt tgatgaggtc ttagatgcag gggcaatgtg tcccttttca gatttcagat 5100
gtttagaaaa agatgtgtca ttttggggc aactgaaaaa ctcttgatat gtaggttttt 5160
atgaagctgt gcagaatgta ggaaatacat tttagaacca acaaagaggc atttaatttt 5220
gagtgtgcct gtctccttg agatgagcaa cagctatttt tctcttcaa agacaatgcg 5280
tgtatttatc agcacatttt atataatcag caaatctaaa cctctgaatt aggttaagccc 5340
tataggtttg ttgccagaat tagtgaattt atacatgcaa agtgcttaga acagtgcctg 5400
gtacacagtg agcactcaat attattttatt gctattatta tgtttattta ttttatactt 5460
ttagagtata attttgatgt taggtttgga ttgctgaggc caagcaaat ttagatagac 5520

caaccagct aatccactag aaagatattt gagggttatt cccatctaaa gatctatggg 5580
atctttggat atcatctgtg agaaacaaca gaagtttgta gataagacag atatagattc 5640
aaatgccacc ttcacaattt attagtgagg gaaccctttg gtaaaatgag catgacaaaa 5700
cctacttctc agctttgtga acgtacagaa gatcatgaat gtaaaatgtc catgaagtgg 5760
taggtgggtca acagatattg ctctagcaaa gtgggttaaga gcaagcaaac tctggagcca 5820
aagggcctgg gttcaaatcc cgattctgcc acttcttata gtatggcctt gggcaggtga 5880
cctcactttt ctgtgccact attcaatgat aatattcctt tattgtccaa cgttttghta 5940
gttaaatcag ttaataaaca cactatgata atgtgttggg aactattctt tttactttta 6000
gcagaataac ttaaagggaac actgcaggta ggaggggttat acataatctc tgagggccag 6060
ctaggacact cgcccatttc ccaccctttt tcctgtgcaa tgaagagtat aagaagtga 6120
agtgccattc taaaggacta gccttgagtt ggctctaatt tatatgactc gtgcctgtaa 6180
tcccagcact ttgggaggcc gaggctggca gatcacnttg aggtcaggag ttggagacca 6240
gcctnggcaa acatgacgaa atctcatctc taccaaaaat acaaagatca gccgggcatg 6300
gtagcacaca cctgtaatcc cagcttctgg ggaggtgag gcagaagaat tgtttgaacc 6360
cagggggcgg aggatgcagt gagctgagat cacgccactg cactacagcc tgggtgacag 6420
agtgagattc tgtctcaa atataataa atttatatga gaaagaagtc attcaaaagc 6480
atcattactt tacatgtcaa attagaaagg cacaccccag tactaaagca tccttgatga 6540
tgaaaacatt tagaaccact ggtttcagga gctccatgca atggtgaaac agcctctact 6600
ccaagggttg ttgctccctg tgggattcct ggggtgaggaa cactactgctc ccgttggggg 6660
ggaatcctgt ggaggaagtg gatgaagagt gtagccaagt cagtgagcct actgcatggg 6720
attagagtac ttcaggttgc agtataattc tgttcagggt catgctcact ccactctggc 6780
taaagaacag agaaattaaa ccattgattc acagagcaat atgagtagct gcctggggac 6840
ttcctgtcca ttctggggtt cccaacagcc aatcaatatt ggccgggttc taacttgacc 6900
tagagctaga ggccactagc acccctccat tcatccttct ttctctccct acccactccc 6960
accagcattc tgaggaggtg agggctgaag ctgcagaggc tgttgactg tcagttactg 7020
taaagtcccc atcctgctct ctgtagtttg ctgtgaagga gtggaggggg ctgggaacaa 7080
aggggggttc caataagcag cttactaata cctacccttg cttctctcac ttctgatca 7140
ataattccca tcctttcttc atgtacctcc cccacatttt tgttctttaa gggaagaagg 7200
gagcagcttt ccacccccc acaatgttgg ggacatttgg tattatacat tatgaaaata 7260
agatttacca gatttttagag atggaagaaa acttggggat gatcttgctt cattctctta 7320
taagaacaaa caatatttgg agaagctgag taacttgctt tttcaacttc acacttgaaa 7380
gtgttttcat gaggaagttg gggttctctg cagcacttgg atgggagtca gggacttgga 7440
ttgtcccaat tctgtcacta aatttggaca agccacttaa cttttccaga atctagttgc 7500
ctcatttcaa atattagga aatttctaaa tggcttaaaa ggagcttggt agctttaaaa 7560
ttttatgatt ctaagtgtat gctgccagag atatgtagca tagcaggaca cattaacaag 7620
attattgaaa ctgttcta ataaaggacatc tttgtgtctt gggtagctac tatgtttaaa 7680
gactgtgcta ggtgggagtt gtgcagaata cacaggttgg ctgtagaggg atagggcgtg 7740
tacacagaca actctactaa caagaacgtt actagaagct tattggaatc acagtatttc 7800
ttgctgaggg tatgaaacat aagagttctc cttggaatat gaggttctat ttggggctta 7860
aagaatggtc aaagggttag tgcaataaac atggattgag atggctttaa aaaataatca 7920
aatggtttgt tagtattaaa ctggtgcaga aataattgca gtttttgcca ttccctttta 7980
tggcaaaaat tgcagttact tttaaaccaa atccctaata ttatttgc atgtttatctc 8040

tgttatggaa gtttttattg acaagtaatg tagatattca cctgatctaa gttaccctga 8100
atcttatatt agcagaatct gaattgctta taaataatta tggctatggt ggatgtagaa 8160
cttattatct gatagtttat gaacagtgtt aagggtctaat ctacttttta cagagaagct 8220
aagaacatgc tacagctggt tgaaaaacaa aaacttcagg cattgaaatg ttttgtcaat 8280
gaaatggcag gactcatttg atgactgatt attatcaact gatttaaag actgaatttt 8340
tggtactgtg tacatctata ctctaagaag gaaattgaaa gtaattctgc tatgcttggt 8400
gccactatat taataactgc atcatctaaa ataattgata gagctcagat ttatcctttg 8460
taataattct agtacttctt taaacatggt tggggattag cagctgtcaa cagttagaac 8520
atgaaacaga ttctgttaca ggagtagaag tcgatccaga catttaatgt cttttcacc 8580
tgtgagagag agaataaaga gaaagagaga tcattattta tgggattatg tgaacttcaa 8640
gtccgttttc attattagga gaagctgtgc tttaaaggac agtcaggagac tttactttca 8700
tgaaatgcct gagctgtaaa taaagtattg ctttattttt tatttcttga acatttgaaa 8760
taaaaaatta gctatgagtt atgttcaaat tatattataa aaatttgctc ttagcattgt 8820
gcatatatat tatacagaaa aacacagagt aaaaagaata gacttcagtt cctgttcaga 8880
aaagggttaa aatttgaata ctgattttgg aaaccccaaa ccttaagaat tcaagaagct 8940
tacggctctc ttgagggaca cctattcaaa ctcttaataa tgggtgattgg gtagaaagtg 9000
cagaaaagcc tgctgataca tgccctaaaa caccttgga aagagagggtg gtagttgctt 9060
gaggtaggac ttaagtacta gttggaaata gaagacaagg atggagactg ttggtagatg 9120
actctccatg ggtccttcct gtttctacac accttgtaag cagggcattg agtgctgtg 9180
ttccaaacta ctttttccat catgtttcta cagcaaacag tcatggaaga tagaaataga 9240
gtcttctctt ggagcaaagg gcagacacgc ttgcttctctg tacttcccac tataagatat 9300
tccggctccc taaactcagc tgcccttcct gtaaccacc atgatacaga tgtcacctga 9360
cctgtgggaa ttgggggtca gggaaaccaag agaaatgctg actgtctggc tactgtgact 9420
gccctgagta ataaattgtc ctctgtctcc aaccaggag tctcatgttt tctaccagca 9480
ggataactgt ggcrggctaa cgtgttagtt tgcaagtaag gtaaaatctc agaccctttg 9540
cagtttgtgg cagggtattt attctgagga gagaggaacc gtatgcacca tggctcagag 9600
gcatgagaaa cggggaacca taactagttc tctatcttca gagcctttta aagggtgcacc 9660
aaggagggca ttttagggga gaatataaag ttggagatat agacacagcc agattcctga 9720
gagaccttat atgccaggta gaagacttca gattgtatgg gggaattatt agagaatttt 9780
tagcaggggt gtgatatgat aaattttgtg ttgattaagt tactccagga aatatgcat 9840
gggtggattg aaggatgggg caccttttct ctaggacgaa aaagaaagag tagttggtga 9900
agtcagttag aggtagtaat aggatgaaga agggatctga atgaccctt ggccattcag 9960
tgagtagtga tgctattcac ctgatacag cacatagnng ggaaangaaa tncgtgggaag 10020
gaggagatg agaccgagtt agctttaaaa taactaaatt caggcctagg agcctatagg 10080
ctatccagat agaaatattt aatngcctat atggatctgg aactcaggaa ggaggcttcc 10140
gtgggagcag aacacttggg caccattagg gtgtatgtgg tagatgcatt cttgtgcagc 10200
agtcaagggg atgggattta gactcaagtg caaattgccc cccatctcct gtgataagtg 10260
actgaagctc tccgggcttc agtttcctag ttcacatag tgggctctag cggataaatg 10320
ttacaaaggt taaatgagac aacataggca aagtgcgtgg tactcaatag aagtcagctg 10380
ctgtcatcag cagcaggatc accagaatgt ggtgcttgac accaaaagat taggtgagat 10440
tgcccaaac agcaggtgaa atgaggggag aggatgnaag tcaaacacag gaagaaaagc 10500
ctttgaagta tgtggaaga aacaaccaga aaggtaagat aagaaccaga agagattcaa 10560

gaaggaaggt gtggccgggc gcggtggctc aagcctgtaa tcccagcact ttgggaggcc 10620
gaggcgggcg gaacacgagg tcaggagatc gagaccatcc tggctaacac ggtgaaaccc 10680
cgtctgtact aaagatacaa aagaattagc cgggcgcggg ggcaggcgcc tgtagtccca 10740
gctactcggg aggctgacgc gggagaatgg cgcgaaccgc ggaggcggag cttgcagtga 10800
gccgagatcg cgccactgca cntcnagcct gggcgacaga gcgaggagcc gtctcaaaaa 10860
aaaaagaaaa aaaaaaaaaa gtaaggaagg tgtggccaag attgagaaat tcgtcagagc 10920
aaacaaggca gtcaggggct aaatagcctc ctttaaattt tacaaccttg aggacctcg 10980
caactttaac agaatttcag tggatcccta gggcaaacca ggctttacaa accaggaatg 11040
gatggtcaat aggaagtgga gacagtaagt gtagacctta ccttggaggg aaggnaagag 11100
aaagagccat ggccaaggga agtttgaaat caaaggaaat atcttttttt ttttttttcg 11160
attggagaga cctcagttat tcttttaaaa tacttattga gcccctcagt tattctttta 11220
aaatacgtat tgagtcctta ctttgagtca ggcacnatgg cagacacgag ggngatagca 11280
gtgaatcaga cagatgcaac gcctgccttc atggagtttc accttagcat ctgtccatat 11340
gctaggggag tggggcaggg gcaggagct ggatacagga gagactgaag atccagggag 11400
caagtgagta aagaataggg cttgagatcc cacagacaac tcagctttga aaaaaagggt 11460
tttgtcatcc aataggacaa gaaggcggtta ggatacatca aacgtggttg ttgaaaacag 11520
aaaagggtcg ggcactgtgg ctcatgccta taatcccagc actttgggag gccaagggtg 11580
gcagatcact tgaggccagg agttcgagac cagcctggcc aacatggtga aaccccatct 11640
ctactaaaaa taaaaaatt agccagggtg ggtggtgcat gcctgtaatc ccagctactt 11700
ggaaggctga ggcaggagaa ttgcttgaac ccaggggggtg gaggttgag tgagccacga 11760
tcgtgccact gcactccagc ccgggcaaca gagcgagact ctgtctcaaa aaaaaaaaaa 11820
ggaagaaaga acatagacag ggaaatgtag ttaaggnnag tttgggtttg ggtttggtag 11880
aagcgttttc tgttnnttgt ttgtttgttt tcagaaagag tctcactctg ttgtccagac 11940
tggagtgcag tggcacaatc ttggcttgct gcagcctctg cctcctggat tcaagcaatt 12000
ctcctgcctc agcctcctga gtagctggga ttacagacac ctaccaccac accaggctaa 12060
tttttgtatt ttagtagag acgggggttc accatgttg ccaggctggt ctcaaactcc 12120
tgacctcagg tgatccacct atcttggcct ctcaaagtgc tgggattaca ggtgtgagcc 12180
actgcacctg gcctaacatt gatatctgtt gatgagaaga agccagggtg tggagtgata 12240
gcttatagca catgaactga ataaaacagt gttaagaca atgtttgcaa cataataggc 12300
actgaagaca tgttaatgga aggtggattt gtgattcaga acctctagac tacctgggcg 12360
agtcttttaa aatgtaagta atatcttaag tgatattact tgtcccagat cagttgttta 12420
aaactgaggt ttaatgctgt cagagtagca ctgtatcgtc ttctatcatg ggggcctttg 12480
ttggcttttag gaggtttgtg tttcatagta gtttccagc gggctctttg ttacctgtaa 12540
tgagtgtgac agttatgcca taaccagggt ttatatggaa tacaattttg agaaagtctt 12600
ttctaggcag agaagcttat ttgaacctct tattatattt gggtttcagg cttttgagtt 12660
cttctgaaat aatagccctt tgaaggtagc tattgctatg acttcattaa attctaagc 12720
ctctggtttt cttccccagg tttctgcata tgaagtgtgt aaaatagatt gcttgatcca 12780
aaacagaaaa acagtgataa ctgttttgct gagttcccag acccttccca agatggaacc 12840
aataacattc acagcaagga aacatctgct ttctaacgag gtctcggtgg attttggcct 12900
gcagctgggtg ggctccctgc ctgtgcattc cctgaccacc atgcccattg tgcctgggt 12960
tgtggctgag gtgcgaagac tcagcaggca gtccaccaga aaggaaacctg taaccaagca 13020
agtccggctt tgcgtttcac cctctggact gagatgtgaa cctgagccag ggagaagtca 13080

acagtgggat cccctgatct attccagcat ctttgagtgc aagcctcagc gtgttcacaa 13140
actgattcac aacagtcattg acccaagtta ctttgcttgt ctgattaagg aagacgctgt 13200
ccaccggcag agtatctgct atgtgttcaa agccgatgat caaacaaaag taagtgagat 13260
ggagatccaa aagactaagg tgtggctggc tgggtttttat tgtatggggg tcaggatatt 13320
tattttaagt atactgaaat gaataaggaa ttaatgctgc agttataaat tgattactta 13380
gctgaatttt tgttttatgg tgatagttta tagtttttaa gcacatttga aaacagatac 13440
gagaaattat cagtttttga gttcaaaaat tcaagagaaa tcagtctaaa actactaatt 13500
aagagcagaa gtgttaagat gtacattatt tcagatgaat gttctaaagc catgcctctc 13560
aaactgaaat gagcttgtga gtcacctggg gatcttggtta aaatgtgaat cttgattcag 13620
taggtctggg gtggaccca agactgcatt tgtaacaagc tgccaagaaa tgctgatgct 13680
gcccttttgc aggttgcaact ttgagtggca aagttctaaa tctccacatt tgtaatccta 13740
ttaagaaaaa tatagtcatt cgtaaactgt gtaaaaatgc tactggccag tttcccaagg 13800
cataatgttc acttaggcaa aggtcattga taagaacgct ggatatgcat ctaagttttg 13860
atgcgatcag gggttctttg tgtttttttc tttcgcaaac ctcaggtcag atctgattag 13920
cttgttatta tcacatgata tggctgaaaa aaaatgtgag acatggtaaa agttctgctc 13980
tttctcgtt catttgtgct tgctttgtta ttagcattcg ttgtagctct gggcaggact 14040
catttgaaga tgcttgnccc attttatgag gattagctta gataaaattg aaaatataat 14100
gcaaatagca actttctcag ttgggctcag ggctccacag ctaaccccat ggactgtgga 14160
gtcttgccgt tgttttgggt gccaaagcaag ccaagtcaca tgtgattcaa gctgtctgcc 14220
acatgtacag ggcgaggatg cgagtgtcaa tccacctgtt aactgtcagt gaagccttga 14280
aagcttctca tattttcaag gttaaaatct ggatagaaat gctaaagttt tctctctgca 14340
ctccattagg ttattttatg tactctctag ggtgtaagga ccttatttag aaattaatat 14400
tcttggtatc aagtagatgc ccttttgctt gttcatttgt tggttcttct agtcattcag 14460
aattgctgtt gcaggtagt ttggagatga tattagcaga ggctttagg aaggcaggag 14520
catcagtggg gaataggacc aggtgatcta tgtataggac ataatggaag gactgagaag 14580
ggagcctaac acacacccaa agggtagaga aggctttgtg aaataaaggc taatatggag 14640
ctcaaaaacca ccatttcaact cacagaatca aactctcata ttataaatca tttcatgtta 14700
ttgtccacac atctcaagtg ggcacggcag catcaggctt ggagattcag agggactaac 14760
ttcctgtact ctaatcctac ttctgcaccc ataaactggg tggcctcagg caattgagtc 14820
tgttttctta tctgttaaat ggggataatt acagtattta tccaatagag ttgctggaaa 14880
gactaaatga ggtagcactc gacctgaaac ttagtaagca tttatagcca taaaaacatt 14940
ttcattcaag aaaattttac tagaggcaga ttatatgcta atttcatttc acgtcttagg 15000
taaaaagaaa catgatacct agatgagtgc cttcagcttt caaagatgag attctggtca 15060
tatttgagga acatttttaa aactacacgt ataacttaat ggctcctatt atttgacaa 15120
attccagaat gaaaatgaga ggactgaaca gcctgtacct cagtccagct ctatatagta 15180
tttgagactga atttccttgg ggagagtttg tgcgtggaat cggtgttcag cattttacac 15240
atttgactct ttcccaaaat cttttacggc catctgagaa taggcttctg gccagtcatt 15300
cggatgcctg acaagagaaa gagatttata accaaattct gtaattggga cttccagtct 15360
ttcccaagt agagaattgg acttactcta tatgctaaaa acccatgggt gaaatatgaa 15420
ttagttctta agtgattttt ggcttgcata ccatttttgc aaacacaaat tgtcattact 15480
ctgctcattt aataaaagaa taattttag tataggtata tacctcaatc agtgattttg 15540
ttgttggaag cagaacagta aatcacactg gccatgatgc taacagcgtg atagattttc 15600

tgttcttggg acaccaatgt cactgtatct catagcgaag gattatctgc tgtaggagca 15660
ttctcttgac tacttataac atttgctggg tgaataaatt ctccagggtta aggcctcttc 15720
taaacagatg aggtcagcac taactgcatt tgccagagaa gacatatgca tttactgcca 15780
gcatcataaa cacaaaacta cagtttgcca ggaaaccctt tgaccagcat ctaattaatt 15840
cactgagtaa tgtcttggga gaagaggcat gtaaaggaac aattttataa gcatgccatg 15900
agattgtttt cnnattgtat gttccataga atatgaggaa acttcaaac attttgtgga 15960
aaaattgaat taaaaagtaa aaaacacata tatacataag ctttatttct caagataaac 16020
tttatcaagt tcaagacact tttgtaagca atgttaacag ccattgagtc ggtctctaaa 16080
gaactgaggg tcttgggaat ttaaccatgt ttatacagtc ttttatacat tattaactgg 16140
agaaaaattg gcgctcttta aagattttttt aaaattgaga agcaaaagga cgtcagaagg 16200
agccaaatta ggctgnnaa gtggatgctt aatgatttcc catggaaact cttgcaaat 16260
tgctcctgtt tgatgagagg aatgagcagg aacattgtca tgggtggaca ggactctggt 16320
gaagcttttnn caggcgattt tctgctaaag ctttggttaa ctttctcaaa acactctcat 16380
gataaacaga tgttatcatt ctttggcctt ccagaaagtc aacaaacaaa atgccttggg 16440
catcccaaaa aactattgca accatttgcc cttgaccagt ccactttcgc tttgactgga 16500
ccacttctgc tctcagtagc cattgcttaa atttgtcttg atcttttagga ttgcgctggt 16560
aaaactatgt ttcacacct gttacaattc tttgaagaaa tgcttcagga tcttgatccc 16620
accggtttaa aatttccatt agaaactctg ctcttgtctg cagctgatct gagggcaatg 16680
gttttggcac ccatctagta aaacgtttgc tcagtgttaa tttttcatcc aggattgtgt 16740
aagctgaacc agcagagatg tctatgatat tggctagtgt gtctcttca atgagggcat 16800
gaacaagatg aatattttcc tcaacaatt atctggatgg tctgctgctg caggcttcat 16860
cttcaatatt gtctcgtecc ttctttttct tttccccccc gcttgagaca cagtcttgtt 16920
ctgttgccca ggttggagtg cagtggcccg atttcggctc actgcaacct ctgcctcccg 16980
ggctcaagcg attctcctgc ctcagccac caagtagctg ggattacagg tacacatgat 17040
cgtgcctggc taatttttgt attttttagta gagacagggt ttcaccgtgt tggccaggct 17100
ggtctcgaac tcttgacctc aagtaatcca cctgccttgg cctcccaaag tgctgggatt 17160
ataaacatga gccaccacac ctggcctcat ctttcttaa aatgagttat acatttgtaa 17220
gctgctgatt tctttggaca ttgtgcctat aaactttttg taaagcatca gtgatttcac 17280
cattcttcca cccaaacttc accataagtt tgatgtttct tcttgcttg attttagcag 17340
gattcatgtt tctctgatag ggggtctttt caaactgatg tcttatcctt cttagagcct 17400
catccagat cctgttcaga catgctacaa gtttaataca gtttatttgg tgccaaaaaa 17460
tggaatcca tgcatagttt ttaaataata tgcatttttc atgnactttt tgaagacccc 17520
ttgtatactt aaactgctcc acatggaaaa gcttccatga tcaaatgcag taaggcagca 17580
tctcaaacat 17590

<210> 2

<211> 99960

<212> DNA

<213> Homo sapiens

<220>

<221> exon

<222> 4661..4789

<223> exon A

<220>

<221> exon

<222> 6116..6202

<223> exon B

<220>

<221> exon

<222> 9919..10199

<223> exon C

<220>

<221> exon

<222> 14521..14660

<223> exon D

<220>

<221> exon

<222> 50257..50442

<223> exon E

<220>

<221> exon

<222> 56256..56417

<223> exon F

<220>

<221> exon

<222> 63326..63484

<223> exon G

<220>

<221> exon

<222> 76036..76280

<223> exon H

<220>

<221> exon

<222> 78364..78523

<223> exon I

<220>

<221> exon

<222> 85295..85464

<223> exon J

<220>

<221> exon

<222> 93417..93590

<223> exon K

<220>

<221> exon

<222> 97476..97960

<223> exon L

<220>

<221> misc_feature

<222> 97961..99960

<223> 3' regulatory region

<220>

<221> allele

<222> 1443

<223> 99-20508-456 : polymorphic base C or T

<220>

<221> allele

<222> 5247

<223> 99-20469-213 : polymorphic base C or T

<220>

<221> allele

<222> 6223

<223> 5-254-227 : polymorphic base A or G

<220>

<221> allele

<222> 14723

<223> 5-257-353 : polymorphic base C or T

<220>

<221> allelele
<222> 19186
<223> 99-20511-32 : polymorphic base C or T

<220>
<221> allelele
<222> 18997
<223> 99-20511-221 : polymorphic base A or G

<220>
<221> allelele
<222> 19891
<223> 99-20510-115 : deletion of TCT

<220>
<221> allelele
<222> 29617
<223> 99-20504-90 : polymorphic base A or G

<220>
<221> allelele
<222> 42519
<223> 99-20493-238 : polymorphic base A or C

<220>
<221> allelele
<222> 69324
<223> 99-20499-221 : polymorphic base A or G

<220>
<221> allelele
<222> 69181
<223> 99-20499-364 : polymorphic base A or T

<220>
<221> allelele
<222> 69146
<223> 99-20499-399 : polymorphic base A or G

<220>
<221> allelele
<222> 76458

<223> 99-20473-138 : deletion of TAACA

<220>

<221> allele

<222> 78595

<223> 5-249-304 : polymorphic base A or G

<220>

<221> allele

<222> 82159

<223> 99-20485-269 : polymorphic base A or G

<220>

<221> allele

<222> 84522

<223> 99-20481-131 : polymorphic base G or C

<220>

<221> allele

<222> 84810

<223> 99-20481-419 : polymorphic base A or T

<220>

<221> allele

<222> 89967

<223> 99-20480-233 : polymorphic base A or G

<220>

<221> primer_bind

<222> 988..1006

<223> 99-20508.pu

<220>

<221> primer_bind

<222> 1509..1529

<223> 99-20508.rp complement

<220>

<221> primer_bind

<222> 5039..5056

<223> 99-20469.pu

<220>
<221> primer_bind
<222> 5534..5554
<223> 99-20469.rp complement

<220>
<221> primer_bind
<222> 5997..6015
<223> 5-254.pu

<220>
<221> primer_bind
<222> 6332..6350
<223> 5-254.rp complement

<220>
<221> primer_bind
<222> 14371..14390
<223> 5-257.pu

<220>
<221> primer_bind
<222> 14798..14817
<223> 5-257.rp complement

<220>
<221> primer_bind
<222> 18751..18771
<223> 99-20511.rp

<220>
<221> primer_bind
<222> 19198..19217
<223> 99-20511.pu complement

<220>
<221> primer_bind
<222> 19605..19625
<223> 99-20510.rp

<220>
<221> primer_bind

<222> 19986..20005
<223> 99-20510.pu complement

<220>
<221> primer_bind
<222> 29529..29547
<223> 99-20504.pu

<220>
<221> primer_bind
<222> 30041..30061
<223> 99-20504.rp complement

<220>
<221> primer_bind
<222> 42268..42287
<223> 99-20493.rp

<220>
<221> primer_bind
<222> 42732..42752
<223> 99-20493.pu complement

<220>
<221> primer_bind
<222> 69026..69046
<223> 99-20499.rp

<220>
<221> primer_bind
<222> 69525..69543
<223> 99-20499.pu complement

<220>
<221> primer_bind
<222> 76323..76343
<223> 99-20473.pu

<220>
<221> primer_bind
<222> 76771..76790
<223> 99-20473.rp complement

<220>

<221> primer_bind

<222> 78292..78309

<223> 5-249.pu

<220>

<221> primer_bind

<222> 78704..78721

<223> 5-249.rp complement

<220>

<221> primer_bind

<222> 81893..81912

<223> 99-20485.pu

<220>

<221> primer_bind

<222> 82353..82372

<223> 99-20485.rp complement

<220>

<221> primer_bind

<222> 84392..84412

<223> 99-20481.pu

<220>

<221> primer_bind

<222> 84909..84929

<223> 99-20481.rp complement

<220>

<221> primer_bind

<222> 89746..89765

<223> 99-20480.rp

<220>

<221> primer_bind

<222> 90179..90198

<223> 99-20480.pu complement

<220>

<221> primer_bind
<222> 9475..9493
<223> 99-430-352.mis

<220>
<221> primer_bind
<222> 9495..9513
<223> 99-430-352.mis complement

<220>
<221> primer_bind
<222> 1431..1455
<223> 99-20508-456.probe

<220>
<221> primer_bind
<222> 5235..5259
<223> 99-20469-213.probe

<220>
<221> primer_bind
<222> 6211..6235
<223> 5-254-227.probe

<220>
<221> primer_bind
<222> 14711..14735
<223> 5-257-353.probe

<220>
<221> primer_bind
<222> 19174..19198
<223> 99-20511-32.probe

<220>
<221> primer_bind
<222> 18985..19009
<223> 99-20511-221.probe

<220>
<221> primer_bind
<222> 29605..29629

<223> 99-20504-90.probe

<220>

<221> primer_bind

<222> 42507..42531

<223> 99-20493-238.probe

<220>

<221> primer_bind

<222> 69312..69336

<223> 99-20499-221.probe

<220>

<221> primer_bind

<222> 69169..69193

<223> 99-20499-364.probe

<220>

<221> primer_bind

<222> 69134..69158

<223> 99-20499-399.probe

<220>

<221> primer_bind

<222> 78583..78607

<223> 5-249-304.probe

<220>

<221> primer_bind

<222> 82147..82171

<223> 99-20485-269.probe

<220>

<221> primer_bind

<222> 84510..84534

<223> 99-20481-131.probe

<220>

<221> primer_bind

<222> 84798..84822

<223> 99-20481-419.probe

<220>

<221> primer_bind

<222> 89955..89979

<223> 99-20480-233.probe

<220>

<221> misc_feature

<222> 3698,12593,13035,21712,27644,27655,31143,43084,43129,64585,66950
67301..67302,67926,75425,98821..98822

<223> n=a, g, c or t

<400> 2

ctcaagcttg aataacttgaa tccaaacttt catgcttaga gtttacccca tctgttgaag	60
gatgtgcaat ataatgactg caatagaatt cactgtggag cctccaaatt agaaattatt	120
gtctgtgagg gccaggcacg gtggctcacg cctgtaatcc tagcactttg ggaggctgag	180
atgggaggat tgtttgaggc caggagtttg agaccagctt ggtcaatata gcgagacccc	240
catctctgtt tttttttttt aaagaaatta ttgtctaaga accagtgtca tcttccaagg	300
agaaacttct agatacttgt tttaagataa ataagaaaca agtcatttct aaatgtgaat	360
tattttttta atgcaatttt ttaaacattt tatttttaatt atggcaatag acgtggaaaa	420
gactcttttt tgatagtagg ggagagcaga agaaacattg aattaagtac acagagattc	480
ttcagacctg ctttaaaaac acatgcatac aaatgcactt ctgtctctta ggatctacta	540
actgatgctg cttgcttttag tcttttagct aatattttct ttctttcttt ctttcttttt	600
tgttgaggac agagtctcgc tctgtcgcca ggctagagtg cagcggcaca atcttggtc	660
actgcaacct ccgcctcccg ggttcaagcg attctcctgc ctcagcctcc tgagtagctg	720
ggactatagg cgtgcccac cagcccgagc taatttttgt atatttagta gagacgggg	780
ttcactgtgt tggatgggat gttctccgct tcttgtcctc gtgatccgcc tgccttggcc	840
tcctaaagtg ctgggattac aggcgtgagc cactgcgcct ggctcatatt ttctttatat	900
atcaaaaaca ttcagcttgc ttcactttta tgaaagcttt attatgagtt tgaaagcaat	960
tctgcatttt cttaacattg taactgggtg tgagttgaag gcaggcccct gggagcccct	1020
tgtgggcaat tcccttcaact ctggaggctg cctcgagcct ggacaggcac ttacacttgg	1080
tcagtgattg cacagaaccg gttgcaacag attctgtgca cctccctgtg gcgcgtagca	1140
tttagcagge acttggtcac tatttgctga gtgagtctgt taccttaggc gtgtatttcc	1200
cgtggacctg cctggggatc attgctcatt cactcatttt gaacaagcca atattacatg	1260
tccagggtac gctctatagt gtgaaacaca aaggtaaag atagttcccc ttctcaaagg	1320
aatttctaag gtagtagcca ttcttttgat gcatattctc attctcatag agagtccaat	1380
tatggataat tggacaaagc tgaatgtcgc ttttatgaga atccattctt tctcttttat	1440
gcyttgaaaa atgtgtagca ttcattagtg aattaggatt tcattattca aagaagacat	1500
aagggtcttcg aacagcagat gactgaataa aataatacct aacagcagta gaatgagggg	1560
aggacatatt caaggaacat tttatgccc ttagattggc agaaattttt aaaaagtgac	1620
aataccgtat aaaggtgaac tttcctatac tgatactggg aacatgaatt tgtaccattc	1680
aggggaagaga aacttgataa tatctggtgt agtctgaagg ggcacagtcc ctgtgacca	1740
gtgaggacat tcctcattat ttccttgcc aaacatttca catgagtcta taaggagctc	1800

tatataagag	aggtcactgc	agcctccttt	gtaagagcaa	gaaaaaaaaag	caaataagtg	1860
tttaacaata	ggaacataga	taaattaggt	tatgcagtga	atatttgcac	tctgactaaa	1920
gtgagtgaat	caaaaaaaaaat	ttgtcaacag	gaataaatct	caaaaaataat	attgaaagaa	1980
gaaagctaata	ttacagaagg	atgtgtacag	tatgacacca	ttcatttagt	ttcaactaca	2040
tatctttttat	ggacacatac	atataaaaagc	agaaaaacatg	aattgatagg	ataaacacca	2100
aatattttctg	catatggcca	ggtgtgggga	agtagtggtg	attaagcttc	aaagatgtct	2160
gcagtgggttc	ccattaaaag	tagaaagtag	gctgggcaca	gtgggtcacg	cctgtaatcc	2220
cagcactttg	ggaggccaag	gcaggtggat	catttaaggc	caggagttcg	agaacagcct	2280
ggtcaacatg	gcgaaacccc	atctctacaa	aaaaaataca	aaaattagcc	agatgtggtg	2340
gcgcacactt	gtagtccag	ctactcggga	ggctgaggca	tgagaatcac	ttgagcccag	2400
gaggtagagg	ttgcagtaag	ccaagatcgt	accactgcac	tccagcctgg	gtgacagagt	2460
gagactccat	ccccaaaaac	aagcaaacaa	aaaaagctca	tagagtaggt	aatagtcatg	2520
atatctgatg	ttttttgatt	gtctggttta	catttttttat	ttttattttt	tgagacaagt	2580
ctcacgctgt	cacccaagct	ggagtgcggt	ggtgcgatgt	cagctcactg	caatctctgc	2640
ctcctgggtt	cgagcgattc	tcctgcctca	gcctcccaag	tagctgggat	tacaggcgtg	2700
caccaccaca	cctggctaata	ttttatatatt	ttaatagaga	cagggtttca	ccatgtttggc	2760
caggctggtc	tcgaactcct	gacctcaagt	gattcatctg	cctcagcctc	ccaaagtctt	2820
gggattacag	gcatcagcca	ctgcacctgg	ccttggtata	tgtgttttaa	tttgtattca	2880
ttcattttaag	cctcatgaca	gctctgcgag	gaaagttcac	tatacgtctt	caggctgcag	2940
gtagaggacc	tgaaagggac	aggaggtaac	agtctggcca	agaccacaga	gccagggaat	3000
agcagaggaa	catttcacct	gggcattgca	ctccagagct	gggcttctca	ctgttctcaa	3060
cccttgcaa	atgctcactt	gaacaaagcc	aggtggtgat	acaaaggat	ttgttatatt	3120
agtctctaca	cttttctgtg	tgcttgaaat	aactgcaaca	aagaatatat	cagtatttag	3180
agtaatgggg	gatttgcttg	tgtgtgtttg	tatttttgag	atggagtctc	gctctgtcgc	3240
ccaggctgga	gtgcagtagc	atgatcttgg	ctcactgcaa	cctccggctt	ctgagttcaa	3300
gcgattctcc	tgccctagcc	tcctgagtaa	ctgggattac	aggtgtgcgc	cactacaccc	3360
ggctgttttt	tgtattttta	gtagagacag	ggtttccccg	tggtggccag	gctgatctca	3420
aactcccgac	ctcaggtggt	ccaccacact	tggcctccca	aagtgtgag	attacaggca	3480
tgagccactg	cgccctggccg	tttttttttc	taacaaaatt	attttctaac	agaaagcaat	3540
caggtgagaa	tccacataag	aaacaattta	attcagagat	ttttgttgca	tattaaaaaa	3600
aaaatgtacc	ttcggctggg	tgtggtagct	cactcctgta	atcccagcac	tttgggaggc	3660
tgaggcaggt	agatcacttg	agctcaggag	tttgaganca	gcctggccaa	catggtgaaa	3720
ccccgtctct	acaaaaacta	caaaaaatta	gctgtgtgta	gtcccagcta	ctctgggggc	3780
cgagggagaa	ggattgcttg	aacctgggag	gtcaagactg	cagttagcca	tgattgtggc	3840
cctgtactcc	agcctgggca	acaaagttag	accctggcac	cctgtctcaa	aaaaaaaaaa	3900
aaagtacctc	cttgtaaata	agtaacacta	agacttcatt	tagtggttgt	caagcaact	3960
ccattgtatt	tttattttca	gtttttatgg	ctagtagtta	agggagagaa	gcttggttgc	4020
agagaagaat	gaaaggatga	tgggaaaata	aaagtaggag	agaggaaaac	gcaagaaagc	4080
aagagatctg	tagaaaggga	tgaaggaatt	gtataggcag	agagaatagg	ttctttaatt	4140
gagaaattta	tgttgtctca	ccttctgaaa	tgcccccaaa	ggtaagtatt	tgttttattt	4200
tgaaaagcta	atgatagcta	cctttctacc	acgctgtgtt	caatgtttta	cacactttac	4260
ctgttttagt	ctcacaacac	agtgttatga	ttcgatcttg	ccattggtct	cactttactg	4320

aagaggaagt	ttgaggctca	gaaaagtaag	aaactggccg	aagaccacgg	ttagtgaaga	4380
cagatctctg	atccagttgc	agagtctgag	caataaacta	cttcaactga	ttggtttcaa	4440
agcacatttc	gtcattttac	ttggggtaat	caaagcaact	ctctgaggca	aaattatttc	4500
ctggacttgc	agccatgtca	ctaaggagca	gatgagggtga	gatcacagac	aggatcagaa	4560
tgatggcctg	gtgccaaaaa	gatgtgtcct	agagatTTTT	cattccttta	agaagcagag	4620
aagggagcga	taaatgactt	ttcgTTTTtc	actTTTTtag	acatcgcaga	tggcagcaga	4680
gaatattgga	agtgaattac	cacccagtgc	cactcgattt	aggctagata	tgctgaaaaa	4740
caaagcaaag	agatctttta	cagagtcttt	agaaagtatt	ttgtcccggg	taagtagcat	4800
aatttctcct	gatttaagtt	aaatcacttt	ttaggagagt	gtaagattga	gttctatgct	4860
tttattccat	caatgttcat	cataaaggta	aaagtataaa	acctTTTTtt	atgttttctc	4920
aggcttataa	cagtattatc	tacattttta	attgttttta	atttggccta	ggtttaaaaa	4980
aaatattcct	tactcttttg	tattatatcc	aatgggattt	ttttgccgct	ccaaagaata	5040
tttgtagcc	agtccttata	aagagcatgc	attagataca	ctgaagtgtg	gcttctgttc	5100
tcctactat	cactatgtat	aacttaaaaa	acagttactg	tcagctgctg	gtgtagcta	5160
tctaaaaggc	tatatagtag	gggtcagcaa	actatgccca	tgggccaaat	tctaccacc	5220
tcctattttt	gtaataaaag	ttttgtygaa	acaccgccac	atccattcat	ttccagtta	5280
tctaaggctt	cttttttgca	gacttcagca	gttgccacaa	acactatatg	cctcaciaag	5340
cataagacac	ttactatctg	gccctttaca	gaaaaagttt	gccaaatata	gctctataga	5400
aagaacaaag	tacacatgta	catcaatctg	ggagtctctt	aagaaattat	ccctccctcc	5460
catgagtgtg	aatagcctga	tggcacgtct	gagaaatcaa	atctgatttt	ccctcagagt	5520
ttcacacctt	tctggagtgt	gcagtatctt	attatagttc	tttttgattt	tatggcacac	5580
ttcttttgaa	acatctgatt	ttattttatt	ttttaattaa	ggaaagttaa	attttatttt	5640
cttogaagat	gtttctgaga	attttgcaat	atcttctgag	atcatgaaaa	acagttgatt	5700
tacaaaacca	gagttgggag	gggctgcatt	tgagagctcc	caaagggata	gagtgtgtgc	5760
cgagtgacat	gcggctggcc	gttatgatga	cttgtgaccc	aggggaggga	gttagttgct	5820
gagtgggctt	gagcacttga	ttttccttat	agacgaattg	tcttgtcttc	ctgcctatca	5880
ctcatgccaa	attacttagc	caccagggtg	tttggaacgt	ttaggttagt	gtcttattta	5940
ttttttaaaa	aaatgatgga	aatgttgatt	attttaatgt	acaaatatcc	ttagtagcat	6000
ttctcagtag	ataacatttt	tttcctgagc	ttatttaaat	ggaccaatct	gcttctagct	6060
gatgcctttg	caaaagcctc	cagagtcata	actcgactgc	cttttcttta	tgtagggtaa	6120
taaagccaga	ggcctgcagg	aacactccat	cagtgtggat	ctggatagct	ccctgtctag	6180
tacattaagt	aacaccagca	aagtaagcac	atttctcttt	atrcgacacc	ctgaagaaac	6240
caacaaatag	gtcttgctca	tctcctgtct	acatacctcc	aatcataaaa	cgtttgctgc	6300
ttgcaaattt	cttggcacag	gtggaggact	ggtcatgcag	ttctatcata	acataaaagt	6360
tttacataaa	agagcagatg	gggctgggtg	cagtggctca	acgcctgtaa	tctcagcact	6420
ttgagaggcg	gacgcgggcg	gatcatgagg	tcaggagatc	gagacgctcc	tggctagcac	6480
agtgaaaccc	cgtctctact	aaaaatacaa	aaaattaaaa	attagccggg	cgtgggtggcg	6540
ggcaccgcta	gtcccagcta	ctcgggaggc	tgaggcagga	gaatggcatg	aacctgggag	6600
gcggagcttg	cagtgaacca	agatcacgcc	actgcactcc	agcctgtgtg	acagagagag	6660
actctgtgta	aaaaaaaaaa	agcagtagat	tttcctatta	aaaaaataat	taatattggg	6720
aaaacatcag	aaagtggatt	tgtgaattta	gagaagtata	cagcttaaat	ttttcttttt	6780
ttaagaaaat	tttatttttg	atttgggggt	acatgtgcat	gtttattacc	tgggtatatt	6840

gcatactggt	ggggattggg	cttctagtgt	acccatcacc	caaatagtga	acattgtacc	6900
cagtaggtaa	tttttcaacc	ttcacacccc	ctttcattct	ccccacttg	tggggaaatt	6960
aaatttctga	aactttatcc	tgtagctggc	tctatgatta	taatgaaaca	ttactgtttt	7020
atttaaataa	gcaagtatct	atgtcccttct	tttaataact	tgctttctag	acatttaatc	7080
atatttaagc	ctggtcagtt	caactttata	actcctgaaa	agtgggtttg	ggttttgtgc	7140
tagggaggcc	agctttccct	tctgctacca	gaggactctc	tttggcagta	gtgagggagg	7200
gagtgtttgt	ggaggccagc	tccttaccac	aggcaggggt	tacagtcctc	tgccatccct	7260
cctagacata	tggctttcag	aatttttcta	acctacagta	agaagcacat	ttaacattgt	7320
ggcgtagttc	acaaacacac	atacctacac	attcacacac	aaaattaaaa	gttcacaaaa	7380
caatatttac	tgtgaacaac	atacaatata	tactgatatt	ttgttctatt	ttatttttaa	7440
aatgctcatg	gcaaactact	cagttgtacc	acctactaac	atgatagagg	gagcagtttg	7500
agaaacactt	ccttagatgg	atgagtgcct	ctcaaatttc	aggtgctccg	cctcccgggt	7560
tcaggccatt	ctcttgccct	agcctcctga	gtagctgaga	ctggttaaag	tgcagattct	7620
ggttcagtag	gcagggcggg	gggagccctg	aatgctgca	tttctgacaa	gctccaaggc	7680
aatgctgctg	ctcctggctc	gcagaccgcc	tctggggagt	gaggtcctag	acagcagtct	7740
tgtaaatgtg	agtttctgag	ttaaaatcca	ggggaacata	gtgtcgtcca	gcctccatct	7800
aatacacact	gatccccacc	tgcaattcat	tgcaagtgtg	ggaaggctat	ttgcttattt	7860
gttgtgtaca	gatgaaccac	acaccgcccc	tttcatgtag	gaagttacct	aggaggagag	7920
agatgacaga	tacagaaaca	gccccagcat	caagcagagt	gtggtaggag	cccagaagtt	7980
acaaataaga	gacattggta	acttcagtgt	cagaagagca	aggggaaggg	aagttagggt	8040
tggtcagtg	aaccagggaa	aagggtgaggg	gtgagggggc	agtgatgctt	ttggactaaa	8100
tcttggtgta	ggaattgtgc	ctatggaagt	gaacagagaa	gaaggcattc	tagacagacc	8160
agtgtcaata	gacataccat	gaagacattc	atgtcactca	gtgggctttc	cagtaagcct	8220
tattgcttgc	tttttatttt	tttccaaaag	gcagatctag	gaatatatac	atattcattc	8280
ttcaggactc	gatagttgtg	aagattcttt	taaaaggatt	taaaagtctg	tctaagattg	8340
caatttctag	agtcattcta	agagagatgc	aacttttcag	aagctgcttg	tatgtattgt	8400
atatgtttta	gtgtacttta	catctttctt	ttattcatct	tgaattgaga	aactactata	8460
ttctatttta	tgtaatggga	tcccttctaa	aaaattgatc	acctaggagt	tgcaaagaaa	8520
ccaaatagcc	ctgaaacttg	acaaatgaaa	atggcccttt	cagttgtcca	attaagctaa	8580
gggttagctc	tttgatatga	tttggagggg	tattagtaag	aatttagatc	aacaggtttg	8640
catgatggag	attgtgttct	gtgatgtatt	gtcttagaga	gactttttaa	tccttaaaag	8700
aatcttcaca	actggtgagt	cctgaagaat	gaaaaacttc	agttatgaaa	gtaatcaata	8760
tttcatagta	tgttggggaat	tttttcctaa	ttcttataca	attaaatgta	tgtaaacttct	8820
cccttggtga	aacacatttc	tttttttttt	tttttcaa	taaaaccctc	aatacttggt	8880
acctaaaagg	cactcaactg	tgtaaatgaa	caggtagaat	tcagagtctc	cagtccactg	8940
ttagatgcat	tcattcttgt	ttactctatt	cctgttgatt	tattttttct	cttccaacaa	9000
tttcaatagg	agcaagctgc	tacaattcct	ctttttgaat	attttgaata	tattaaaaat	9060
atattggcca	ctagccacgt	cctgggtgca	gtgttaaaca	tcagtttgct	tgagtggtag	9120
tagttcattc	ctttgaaaaa	gcgtgcatcg	tgaaggcata	caactttaaa	atattgtcat	9180
gattctcaac	aaatgtttga	gcactcactc	catagattta	ttgcatacct	aataaaacaa	9240
taacttatgt	ttgtgtaaca	ttttacaaca	taaaaagtac	ttttggttgt	atcatcttgc	9300
tttgttcttg	aaactcagat	acattttttac	tttaccctct	tacagaagaa	attgaggtgc	9360

agaaagaaat	tatttgccct	gaattgcagc	agtaagtgcc	tacagagtga	ttttccatat	9420
tctaagaata	ttgatacagt	tcttaatctc	aaattatgaa	gtcgaatctc	aacagtagat	9480
cagattcgga	gagagcctta	aaatgtgggt	ttaacatgag	tgaacacatg	tggaagat	9540
aaagaacttg	gttaagcagt	ggagacaagt	tctccagcac	tcacaccct	tagaagctgc	9600
agtaaacagt	cctgttttct	agagagaggg	cactattcat	ggcgttgttc	agaacgttac	9660
agattgtggc	ttatgtcctt	cactcctgca	cttggccagt	ctccccattt	ctccagcaag	9720
ccagcagtg	gtccttgagg	agcgggcatt	tatttaattg	accttcattt	tcttctgctt	9780
ttggtggtg	cttctagatg	gcattataat	cagaacacat	acttagatac	tgcaatgttt	9840
gcccgtgcag	gaactagaga	tttataaatc	ccacatattc	cccatggtgt	gtctgatctg	9900
ctgtgtgtt	gtccccagga	gccatctgtg	tgtgaaaagg	aggccttgcc	catctctgag	9960
agctccttta	agctcctcgg	ctcctcggag	gacctgtcca	gtgactcgga	gagtcattct	10020
ccagaagagc	cagctccgct	gtcgccccag	caggccttca	ggaggcgagc	aaacaccctg	10080
agtcacttcc	ccatcgaatg	ccaggaacct	ccacaacctg	cccgggggtc	cccgggggtt	10140
tcgcaaagga	aacttatgag	gtatcactca	gtgagcacag	agacgcctca	tgaacgaaa	10200
taagatttgt	ttaaatttgt	tgcataaata	gctggggcat	atctgtgact	agccaggtat	10260
gtgcatccca	ggatatgttt	ttgagtgaga	gaaatgagtc	aggctttact	cttggtttgg	10320
agataaaact	ggaagcagtg	acatgttctg	tcgagctgct	tgtgagtata	caagcaatgg	10380
gtacttgtat	tgtcaggaag	caagtgaag	tgagcaaaaa	tggtacctaa	catgcatagt	10440
cattactcct	caaacaaagt	aagagacgtt	gttgactgtg	gaactttgct	gctgtgagga	10500
agagggcaag	cggatgagtc	tccccatctg	aagccctgga	gcaggggtat	aatgggaggg	10560
agaggcgctg	atccttacag	gcagagcaag	agaggatgct	tggcctcata	gggtgacagg	10620
gggtgcttcag	cttctggtec	tagctctgcc	gtgaactaat	tgtgacctgg	acgaatttgc	10680
taaattctct	gaataacaaa	attggagtag	atgttttcta	aaatctctca	ctgtaagaat	10740
tctagattct	tctaacaaga	tttattcatt	gtaatagttg	ggttcctgtg	accagttaga	10800
atcgtctgg	tatggagaag	agtaatcaga	agttcccccc	attccttcca	agtgtccctt	10860
agtgattcat	ttaattctgt	gtgccagaga	ctataaatgg	acacagttat	ctttaaaaac	10920
aactttaaac	aattttaaaa	atctctcacc	taatatgaat	caaggtcaca	cctgtgtaca	10980
gtcgtgcct	tcttctgacc	agcagccgca	gaagtcccag	gacctatgtg	ttcgtgtagt	11040
tcatacacgg	atcattgaga	gtgtgagtta	gtacagaagt	gtttggaatg	ttctgagtaa	11100
agaagtgtga	gcattaacag	tcctggatga	tggagcagag	cctcccagct	ttgttttctg	11160
tcagccattg	gaaagagttc	ttggttcttt	ggaattcagc	ggggtagtgg	tgatcccaaa	11220
agcaggggac	atgtcagaag	gtactgctta	ataaatacac	gcttttagag	acacacatcg	11280
ttgggttgta	gctgtgtaag	tttcttgtgt	ttaacaccct	gtctgcacat	tacttctgtg	11340
ctgcctcacc	actgcctgcc	cactcctctg	ttgttggcgt	tttcagtgat	cattgaaaca	11400
ttcctgtctg	gagagtccta	gttctcttgt	gaagtctgct	gtttctcaaa	agccagagtt	11460
gataggactt	agtatcagta	cttttctttt	ctccatgaag	aatgtagctt	tataatagat	11520
gatgtcacac	atccgtaatg	ggagggatga	ggagatgcct	gtctgtctgc	ctctctagca	11580
tggcccatte	tgttttcttt	cccccttggt	agctcttttc	cgatttatct	acaggaaata	11640
agacattgaa	attcagggca	ggatattgtt	catttttaaag	ggaaatgtat	tttttaaagt	11700
tcagtttttt	tttgcttttg	tttatacttt	aattaaaaat	tttttttctt	gccagttcct	11760
gaaaaagaaa	atagagaaa	aaatattatt	gttcctgggc	gaagtggctc	actcctgtaa	11820
tcccagagct	ttgggagact	gaggtgggag	gttgcttgag	gccaagagtt	caaggtcaca	11880

gtcagctgtg	atcgtgctac	cgcactccag	cctgggtgat	agagtgagac	ctattaaanaa	11940
aaaaaagtat	tgttgggagc	ataaacacgt	gggaaatggt	caagaacggc	cgtcaatata	12000
ctctgttttt	cactgaaaac	tacctttgcc	agagagcgag	cagagatgag	gaaaaggagt	12060
ggaagaagtc	ctccactctg	atagtgttac	tggacaacag	agacaaaagc	ggtgtgctcc	12120
ttccacctgt	ttgctcctg	tccctgtcgg	cgccccctct	cctgctaacc	ccccctgct	12180
ttctctgatt	gctgttttagt	gtggatcctt	cacctgtggg	tgagtctaag	caccgcccag	12240
gtcagtcttc	agctcctgct	cctccacctc	gtcttaaccc	ctccgcctcc	tcgccaact	12300
tttttaagta	cctaaaacat	aattccagtg	gagaacaaag	tgggaatgct	gtgccaaga	12360
ggtgagcaca	ctcacgtggc	aagtttggtg	ttgtctgttt	tcctggggag	ttcacactga	12420
tgaggatgtg	ctgaatgggg	ggaatgtcca	tgcaggaagc	agagccactg	tgtgtgtgtg	12480
tgtgtgtgtg	tgtgtgtgtg	tgtgtgcgcg	cgcgcgcgtg	tgtctttgtt	tatatattgt	12540
cttattttca	gctgtcattt	gaaccaagtt	aattttacta	ttgatgactt	ttnttaagat	12600
tattatgaaa	acagatctta	atggcagatt	ggtttgtgtt	tgtgtttgtt	tttttttttt	12660
ttgagacagg	gtctcactct	gttccccagg	ctggagtgca	gtggcgtgat	ctcggtcac	12720
tgcagcttct	gccttgtggg	ttcaagcagt	tctcctgcct	cagcctcccg	agtagctggg	12780
actacaggca	cacgccacca	tgcccggtca	atatttttat	ttttttttgt	agagatgggg	12840
tttcaccatg	ttgaccaggt	tgttcttgaa	ctcctaacct	caagtgatcc	gcctgcctca	12900
gtctcccaaa	gtgctgggat	tacaggtgtg	agccactgca	ccctgctgca	aattgttttt	12960
ttatacttat	tttcacattt	ccttgcccta	gtggacactt	acatgcatgc	gtatatacac	13020
acacacgcgc	gcgcngtgcg	cgcacacaca	cacacacaca	cacacacaca	cacacacaca	13080
cacacaggat	aacatctgtg	tttgatcatg	tacactgcaa	tttgtgccat	atcagaaact	13140
tcctgattga	tttaggggaa	ttatttttcc	cagtttgaaa	ggaagagtta	tttggaanaa	13200
ggatggattt	tcttttttaa	aaaattattg	atcccatcca	tttaaaatca	aattttattg	13260
gtgaaaatga	aaattaaatc	tcgttcgtga	actactttta	atttcttacc	tagttttctt	13320
ttcttagcat	tagaacaanaa	atgtttcttt	tattttgaag	cttatatttt	atactttgtg	13380
tttttatgtt	tctttatcct	aaactctttt	ttcaacaaa	ctcttagcat	ctcctactgt	13440
aatgccctgc	ggaaaaaact	tcattcttct	tcctctgtgc	caaattttct	aaaattttctg	13500
gctcctgtag	atgaaaataa	cacctctgat	tttatgaaca	caaaaaggta	gggcttaatt	13560
tagatatatc	aagcctgggt	gttactaagt	gttgaatatc	attagatata	caagggtgtt	13620
ttaattacta	ttttgccatt	taaaaaatca	tttcagctaa	atctgttgta	tcttctttct	13680
tatacttttt	tcttactgaa	tgccattttt	aaaaatgtgc	aaccaacctg	ttctctagtt	13740
ttgacgagga	ttagtttaag	tgttgtctta	agaaaagtct	ttgccaagtc	tctgagacca	13800
gtgtttcttg	ttagttagca	tatgtctgtt	tcaaatcagg	atgtctgata	tgttcaggac	13860
gtctaatactg	taagttgagg	ggattgctta	cttacaggta	cataacttgg	gtataaattg	13920
gaagggcctt	ctcaggttgt	cctgtgaata	ggagaaaaa	tttatgattg	tgtttatata	13980
ttgataactg	tattttgtag	tttaaaaaat	acacacgtta	aaacaattat	catcatcaag	14040
tgactgcata	gttattgcct	tgctggttct	gtgtaattaa	attgcaagtt	ttttcatttt	14100
ttgtgggaat	ccttgagagc	atgggcctgt	gctgagcaga	tattcccatg	cacagaagag	14160
ggcagaatgg	ggcccccttg	catcaccccc	tttccccctt	taggcagttt	ctctttatca	14220
aagtggcacc	aagagaggcc	caattggaac	tatgatattg	ggaacatggt	tcttaatctc	14280
tgttacaatc	gaaatcactt	aagggcatgt	aatctttctc	ttttcatgaa	aagaattctg	14340
taagaaagca	gttcttttagg	aatgatgacc	cactgtgagc	ttgatataac	ttctgtgatt	14400

gattatttgt	ttatacaaaag	atagttgata	atttagtgat	ttgttttaaaa	aaatgttaag	14460
ctaacaaaat	cccgtgaatt	cctccccact	agtcataaat	caatcatctt	ataatttttag	14520
ggactttgaa	tccaaagcaa	accatcttgg	tgattctggg	gggactcctg	tgaagaccgg	14580
gaggcattcc	tggaggcagc	agatattcct	ccgagtagcc	accccgagca	aggcgtgcga	14640
ttcttccagc	agatatgaag	gtaaggccgg	tacctgaaat	gaaacctcaa	agagagcacg	14700
ctgacagagg	accctgggag	ccycatcata	ttggtaagaa	agcagagcgc	cgtcctcttc	14760
agtattggca	ggctctgaggc	aatcacaaaag	gtaactaggg	aggggaattta	gaggttaccc	14820
tccatttctt	aggggaaggaa	tttaaaagcta	atttagggta	acctctccat	aaacaggagc	14880
agagctctga	tgtttagagt	ggtcacagtg	ttaaccagcg	gtgaatccag	acaggtctgc	14940
ggcaacctca	cttcttgccct	cctaggacat	aaggcaaaaag	gagagactga	ggcaagtttt	15000
agagcagcag	tgaaagttta	ttaaaaactt	cagagcagga	atgaaaggac	gtcaagtaca	15060
ctttgaaggt	ggttaggcgg	gcaacttgag	agatgaagtg	tgagatttgg	ccttttgacc	15120
tggggtttta	tatgctgccca	tacttccggg	gtcttgcggt	ccttcttctc	tgattcttcc	15180
cttgggggtg	gctgtccgca	tgtgcattgg	cgtgctagca	cacgggggtt	gtgggggagc	15240
gtgcgaggg	tgtttactgg	agttgtaggc	gtgctcactt	gaggcggtct	tcctgtcca	15300
gtctagcatt	cctagaggaa	cgtcatgcac	caggtaaatt	ccgccatggt	gcctcttaat	15360
gcgcagctt	gagcccactc	gcccagctcc	cgagatctta	ttgggaagct	gcagctcccc	15420
agttttaggt	gttttctatc	tactgggagc	ccgcccttcc	ttggtgccc	ctgtgaccaa	15480
cgatcacttt	agagaaaacag	ttgacaactg	cctgaccaac	acctgatggg	cgcctgacat	15540
tgctggtgca	tatctggaaa	gggccctctc	ctgccgtcct	catgtctgac	gagctaccgg	15600
ctgtaaccaa	agcgtgggct	tcggagtctg	ctttcaaate	ccagcttttc	cccttaggag	15660
ctgtgaacta	gaataaaactg	tctaaagtta	ccacctataa	cctgggatta	attatgcctg	15720
ttgccacact	gatagagaca	aggcagcatg	atatcattac	tgatacattt	tttttaaaagc	15780
attcaaaatt	catagtactg	gaaagaaaat	cagtgatgcg	aatgtttcca	gggtaatgtc	15840
acctcccatg	ctgtggaagt	ccttcgggtg	agcctggccc	cttgcttctt	ttgccccagc	15900
ctttctatgt	gggggcacca	tggagctgcc	actcaccagc	accttttttc	cctcaagtag	15960
tttgtacct	taaagtattc	ctgccgtggg	tggccccctc	gtggagctgc	tgagcctagc	16020
cagggtttga	tttctcttcc	tgccagtgtg	agccagatgg	ccacatctct	cttccccctgc	16080
cccgtggaga	ggctctgctta	ccgcaaagaa	gggctcttcc	tcccaggctc	tgtagcacc	16140
tgtttagagg	tgtggagtgg	agcagtggga	accagagcca	ccagagggag	gccctggagg	16200
aggaacgaag	ctgattcatg	tctgaaagg	gtgccagaac	ccaagtctcg	gtgtttaata	16260
aagagtgcct	cgggtgttgcg	gtggccatac	ctcacagggc	atggctcgctt	ggaaatttct	16320
gctcggaat	gctttgtgca	gtggccagga	tgcgttaggg	gccacagatg	actgcttgct	16380
ccatcataga	acagttccaa	gttttcaaac	gagcattcac	agactgagcc	gcacctgcc	16440
tcctgtcct	ctgattcctg	gcttcttctc	tggctctctga	agccacacgg	aaatgtgttt	16500
gcacctgttt	cctgcccttc	agatgacaga	ggaccatgga	agctgctgcc	tccttttagct	16560
ctcttctcca	ggggaattgc	cctcgtcact	gtttgggaac	ccctggctcg	agtcctgtcc	16620
tccgaagagc	ctctgccccct	cctggagtcc	tgagttgaac	ttgggtgttca	cttggcctct	16680
ggctctggca	gtgtgttget	ccttccgttg	acctgccact	gctctgttaa	tgcagattga	16740
tcttcataat	ctgtttctgc	tttaagtgat	taactcaaac	attcttggct	cttattctat	16800
cttgtccttt	gggatatgaa	ccattattta	aatttggact	ggtttctctg	cttggcacag	16860
ttgaccatgc	ctgtaagctc	agtgttttgg	gaggccaagg	caggaggatc	cctggaggcc	16920

aggagttcga	ggccaccctt	ggcaacatag	tgagaccctg	tctctacaaa	aaaataaaaa	16980
ttagctgagc	gtggtgttgt	gaagctgtag	tcctagctac	ttgggagact	gaggcaggag	17040
gattacttga	gcccaggagt	ttgagtttac	agtgagctgt	gatcacaccc	ctgcactcca	17100
gcctgggcaa	cagagtgaga	ccttgtctcg	cggggtgggg	gagtcagtgc	tatacttgag	17160
aagttttttt	ccctcgcata	gtttgtacct	ataaagtatt	catcagtttt	gagcagtcct	17220
tttgcgattg	ttttgagtct	tgattctggg	gaccagtaag	ttgtatatat	ttgcctgtca	17280
agtggacaaa	catggccttt	gtgcctttaa	gtaatggcta	aaagtaccaa	acagaacagg	17340
gcctggcata	gatgctgctc	ctcctgttcc	taggccgtaa	tcacccctga	ttcatcagac	17400
cccaaacaag	tcagctcctc	tcctcctgt	gccccaccac	ccaatctcct	gcaggaagat	17460
gtctggagac	ccctgtcagc	gctaggcaga	gatcaccatc	catgtccacc	tttctcttga	17520
tgagggctcc	cactagcccc	tctggcttgt	gccatgccag	ccatgaactc	accctcatgc	17580
cccacccgag	ccctggcaca	ggctattccc	tctgcctgga	atgctcttcg	tcagtatccc	17640
catggctccc	tcctccccct	tccttctgat	cctgactctc	ccatagcagc	tctctccctg	17700
taacacatgt	tcacaggttc	atcttcgtca	cccatctccg	gcagctcctg	caggcttgat	17760
ggctgctaaa	ggcaggcaag	tcagtggctc	agattcttgc	aaacttagtg	attagtgatt	17820
tctcaactcc	ctcctcatgc	ctcctctgtc	tctataggca	catattatct	cttatctctt	17880
ctcagaacca	agccgcctga	atttctgaat	aacattgttt	aagtgttctg	tgtatgcaaa	17940
agaaaaacga	gaataaaagg	attattaagg	aagaattaat	ataataatag	ccacatatta	18000
tgctcttttt	atactctgct	aagtgtctta	catgaattat	ttcctttaat	tagacaatct	18060
taagaacatc	gacattttta	tgaagcccat	tttacagggtg	ggtgagtggg	ggctgggagt	18120
ggcttaaatac	actttcccca	aaccaggagg	ttagtgggag	ccagaggcag	gacctgagct	18180
cgcggtctg	agctccaaag	ctcattctct	gaactgtgca	cagcactggg	ctgcagccag	18240
agatgcagga	cgctgcggga	ccctctggag	gtggctctgc	ctgtgcttcc	ctcttccac	18300
aggaagctcc	ctataggcat	ctgtgttggg	cgtggactct	cagtgtacct	gcatgtctcc	18360
ctgttgccca	gacaccaaca	ctgaatggaa	aacatgtttc	tgggcatttt	aatgtacgta	18420
cttgccctta	gtcaatctcc	tcgccccct	tccatcctga	ccgcctccct	aatagttagc	18480
agtgggactg	gagcttgaat	ggcaactgat	ttctgtctga	gaggacaaat	caggcatctt	18540
tgtcctctgc	cactgtctgt	tccccatcct	taggatgcac	gatgccagag	ccctccactg	18600
tggctctgta	ccactttgac	ccacactagc	aggtctccat	atgttctctc	cagctgagag	18660
acatcacatc	caaagacagt	ttagagctct	gagggttctt	tccccagagg	tccctgtctt	18720
gtgcaaactg	tctccagcca	agcgtgcaca	agactctgtt	cctgatttgc	ctgggcggct	18780
gagccatggg	cagctgagcc	tgacgcgct	ggactcactg	cattcccact	ctgactttgg	18840
catgaaagac	acacaagtgt	gcttgtgaga	aatagatctt	aacagtacct	tttaaacacct	18900
atttcagggtg	ctcaaaatga	ctgcctgttt	tacatttata	ttctggcagt	gcaaacttca	18960
attggacagg	aaatcttaca	acctctcttc	cagggtgraaa	agcgaggcag	ggatgtttat	19020
acagttccat	ccatgtcatc	ccacttggaa	gatactagta	aaacacacca	acagtaatac	19080
aaaaaccatg	gtgtttgcaa	tagtgataat	gttactagt	aaggaaaaat	agaaactttc	19140
tgtaatttgg	agattctaata	ttttataggt	gggctaaaaa	aaaaayctgg	agagaagggt	19200
gttaagttag	taaggagtgt	gtctctaact	aaatatagt	taaaaagaga	agaaaataca	19260
aagtcaggca	cagtggatag	aggtggatag	tctaactctc	aatagtataa	tgggcaaaat	19320
tgtctcaaac	aaaattagtc	tgccctcttg	ttactcaggg	atgtgtgact	gttttctatg	19380
cacaaaatcc	ccatgaaata	attaagttgc	aagaatctga	actttatatt	ttggaaacct	19440

atctgaggta ggtaggaagt taatttatat ttagaaattt gcttgcata gtctagtagc 19500
tccaggacaa atattcccaa atcccagact attttttttt ctttttaa atcaacagtga 19560
ccagttgggc tcttgtaaga attacagcct taagtttagca aagtctaaga gggctgggtt 19620
taatcctgaa cctcagaggg tccctgcttc tcaaatacta agtaggtcac gtgcacagca 19680
ggtactacat tgaagggaaa ttgtatgata aataggaaat cagcgatttt tacttggaga 19740
cttggcaagg caaatgtttt tgtaataaaa atagatcgtg aaatagaatc ctgaaagctg 19800
cctgtttaaa tgtaaagcaa atggcttttag tgatgcttta agtggtggcag tcacttctgg 19860
ctgccgcaga aactatagaa agtgcattct ctcttggtgc tgtgggttct tagggtgaat 19920
gccttggtg acgctgagta tgtggaagga ccattcattc ttggtaacta tacactaggc 19980
agaggggtggc gttagcgaag ctactgcagg ttgggtgtgt ttaagatttg gatttatatt 20040
tcttttaatt tttattttta gttccagggt acatgtgcag gatgtgcagg tttgttacat 20100
ggttaaacgt gtgccatggg ggtttgctgt acctatcaac ccatcaccta ggtattaagt 20160
ccagcatgtg gttatttttc gtaatgctct cctgctccc tgccgcccc caacaggctc 20220
cagtgtttgt tgttcccttt cctatgtcca tgtgttctca tgattcagct cccatctatg 20280
agcaaaaaca tgtgggtgtt ggttttctgt tcctgcgtta gtttgctgag gataatggct 20340
ttcagcttca tccatgtccc tgcaaaggac atgggtctcat tcatttttat ggctgcatag 20400
tagtccatgg tgtatatgta ccacattttc tttatctagt ctatcattga tgggcatttg 20460
ggttgattcc atgtctttgc tattgtgaat agtgctgcag tgaacatatg catgcatgta 20520
tctttgtaac agagtgggtt atattccttt ggttatgtac ccaggaatgg gattgctggg 20580
tcaaattggt tttctagttc tagatctttg aggaattgcc acaccatctt ctacaatggt 20640
tgaactaatt tacattctca ccaacagtgt aaaagcattc ttacttctcc gcaacctcac 20700
tagcatctgt tgtttcttga ctttttaata atcacgcttc tgactgggtg gagacagtat 20760
ctccttggtg ttttgatttg catttctcta atgatcagtg atgttgagct ttttttcatg 20820
tttggtggct gtatgaatat cttcttttga gaagtgtctg ttcattgagag agacatat 20880
gctcctctga gtaaagggtg aggatgctta cgtctgtgtg acagccttct ctctttttca 20940
gaacctcact gtggatcgcc atcgttggcc tgtactgaag gtaaagcaga tagaggcagt 21000
ctcatctgtc agatgaagac ctcatacacc tgttgattaa gaggctttct tcagatcatg 21060
gttttagagc gtgttttaca aacttgacgt gcttggagtc ttctggaaat cttgttaaaa 21120
ggcagactct gttgcagtag gtccgggtgg gttctgaagt tctaacaagc tcccagtgta 21180
ggctgatgtt tcaggtccac tgtgaggagg cagggttag aataaacaac cgtgggaaat 21240
ccagtcaga tctttgatgc atcctagggt aggcctgtct gtcaggctgc cctgggtctc 21300
tagtgatgga ctccaggagt ctctcaagtc tcaaataagt ctgagtcac agggatattt 21360
tttgagaaga gttgtgctgt ctgaagaagc aaagagtga tgtgatgggg aaaatgcagt 21420
gattaaaaac atggtaaggt ttaaagaaag atttgaccat atgccagggtg aacccaaatg 21480
tatgggtgct tgcgtctttc ctgccctttg gttttccagg gaggcaaggc cttatctctt 21540
atggagcaca ggagacacag tgtgggcgtt tgttttctca gccgtgggct ctaacctaat 21600
tgtcaagcct acaaaaaaaaa aatgctgaaa atcaacttct gactagatat ctggtagtac 21660
ataatctcca taattttctc tctgggtgta ttatgcaaaa gataatcctt tngttattaa 21720
gaaacaattt ttaaggcaac tccaacttt gaaacgggga aaaatcattt tatttacctc 21780
tatgtgctag ggaacaatat taaatttagt tttatacttt tcctttaagc atttcagatt 21840
atattgtgca tttcaccaac aatagaagct ttcagacttt atatgtcttg taaaaaaaag 21900
cctaatatag ataagaataa tttattgatt tgaaacccat tgtataagaa atagtccagt 21960

gaaacttaag ttcaaagttt tttttgtcct gtggatgtag ctatgtcaat atgcctagtt 22020
tatagtaaca ttaagtctag tggattagat attagatatc aattgagatg taagcagtaa 22080
taaacagtaa tgcctaaact gaagtatata atctgaatct ttatgatgac caatttatat 22140
tattgtgaaa aacttaggaa ctgatttgaa acatgattta catgttttac atgaaacatg 22200
atttacatgt gtcatatata gttttcaata atttacgtac cagcaggaaa ttttagtgga 22260
taagtggaaat aaactgcagg tgaaactttg ctggaaaata caagcatagt gacatctgtg 22320
caccaaaagc acctggggag attttttaaa acatgggcca gacatgcctt cctggctgtc 22380
tccctcactg tcagtgaagt tggggatggg gtctgggcct gaattctttc tttttagact 22440
cctcaggatt ctgattgctg ccacgttgag agggttgacc tcaattcgga cctcagaggg 22500
tgacttgaga aactgtcacc acttggtggc agtggtgctc cccgcatctt gattgccctt 22560
gtttctttcc aatcccggaa aagtgtgctt gttttttttt ttttccctgc gtgtttttgt 22620
ttttgatctt gctataatat ttatattcct tgctcatttg caacttattt gaatggagag 22680
ctactttctg aaatctagat gtttttcttt ttctacaggg ttttagggca tgggcaaaac 22740
acggaagaaa aaagttgtct tcagttggca gagacgtgga tttttaagat tgttcttaat 22800
ttactttctg tataactttg cttttctgtg gtgaacaaag accagggttca agataaaata 22860
ttgcaagcca agaactctgat tgttcattga tttctatggg taaagatact tgatcacctc 22920
cccatccgcc ccctaccca cccaccctgc gccgccccca caccctattg tgcttcttgg 22980
cttgtcattt caaaagtcaa ggaagtcaaa gtgaatggca agattttacc tcgacttgct 23040
atttttgtgc ctgttaacaa ttgtgagtta aacttgactg agcttttcct agtgaacctc 23100
cggcgtttaa acagccagtc cataacactg tgtgagggct ggagctaagg ttattgggtga 23160
cacaagatag cacctgagcc agtgctgctt ggtaggaggg ctgaggggaa gagggctgag 23220
ggcttgagat ctgagatgct agagtcacat cgcctggatt tgaatccctg ccctcctgtt 23280
ctgataccag ctgacccatg acgatgctac agcacctgac agcagattcc tccttagggc 23340
tggtctaact ctagagtgtg tgccctgtgt cctgcaggag aatgtccaaa gtgggtgatc 23400
ttgatctgtt aacctttgaa ttttaaccta taccaggag ccattgaaga gtttaagca 23460
agtgaatgac gagtagtttg aaaatatttc cagggtgata gaatttgtgg acatacatga 23520
acatgagcag cctcaaaatc agggctggga ctagagttag gccagcacgt gtccagggtg 23580
caaaatgtaa ggaggcattc actttcaggg cctggcaggt gtggaccctg aacttccagg 23640
accttgagag tgagtgtctc ctaaggatta caccctgggg gcctatttgc ctcacctggg 23700
tccctgggcc tctgtgtacc ctattgcctg cttcagtaaa caggcagccc tgcaagggaa 23760
ggaaggggtg gatcagctct gaggagggag tttttttaga aggatagatt tgttttgttt 23820
aaaaaacagc tttattgaga tataattcac atcctataca gtttgttcat ttaaaatgta 23880
caattcaatg ttgtgaggtt attttttggg atatccacag agttgtgtga acatgaccac 23940
aatctaattt tttttatttt tttttttttg agacggagta ttgctctgtc gcccaggctg 24000
gagtgcagtg gtgcgatctc ggctcattgc aacctctgcc tcctgggttc aagtgattct 24060
catgcctcag cgacctgagt agctgggatt acaggcatgc cccaccaagc ctggctaatt 24120
tttatatgtt tactagagac ggggtttcac catgttgccc agactggtct ccaactcatg 24180
gcctcaagtg atccttctgc ctcagcctcc caaagtgttg ggattacagg cgtgagcccc 24240
accacccgca gtctaatttt gaaacatttt ttgtccccct agaaaaaac ctgtagtgtg 24300
cacttgccaa tctactgccg tccacctcta accatagaca gcccctaata tactttctgt 24360
ctctatagat ttgcctattc tgaacacttc atctaagtgc aatcatataa tatgtggtct 24420
tttgtgtctg gcttctttga tttaacatgt tttcaaaatt cattatgtca taatacatac 24480

cagtaatcca ttctttttta atgacttatt aatattccgt tgtatagaga catcacatat 24540
ggtttatcct ttaccagtcg agaggcattt ggattgtttg cacttttggc tgttacggat 24600
aataccgctg tgaacattga tgtatgtgtt tttgtgtgtt gaatgtgagc tgggtgtggaa 24660
actcctcctc cagggggggc ttacctgtga ttctaccac ggggatgggt aagccagcag 24720
ggatgggaag ggtttgggtc tgcctggcct aggccttccg gcaggctgcc atgtgccttt 24780
cttctgccta ggctgaaacg gaggcctgcc tgggtttctgg cactgccctc gtgagtgtgt 24840
gggaaggctg ggggaagcca agtctccatg gtgcctccat cagggaccct gcagctggga 24900
ggcagccaga gggccacagg ttggtagcat tcacacagag ctacatttct tttttttttt 24960
tttttgagac aatcttgctc tgcgcccag gctggagtgc agtgggtgca tctccgctca 25020
ctgccacctc cacctcccag gttcaaggaa ttctcctgcc tcagcctccc aagtagctgg 25080
gactacaggc gtgcgtgcc atgcccggct aattttttgt gtttttagta gagacggggt 25140
ttcaccacgt tgaccaggat ggtcttcatc tcccagctc gcgattcacc tgcctcggcc 25200
tcccaaaaag tgctgggatt acaggcgtga gccaccatgc ccagcctaca tttctttttt 25260
ttttttcttt gagatggagt cttgctctgt caccaggtt ggagtgcagg ggcaccatct 25320
ctgctcactg caacctctgc ctctgagtt caagtgatc tctgcctca gcctccggag 25380
tagctgggat tacaggcaac tgccaccaca cctggctaatt ttttttattt ttatttttta 25440
atagagacgg agtttttcca tgttgaccag gctgggtctg aactcctgac ctcaagtggc 25500
ctcaagaggc caatccgcct tggcctcccc aagtgcctggg attataggtg tgagccactg 25560
caccaccca gcccgtagct acatttctgt cagctgtttg caaactgtgc ccagaatcc 25620
cctggaggac ttgtagaacc accagttact gggttacgcc cccaaatgtc tgatgctgga 25680
gatgaattat cttgggtgga gccctcaagc cgcagcagct gataagcatg gggacctcct 25740
attctgataa aaattccaaa aaagtcctga gtgattaata aacagcacat tgaaaattag 25800
aaatgagttc tatggcaggg gatgaaacag gcaacaaagc ctattttctt tgcaatgaag 25860
cgcatcagat attaataata gccattgtaa ttatctttat catgtattaa gcattttgtg 25920
tttttcaact ttacacaatt agatgatccc cataggtatt accgcctttt tttttttttt 25980
ttttttgaga cagagtcttg ctctatcccc caggctggag tgcagtggca cgatcttggc 26040
tactgcaac ctctacctcc caggttcaag ctattctcat gcttcaccct ctttagtagc 26100
tggtgattaca ggcgcctgcc accagaccca gctaattttt tgtatctttt ttagtagaga 26160
cagggtttctg ccatgttggg caggctgggtc tcgaactcct gacctcaggt gatccgcccc 26220
cctcggcctc ccaaagtgtc gggattatag gcgtgggtca ccacaactgg acttactgcc 26280
catcttttaa gagatgagga cagaaagatt gagtgcaca gttatgtctc ctgcagctct 26340
tggttcacat agccaggatt cgtatcaatc tatttagctc taaatctagt ctcttaatca 26400
cagtaatgaa ccgttgacag ttttacgagt aaattatcaa gagttttgat aggtttgtct 26460
acttaaatta gtgcttgtag agtaatgggc tgtgttagtg tgaaggaatg tatcttatgt 26520
tggaagtact ctagaattaa atgttaactc ttgctaataa agcatacatt tggggcatta 26580
ttagcaactt tttttttttt ttttttagcaa aattagaggc ttcttagttg agtggtttat 26640
gttatttata tttatttatt tgtttgtttg tgacagggtc ttgctctgtc acccaggctg 26700
gagtacagta agtagcacia tcatagctca ctgcagcctc gacctcttgg gctcaagcag 26760
tcccctgcct cagcctccta agtgccctggg accacagggt cgcataccca cgcctgcta 26820
aatgtttaca gttttttag agacagggtc tcaccatgtt gcccaggctg gtcttgaact 26880
cttgaattaa agcaatcctc ttgcttcaga ctcccaacat gctgggatta caggtttgtc 26940
cactgcgcca ggcctccatg tatttgaatg aaagagcaga catctcctgg aggtggcaaa 27000

gctatgcatg	ccccccctgg	aggggagctg	ggggctctgg	ggttacagtg	atggcacatt	27060
cagggagctc	tccgctttgt	gagatcctga	gataaagcca	aaggatgcat	taaactgctt	27120
ctaaatgaac	tttttccaag	tgaatttggt	atatcacttc	tatataaatg	aaaatatttg	27180
cagcatgagt	actaacaaga	tttttttttt	cttttacccc	gatggagtct	cgctctgtcg	27240
ccgggctgga	gtgcagtggg	gcaatccttg	ctcactgcaa	cctccgcctc	ctgggttcaa	27300
gcgattctcc	tgccctcagc	tcccagtag	ctgggattac	aggtgcgcac	caccacgccc	27360
agctaatttt	tgtattttta	gtaaagatgg	ggtttcacca	tgtaaccag	gatggctctc	27420
atctcttgac	tttgtgatct	gcccgcctcg	gcctcccaaa	gtgctgggat	tacagggtgt	27480
agccacgctc	ccggccaaga	ttttaaacat	tatttaccaa	agtaggaacg	tggttaattat	27540
ggtcttatat	aattctgaaa	atgatttcta	gtaccaaact	atgaatttta	tacttgaaag	27600
aatgatgggt	ttttcacaga	aagttgaagt	tattatgggt	tgtnttcctg	ttcanggtgt	27660
ttttgctgga	gaatgttcga	tgaacagcag	ttctgggtgat	aagttatgga	tgtacacagc	27720
tggtgtgggt	tttaggattt	tattttgcag	cagcatcttc	ctcaaacagt	tgccagggga	27780
aggctttcct	tcttcttact	ggtaccagcc	tttctcttgc	agacaaggca	gtatgggagg	27840
gttgggagac	aaaacagaag	ctggttggtt	cttcagcctg	gcaaggattc	agattgcagg	27900
ttatagattg	gaggccgtca	gtggggatac	ctttccggac	aaagtgggtg	ttctgcctgg	27960
cactgcttgc	cagagaagtt	tcagttcttc	attctccgtc	agagaaaccc	atatggacca	28020
cattctgata	gttttcttct	gtttccctaa	caccgaaggc	tcagcccctg	gtgcagggtc	28080
cagtgtacag	caggctgcat	acagtttagc	cagatgttct	tgtagtacga	aaagtcaccg	28140
agtttccatt	cacttggtgg	tggcagggtat	ggccctcctt	acctcccatg	gcccagggtt	28200
ctctgtcctg	ccgttttcac	attttccagg	ctttcacctc	caggtaccaa	aattcacatc	28260
atttagagat	tgtgtctgcc	tgccaatacg	cggatgtacc	agtgagggat	tgttctcgcc	28320
tgacgagagg	tctggatgat	gagagagcag	agctggccct	ggggctcagt	ggtgacaccc	28380
tcgagcttgg	ctgcttctgt	tcttctgctt	cctctgcttg	gattccttcg	cctttggctt	28440
cccctccagt	tccaagcaga	acaaaacagg	agatatcaag	gaggaaaggg	tgacccctct	28500
atatctggag	agcaaaactg	tcgcggaaat	ccctagtgtg	cttccatttg	tgtctcatta	28560
tctgaaaccg	agttacctgg	ctggtcacgt	gcagccacca	gaggcaggaa	ggtagtgtat	28620
ctgcctgtgt	ggaatgttct	agcattccct	ggtagctttt	gtttcttcag	gcagccatga	28680
ctttgcatag	atcatttctt	tttgcccagg	acactcctgc	tcgtttttct	ccctcctcac	28740
caaaccacac	gtgcattaac	agcgacagac	ttctcctcat	cctctcaggc	cacttggatg	28800
tcaccatttc	ttctctttac	ccctcaggcg	tagtcagcct	ctctgtgcct	gatgttttat	28860
ggctttgtgt	atgccccgat	ggagagcgtc	ttactgtgtc	ttcgggttat	ttatctcaac	28920
ctcgcatctg	tgctatcctg	tacagtaacc	aacagccaca	tatcactatt	taaaattaaa	28980
tacaaactaa	ttatacttaa	atgtaataaa	aatgtagccc	ctcacattag	cctcatttca	29040
agagccacat	gtggctaccg	tattgtaagc	agagctcaag	aacattcagc	aatattgtga	29100
gagtggctac	catattgaga	gcagagctct	agaacattcc	cttatcccag	aaagttctct	29160
tggacatgct	gctcaagggt	gtgaactctg	agatctccag	ccccccagc	tccgtcactc	29220
agaaccacaa	atgtggcacc	atcagccttc	agggtgggtg	ttgtgttgta	ctgtctcctg	29280
actaaagaag	taaacttcag	gcagtcaaga	ttttctacaa	cccacactgc	tcctaaaact	29340
agtgttactg	gatatgtaaa	agctattgag	cccagtgcct	tcaaggaatc	ctaaaagcaa	29400
gtggggactg	tcatgtatcc	aggttccttg	ttttgcagaa	gaagaaatag	aggcttttag	29460
ggaaaggggt	ccactcaagg	tcatacaggc	aaggtcatac	agtcagtggt	agaatggact	29520

ggaatcttga ttgttcatct cactccccctt tccattaagc cataactgat tgagtatcac 29580
caacctgttt gttttccctg attatgttcc cttctcrctt gtttaatcag gtttcagctc 29640
cttgtgagag gaagttgttt tcagcttctt agccccctac ctgtaggcgg tgctctggga 29700
acgctcagga agcagatgca tgtgagcctg tctcagccaa tgattgctta gttgcaagaa 29760
acaaaaaaat gactcaaact agcttaatca aaaggaggct tttaaacagc aagataaggg 29820
taaaggctgg gaactaggaa gttgtcagga accaaggctg tctctctgga tctctctttg 29880
aggccatgta attttttttt cctctcggtc tctttattct gcacaccagc tcaccagct 29940
tgcttggttac tatatgcagt catgccaacc ccagatctg tatgacctgt tagcctcagg 30000
ggcaccacaca tagctggctg ccaatctgtg ttttcttcca ggtttcgaga gggagaaatg 30060
attggcccag ctcaggggtca cttaccagc gagaggtggg ggaagtatgg aggcaccgtg 30120
gtatcagggga acccctgggc caagcttgct caaccgcgg cctgctttct ttcatttttt 30180
ctgttttttg ttttgttttg ttttgttttt ttacagctca tcagctattg ttagtgtatt 30240
ttatgtgtgg cccaagacaa ttcttcttct gctgtggctc aggggaagcca aaagattggc 30300
caccctctgc ctaggccatt atctgggctg tgggaggtgt ggagcagggc cagagctgga 30360
gggggagggc atagcctcca gccaccataa gttggtgtgt tcttggtaat tatattgctt 30420
gtcaaccgaa ggcagaatca ggacaatgaa agtaatgaga atccctagct ttgtaacagt 30480
tagtggttat ctaaaagtag gtgaaattgt acatgagtga gtggcatgaa tttcttatta 30540
ctaaagtgtc cagatagctg gctaactttc tgtcaaagat ccctctgcta ggatcaacat 30600
ttgattaata tatttatcct gtaataagaa tttgggattc ttaaagcaaa atagttgtca 30660
tgtggctgac tacacaacca aagatggctc aggtgtcgct ggaagaggag agactgaaga 30720
gctgttgcca ggttcccacg tggaccttcg gcatgaccg gccatgggga ggctcacac 30780
gctcctgcat cgccacatc ttgccaagcc atggaaaaca cttgggattc atatctaaat 30840
cctagttaa gcttggtgag gacagtggcc tgggtgcagag tttgggtcat agatggtgct 30900
tggtttcttt tgtataaagg ggtatatgat tttggaatat ttaccaaagtg tgggcatttt 30960
ttctataaaa attattgtat ctactgagat tatagtatgt aaaaaaaca tacacatgga 31020
gaaagaatac aaagagagca ttgatattct acagaagtgg caagaagatg tgggtgatagg 31080
tgatattttt gccttttggt tcaattttgt attgtagtga cttttttggg agaaaaaact 31140
aantttctaa ttaagggaga aacatttgaa gtacatttag tctttctaga aaacctcatc 31200
ttctcataga agtttaagat ggagacatac ttccattgtg aatcatgatg ctaaccagta 31260
ttcagatttg ttggaaatgg actcagtttt aaaattgctt ctctcttggt ggctggaact 31320
gcaaatgatt gtttggggat ttttccctt ttctctatag gagttattca acttggcatg 31380
accagtgatt tgagctgaga acatggaacc cttgatttgc agaaatcaag ccccaaagg 31440
tacagataca gtggtcattg tctgaagggt ttcttttggt cttggccctc ctgtccctgc 31500
tcttactgtg gcagctgcag ctgcaggtgc ctctgaagcc ttgccatcca tggtcacttc 31560
ctgcctgctc cccaccacc cctgggaaag agcccccaag tgtccaaaag cactgtgttg 31620
cctaattgctt gttgagagtc tacatttctc tagatctagc agaagtaaaa tttcagtttg 31680
ttatatttat agtttcagga atagtttggg aatggattta ataaaaaatt taaaagccca 31740
tcatttttat atctcttttt cgatatattga tggtttaaaa gacatcaaag ttatcttctc 31800
ccattactca tcctatacaa ttaaaacctg ttttttgaag ttgtaatagg taagttagcc 31860
ttaggtcacc ccatatttat gtaaaactcca gccactgcc acagctactt tgattgtgat 31920
ctgtcattgt gttaccact gtagggcaga aatggttcct gcctcatgcc gttgctgctt 31980
tactcttct gaagtgggtg ggttctgtct ctgtagctct tggcacactg taggttctca 32040

gatggcaggg tgaaggttc ttctgtttgc ttaaattctct cataataccc tgagtctgtg 32100
gagctcaata aattctactt ggtattattg atagattatt tggagccttt tatgttagaa 32160
aagggattct taatccaatg ctccgtttta cagatgagaa gactgaggct caaagaccat 32220
acccccagga gccatgattt gcaactgtatt taggaatagt gtctaggggtc agcacctggg 32280
gttggccgac tgcagagcag cctgggttagg agccctgggg ttgggcgggt ctgggctgct 32340
gggtgccacag cagtctccct cccctgggac tttgggcctg ctacccaccc ctgttccttc 32400
ctttgtgaga tagggctagc agtaactgtc ttgtttcatc agaggcagta ttgcataatg 32460
aatgagagct ggggcctaaa ttaggcacaa gtgcaagccc tcagaaaact atgtacacct 32520
agagagagag agagacacac gtctgtatga cagagaggca gggtttggga atgttctgat 32580
ttcatgtttt gaattgggtg gacctttggg aggatatcct tggaatcgca gagcttcggt 32640
tacatcatga ctttcctgcc caccacatt ttctgagaag ccagagtttt aaatgtggac 32700
cccgtgagct tttctctgtt gcctcatttt ggctgtggc cttttgtttt cttgggtatgt 32760
catgaggcaa aataaaatga aactcagtgc tggttaataa ctcccatcat aatgtatatt 32820
tctgtgaatg gcttttttagc catttgagag gaaaaagggt catgtaaatt tcagaaaggc 32880
ctgattggct ggagagtcag tgtagtgtca cagttaagag tatagattta aaaaaattt 32940
tttattgtgg taaaaaacat aaacataata ctaccatcta aaccatattt aagtataaag 33000
ttcagtagtg ttaagtatat tcacattgtt gtgcaatgga tctgcagaat ttttcatctt 33060
gttaaactga aactctatgc ccaataaaca actcctattc cccctctccc agccctggc 33120
aaccaccatt ctactttctg tttctctgag ttgactact gtagataact catttaagta 33180
gagtcatatg gtatttgtct tcttatttct ggtttatttc gcttagcata atgtcctcaa 33240
ggttcattca cgttgtagca tatgacagga tttctctctt tttttccgc ctttttttg 33300
agttatatct tgttgtagt atatttaaca tttctctcat tcatctgtg acattcatct 33360
gcttcacct tccacctttt ggctattgtg aagactgcag ctatgaacat ggggtgcaaa 33420
atgtctcttc aagatcctgc tttcagttct ttcggatatg taccagaag tgggttgct 33480
ggatcgatca tagtgtagt ctgtgagtaa cctcatact gttttctgca gctgctgtac 33540
cattttacat tcccaccaac agtgcccaag ggctccagtt cctctacacc ctcaccaca 33600
cttgtaatct tctggattgc agattttctg gatcaatctt ctggattaca cttgattttc 33660
tgtgttgggc ctggatgtt agaacagtat ccctccttg gagtggtaaa tatgtaagtt 33720
tttattataa aataatggcc atcctagtga gcgtgagga atatctcatt gtggttttga 33780
tttcttcat agttaatgtg gttgggcctc atttcatgtc ctggctggcc atttatgttt 33840
catatttggg gaaatgtctt ttcaagtctc aagtcctttg cccatttttt aattgagtta 33900
tttgattttc tactgttgag tatggattat taaatcagac tggcctgaac ttaaatacatg 33960
gcccttccat ttttgaccaa aagcagctgt gtgtcccat tgtgccttg cttcttcggg 34020
gtaatgccg cataatgata gcccacctt gtagttaaga gtgttggggc agtcagttag 34080
gaagcactca ctccacagga gcttggtacg taaggagaag gcagccgggt cattcctaata 34140
aggggtctga aggaaggaag aagggtgaa ggaagtaaaa agagcctcct ccatgaatgg 34200
cagccattct tgaaatccac cttggctgcc ttcatTTTTA atgtcagtgg acttttaaga 34260
caacaaaaag gatgttcttg gatgaccaga gactgtggca gaggaggat gggtcacattg 34320
ccaaggatct ctctcaacct cttggatagt gtgctgctgg tagtttgcac aattgcttca 34380
gctttttggc aaagtacatg taaaatcctg aagtcactgc cagaggaaac ctgggtcctg 34440
agatagcagc ttgatgtctc tgccccatcc caggtgcaca cctcactggg cagctctggc 34500
tctgaattga gggacagcaa aaacctctaa ccaaccatac tgaaaagcag gcattggggg 34560

ctttagggga aggttctttt caaaactcat gatggggaga gaccaaagac tgggaatcat 34620
tgtaaagaag ttagtcatag atgcttcact ctttacaatc atcccaacac aagggttaaac 34680
aacatgcagt tttcacgatg tcccagaaaag cgacgagtgc agtgaggtga aacgtggcca 34740
tctgagcaca caatgaccag gcttggaagg atcgatttcc cctgtgctgg ccctcagaat 34800
ttaaggcaca acttttaagc tgagtgtgca gcactcgatt ttctatgttg ggcctggatg 34860
tttagaaaag tatctctcct tcagagtggg aaatatgcaa attttttact gaattacttc 34920
atttaaatcaa agcagccgac ttctcctgcc tcccctgttt ctgtcttggg gttgaatatt 34980
tggtcccatg taacaactct tgattcttaa tgatgccaca tggaagctgt gtgtgctggg 35040
atttgccata ttcagttatg gtcagtagag actttcttag tctctctctc tttttttttt 35100
tttttgagac aaattcttgc tctgtcaccg aggctggagt gcagtggccc aatcttggct 35160
cgctgcaacc tctgcctccg ggttcaaagt attctcctgc ctcagcctcc cgagtagctg 35220
ggattgcagg cacgcgccac catacttggc taatttttgt attttttagta gagacagggt 35280
tttgccatgt tgtccagact agtcttgaac tectgacctc gtcacccgcc tgccttggtc 35340
tcccaaagtg ctgggattac aggcgtgagc caccgcgcct ggccacagtc agtagagact 35400
tttgaaagga aatattacct cttaaatgat gtttttagtc caagtaaatt gtggtaatgt 35460
ttaagaaatt tgcttaccac aaaaacagtt ttcaaggagc atttgaactt gtccacttta 35520
agtcataaaa tggattaaaag tgtttgaaat ctattgggat tgtaaattta tgtcagtgtg 35580
ctgactttca agagatcttg atgatcatgt cgtctgtttt cattttctac tacatgagaa 35640
cattgaagcc tgaaacttaa cacaaccg agttccccac ttgcctaaga gtcattggata 35700
cctaaaaagt atgctacttc ccaagttgat ttctttcagg atatgggccc ttcaaaggaa 35760
agcagtgagg ctgggggttt ccagggtggaa aggtcacatt tccacatata actcagcgaa 35820
cattgtgttg ggttgggaga agaattgggt cactatttta aactttttgt ttcattctga 35880
ggacttcccc atccccctc ctcgcgaaag cacaaaagta ttctctaatt ttaagtcat 35940
gggcttccct taatggattc tgaactcaga tcacgtccag ataagcattg tgtaatggga 36000
tggttgggggt tagatatattt agtcacagat gcatgagagg agggaggggtg gaggacagca 36060
aagtttataa ctggagccta tagtagttta tctcttgtca tcggccagggt cacagagtct 36120
cacttcagga cagctgtgca agcagaaccc ccatcacggt tttcttgatg cctttgacag 36180
tcacctgtac atgcctctgg gacctttcct cctcctttct ctttttgttt tttttccctt 36240
ggtcacatgt ttcattctac taaatgtcta accagctctt ctctgtaaatt tacagagctg 36300
tgatggcacc ttgcttgttg attatttctg gttgaatagt ttccaatggg acttctctgg 36360
agataagtcc tgtattagtc cgttctcaca ctgctaataa aaacatacct gagactgggt 36420
aatttataaa ggaaagaggt tgactcacag ttcagcatgg ctggggagac ctcaggaaac 36480
ttacaatcgt ggtggaaggg gaagcaaaca tgtccttcac atggcagcag gagagagaag 36540
tgccgagcaa aaggggggaa agccctttat aaaaccatca gatctcatga gaactaactc 36600
actatcatga gaacaggatg ggggaaactg ccccatgat taaattatct ctgcctgttc 36660
cctcccatga catggggatt atgagaacta caattcaaga tgagaatttg tgggtgacata 36720
gcaaaccac attaaatccc aagtgcgcat gtctggccct gatcccttta tgtgagactg 36780
gggtcatgat cctccgcac ccgtcttctg agccctattc ctacttgggc atgcttaggc 36840
acttcagcat ctgcatccca ttgatgtctt aagggtggtt ccagaccttg gaggtacaca 36900
cgacacactg ctgatgaaaa cctagaatat agaattggaag ttacatttat tcatagagtg 36960
aaaatccaaa aatagaccag agagaagata tgaaaatatc aagaatgctt atcttaggga 37020
ggtaggatta taggtaactt ttttttccct tagataaata tatagataga tatattagtg 37080

tttacagttt	ctctgccacc	aacccaaaata	tttttttcag	gaggaaaaaa	aaccccagcc	37140
agccaacata	cctaaaaacc	atctcctggg	cccagagagg	aaaaattggg	ctccttttct	37200
tgaaattgcc	atttgtgcc	ctgttgatt	attttaccag	taactccaga	ttccaggctc	37260
ctgtatctga	gttctctctc	cttccacagt	ggagctcata	ccttcctgtt	tcctggctgc	37320
cactcagatt	taggctccgt	ttttcagacc	tcagtggctg	taatagctgt	tccttctacc	37380
tcttaggatg	gttctttctg	taatagcctt	tgtcatcaca	tcacagagg	atgatagctc	37440
ttaatgagga	tctaaaattt	gcaggtaaga	tatccctgcc	tctgacatga	gatagatgta	37500
ttgcatgcta	tttaacatac	aactatactg	agtgtgcagt	tgtatgtaaa	agcattgttc	37560
taggtattgg	gttgaaagtg	gatcaaagtc	tagacaaagg	agcgtacaag	tcttgtaagg	37620
aagacagctg	ccaagagaga	agaaaggatg	gggaaatgct	gcgtctacta	agttcaaggt	37680
tctgaattgg	aaagctgcag	ctattgagga	gaagagtctt	ttaaaattcc	taaagggttt	37740
ttgttatctt	ttattgatgc	aaatgctatt	ttgtggcata	aaccttaata	attttgggg	37800
tgaaactctt	atcaggataa	aatgatcctt	ttctatccca	agcttaataa	atattgttta	37860
agtacaaatt	aaatatatga	aatctgcccc	tctatattat	aaatgtcata	tggcagaaat	37920
tataccttga	cttttgggtc	tttcacaaaa	ccttaatttt	tttttttttt	ttttgccttc	37980
aatgaatttt	gtctgatttt	acattaaaag	cctgtaattt	ctcaagtctt	gagctctggg	38040
agccgtcgtc	atcctttttt	cccctctccc	ttgtcttctg	gatgttcaag	cgattttaat	38100
tagatgttgg	gcttttatgt	caagtgtctg	cattgcactc	catgataatc	cagggaactc	38160
gaagcacatg	ttatgcgtca	ccctgggttg	gtgcagtggg	actgggggtg	gttggaagta	38220
gtattctaaa	tctgcttctt	gcgatgggg	aggtcagggt	gtcctgtgtt	gacaaggaag	38280
aagtctgggt	gaggaagcgg	gatgaaagca	gaccagacgc	tagagtccac	tttcaagtcc	38340
gatcccagga	cctggcttaa	agttaaagaa	cagcaaagat	gaaagggtgc	gcacagcagc	38400
acaggtcggt	ggccacgtta	atgacataga	aagcaagtgc	tgtgaattca	aaagaaagga	38460
cagctctgag	ccagagtact	tgggtgactt	gctcaaacaa	atccctttct	ggcaccacca	38520
ggccttccct	cccgttcaa	aaaaattctg	aattgtgcca	atccattgag	gctcagctca	38580
aggccatccc	atgcctttcc	atcgtaataa	agccttggtt	cctgggcttt	aaacatattc	38640
cttttttctt	aggtagagat	tgaacttttt	taaaagggaa	gttgtcagag	gctctgtaaa	38700
acgttaaata	aaacctgctt	tgttttaggg	atggggtagc	ttggaatcag	atttgctcct	38760
gctatggact	gaacatttgt	gtccccccaa	aattcctatg	ttgaagccct	aatgcacagt	38820
gttatgggtg	ttgaaggagg	gcccttgagg	ggtgattaag	tttagatgag	attgtgtgag	38880
tgaagccctc	atgaatggga	ttactgtcat	cccaaaaaga	ggtagagacc	ccagagcttc	38940
ctctctcttc	accctgtgag	gatacagcaa	gaagggaagc	ctctgcaagt	caggaagaga	39000
gagggtctct	actagaatac	acttgtagct	ccaccctgat	cttggaactc	ccctccagaa	39060
ctgtgagaaa	caaagtgtgt	ttgtttaagc	caccaggtcc	ctatgatttt	attagagcag	39120
cccagagctc	attctccact	ccctggcttc	ctgcatggac	tttgcaacca	gagcttcacg	39180
gggtatagtt	taatagctgt	ttctctgtaa	cgtagccact	tttctcttct	cagggtctagt	39240
tttgaccctc	ataacacttt	gttaggggag	atttgagggt	gaggaagttg	gcttgctttt	39300
cttttcacca	tgtctcagta	gaaacagaag	cagaaaggcc	ctgagatact	gagcccacct	39360
ttctcagcag	ggtgtgacag	cccggagtac	cctgggctga	ggaggccagg	gctggagggg	39420
aggctccac	ggtggagggg	ttgaaagctg	ggttgtaatg	agctgctttt	ctgtagatgc	39480
ctaaatgatg	tggtgtgaga	aatcgtgatc	ttagctttta	gtagtatatt	tttctgttta	39540
tgtaggtga	gtcatcagtc	tgtctctgac	tatgttcaga	tctggaagtt	ttctggaagg	39600

aaatttggtta	ttgctgtaat	agtgtagggt	gttgatctgg	attagcaggg	agcggccccct	39660
taatacatctc	ttaagaaaat	ggtatttagt	tcagtccttg	gctttgaact	ttgcctttga	39720
caaagatgaa	agtgcgactt	gactgggtgt	tgaaaaacat	ggtgatatgg	ccaggtgtgg	39780
tggctcatgc	ctgtatccca	gcacgttggg	aggccgaggg	gggcagatca	cctgagatca	39840
ggagttcgac	acctgacttg	gtcaacgtgg	tgaaacctca	tctctactta	aaatacaaaa	39900
aaattagcca	ggtgtggtgg	tgtgcaccca	taattgcagc	tacttgggag	gctgaggcaa	39960
gggaatcact	tgaacctctg	aaggcggagg	ttgcaatgag	ccaagattgt	gccattgcag	40020
tccagcctgg	gcaacaagag	cgagactcca	tctcaaaaaa	aaaaagcaag	ttatattaca	40080
ttttaaaact	ctattttaatg	gtcagggtcat	ccatccataa	tgggtagagt	cattgcttaa	40140
ttaattttaa	acaatgtatt	taaaagggtac	ctttgttccc	tagtgtcaca	taacgtgaaa	40200
tatccaatta	aggtaactgt	aatgtaaaagt	aagtggctaa	aaaagtgtctg	aacgccaaag	40260
gccagagatt	caaccttttg	tgtgcattag	aatttcccaa	ttgttcaaata	ccaggttget	40320
ggatctaccc	cagagttttt	gatccagtag	gtttgggggtg	ggaccaagaa	tttgcatttc	40380
taacaagctc	ccaggtgggtg	ttgaggctga	agctcgtgtg	gggaccacat	tttgagaact	40440
tctcccgtag	actgaactca	tgggtctaggt	tctgtcagct	gtgacctctg	tgtgtctgga	40500
gggagtggtc	agatgtcctg	acctctgtgc	ccacagttag	gtccaagctg	agtaggtttg	40560
accagcagct	gtaatcacag	agtgaacaat	gtaaacgacc	aatgttgggt	gggtctgacat	40620
cttttaaaaa	aaatccacgt	ggatgagatc	acaggggttaa	gtgtgggcag	cagtcaggggt	40680
aactccatgt	ggttactgcc	catgcactct	ctgctgtttt	tcacctcttc	ttcagagtgt	40740
ggtcaggatg	gtggccttgc	ccagcacagg	aggccctttt	ccttctgacc	acctgacctg	40800
accacactct	tagcatctgc	aggcactccc	tgtcccttcg	ctgggccccg	tggggaacta	40860
cttgtagtca	tcaaattcat	catgctgctt	tcttttaatt	cccacacttg	ccaaggtggg	40920
actgccccgc	atctccttcc	cagtcgtgtg	tcagaactca	gcaactggacc	tttccccctt	40980
ccccactccc	accctctctc	accccgacga	acgtctcact	tgggatcatc	tcttctgagg	41040
ttggacctgc	acagccgccc	tctgcaactct	cgccacctta	tgggctgccc	ttgacctctt	41100
ggcacacaga	cctggaagtt	ggcctgctca	gctgtctcct	taggggtgga	gcttggtttt	41160
ctttcatcac	tgttctgcga	tgaattgaat	gcatgattgg	tcacaggaag	gtaggggagg	41220
gataaacacc	ttatgatatg	tttcttataa	ggtttttatat	gtagaaagtt	atatgaaagt	41280
gtcagatatc	tatatatgaa	gtatatgtga	agttttatga	tagttttgca	taatttaaga	41340
ataaactctt	taaaggagct	gagtcccaat	cccttgggtc	gagagtgtcg	tggctccccg	41400
ggcctgcttg	tttccctcca	ctctgcgtgt	tcgttgctgg	cccccatag	gctgtcccag	41460
acctctttga	cttctctcct	ttctgcccag	tcttccctga	gacgtctcag	gctccctggc	41520
ctctgcttc	tggagcttc	tcttgtgttt	gttttctgtg	ctcagggcgc	catgggtgcta	41580
taggccacag	aggaggcgtc	tggggctcct	cggggcaggt	gcagcaggag	gaagccgtct	41640
ccgagggcat	gaccttgga	ctgagcattg	acagaggaga	gtcagccaga	caaagaaagg	41700
ccaaaacccc	accctctctc	cacctattt	ctacgtgacc	atgggcccctg	gacacagcaa	41760
gacggtgacc	ccgggcctcc	tattgttgcg	aggagccccct	gggaaaatgt	tggcattttc	41820
ttcatagaac	aggtttctct	tctccagtat	tcttcagtaa	atcaactttc	ttttttatcc	41880
ccaaccccag	tctgattgctg	aagaagtcta	agcaacagaa	agattttgcc	aaatagatta	41940
tcttttttag	aacaaaatag	atcatgatat	taataggaat	tcagcaactta	ctcttgtcta	42000
agtactgttt	ttaagtgtctc	tcaaggattt	ttcattttaat	ccccacaaca	aagctgtggg	42060
gggtggatgc	tattattatc	ggtgatttat	gaatgaggaa	actgacacag	aggggtgggtc	42120

gaggagcttg	cccatttcct	ggtagttagt	accagggctg	gcacatcag	ttgcctgctc	42180
cttttcctct	ttgcttttgt	gtccattacc	ccaaggcatt	aggatgagcc	agccaagtcc	42240
tagtcctgga	ttcaccacct	aattagctct	gtgtcccatg	tcttgccgtg	gagggataaa	42300
accaattcct	agcttatccg	ttggtggtga	agatgaaatc	agtgggggtac	ttgtaaagca	42360
cactgcccag	cacatagtaa	gtgcccagaa	aatgtgacgt	cggacctctt	taagcttcag	42420
tttccacatc	tgggaagaga	gggggagttg	agctaagtca	ttttccagt	tccctttcag	42480
ctccatgttc	ctgtgagcac	tgacagtttc	cccacaatmc	tgaagaaaga	aggaaaataa	42540
ggggcggggtg	gcgaagggtc	ccactgtgac	gtggctgctg	gtgggaagtc	cctggggagg	42600
caaggcccag	cttcccagac	acagccctca	ggtgctcatc	ctggtggcac	tgaccagggg	42660
ccatggtggg	cttttccacc	ccaccatgtc	tcataaaatt	acaagaacca	cagttgaaaa	42720
tcagtgttac	agaaatggta	ataggatagg	gcaaactgtt	acaaagatca	gcacttaaga	42780
ttctggctga	ggcggaatat	ttgtttctct	ttagttttgt	tgtctttaat	caagaactga	42840
gagccctgac	tttcagctcc	tcaaaaaata	cagcttcctt	ccccttgcat	atgcaaaaac	42900
aaacgccact	tctttccaag	cataattttc	tcccatgctg	tatctcctgt	ctacagcttt	42960
ttcttgatcc	ttctccagct	cctgtagacc	tcccatttag	agccaccagc	cgcccatcac	43020
tggggctgctg	cagagctctt	ggtgctctgt	gccctgggct	cgcccaccca	ggcctgttct	43080
ctgngectct	tcctgggtct	cttccctgga	cttcccactg	ccgtgtggnc	ttcagtgtct	43140
ctctgagctg	ttgtcatgac	ctctaaccag	actgagtcag	gacttttttc	ttcctcatct	43200
ctaagtcatc	cttacacagc	cttgggaagt	taccctaaat	ggctattttg	ggagggagtg	43260
gggataaaga	tctgcaggcc	tcttgctcct	ggtccttggt	tctgcttate	ttggcttctg	43320
tttttaagt	tgtgtgcacc	tctttcctca	tcacaccctt	cccctccgta	tggctcccat	43380
ctcaggcaga	gttaggtgct	ctgttctgtg	tccatagctc	tttttcgagc	ccttcttctc	43440
actgtttggt	agtggccttt	catgtgtgtc	tgatccacta	ggctgtgcac	tccctgcctg	43500
ccaggatatg	gttaaagtgc	taaagaatgt	atatatgaga	tcacttttgc	ttaaaaaacc	43560
cccaatcttc	tgggaattccc	aatttctaac	caattaatat	gtggattgac	tagaccttaa	43620
gcaaccaaga	gtcagccagc	cttgtcttct	atattcaggc	gcatactatc	tggtcggttag	43680
acaaaatggg	tcattatcag	tgatgagtta	ataattacct	gcacatcttg	tttatgctgg	43740
ttctttacct	aaagtggctc	ccatcaatta	aacctgtatg	gattttacct	gttcttccag	43800
aaccacccca	ctttccacaa	aaactgacaa	caatgatggt	aagaagaatg	gtagttgaca	43860
ttttattaaa	tgtttactgt	gtgcaggctt	gtttttttcc	acacatttac	ctacttaatg	43920
ctcacaataa	tcctatgaac	tagtcagttt	tatgcagatt	tcgcagatta	ggaaactaag	43980
gtggcaagt	atcagataac	ctgtttgagg	ttgagtagct	agatcatggt	agagccaggt	44040
tcaatcccag	atacctggct	ccagggccca	tgtccttgac	cttataaacg	gctgaaattc	44100
atcttttttt	gctgaacttc	cagaacactt	tctttgtatt	tcccttattt	tggtagtctt	44160
gtactttctct	gctaccctga	ttcatacttg	gattttctagc	agcatgcctg	gcacgaggca	44220
acaacttaac	agtatttctt	tataccaaat	gaatgttgct	tttttttttt	ttttttttct	44280
tgagacagag	tctcgctctg	tcgcccaggc	tagagtgcag	tggaactatc	ttggctcact	44340
gcaagctccg	cctcccgggt	tcatgccatt	ctcctgcctc	agcctcccaa	gtagctggga	44400
ctataggcgc	ctgccatggc	gcccggctaa	ttttttgtat	tagtagagac	ggggtttcac	44460
cgtgttagcc	aggatggtct	cgatctcctg	acctcatgat	ctgcccgcct	cggcctccca	44520
aagtgtggg	attacaggtg	tgagccacca	tgcccggcca	tgaatgttgt	ctttaaaaaa	44580
ttctgttttc	ctctagctag	actgtcatat	aatgcaactg	taggaaataa	tcaggttctc	44640

tttggagtat tttccataaa agatccacag aagtcatggc aggggttgaga gtggacttgg 44700
gcaaataaat ctgttcattc attgaatatt ccatgcatat ctgctgtttc ccaggcatgg 44760
gatatggcag ggaacacaga aatctctgcc tcctgggctc tgctttctgt tgtagtagag 44820
gtaaagctgc tcatactttg taaacaatat gacaacatta agtctacatg gtcattttac 44880
tttggttttt tctaagaaat tttgagctgt tcgtaacaac agacgctgca gatgttaatc 44940
ccgttggtgt taacttttct ccagagattt aatgttcaat tttctccttt ccagaatcga 45000
tttatgttgt tcaaacagag gtttgagaat aactggaatt tttttaactt cttttttttt 45060
tttcgcatgg agttcagaat tttcaagagg gatgaagaga gttataaaat gctctatggg 45120
gggtaacaca cagaaaaagc cagaaaattg gagaataagg atctgtctac tcgtttcctt 45180
ctagagctcc tctttcttac agggcactta acatgtgatt taatgtcgtg tctttaaaag 45240
gaggagaact gcagttcaga acttaatgtc agtgctttgt gaaagtgcaa gaaagaagcc 45300
ctgtattctg cacttgagag agccagatac tgggcagata ggaggtgggtg tgcacgttgc 45360
tttttgtctt tctcgatcat ggcattgatt ctgttcataa caatgatgca atgtcatcct 45420
cttccccaca catttggtgt cagatagaaa gaatgcaaca gcacagagtt gttggggaat 45480
aatttggeat ctaaaatata gacataccag catagatcat atttatgact ctggtgggag 45540
tgtcacagca atgatttaat aggaggcagt tgtctccaag gcctcctgaa ttatgactgg 45600
ttttaaaatt cttagaacct attggaggct attgtttctg aaaggctaca taatttaagt 45660
gctccacatc cgtcattata ggagatgtca gaatagtaaa atctaactct ggactaagtt 45720
gttatcgctg ccctttgggt tgggtggcttt gccgacttta taaatatgcc tgtcagtgcc 45780
tgtgggtctc acagttgggc agtcggcggt gaatatcatt tctcacattt tacactgggg 45840
gactggaacc cagaaggcat atgttttccc aagaggcacc aacacagttg gcccatgagg 45900
tagagcagcc cctccttcgg ctcagcctcc gctgcactga gccaaagcaa gcttcctaca 45960
ctggcctctg tgcagctgtc tctcagcaag aatgcaagtc ggggagagaa gccggatccc 46020
tgggattgtt ctagagagta gaaacctcag agtagccctc cttagaccac ctaacgcatt 46080
gcatcgctgc atacatgtaa gggactcaat gctggtagga ttggcttagg aatgatgcaa 46140
gtgaaaacag tgccccggtt tatcattaga acaaggttct tagctgacag ttgcctcaga 46200
ctttgatttt gttctccttg acctgccact ccactcgagt ccacatctct caagactgca 46260
cacgcctgaa ggaggactga ttacaaacca aagccttggtg ccagctctgg atctttttgc 46320
attgttgaga aagcagctta ctttcttttg actgattcag caggccaaat ttagaacaaa 46380
gatttttaac tatctccctt tataaattac tgagctatth tgtagccagg ctactcttaa 46440
tatgaacaaa aatatattata caaatttggt gttaatcgta aactataaaa aaatcagtaa 46500
ttgttaccac gtgaaatgaa tttggataaa agagatacgt ttttgccctt tcccagggtt 46560
taggagagac gaaatgggtga gatttttagct ctgaatcaga ggttcttatt agaggtgggt 46620
ttgttctctc tgacctctag gggatattta gcaatgecta gaggcattga tgggtggcag 46680
atgctactat gccctctgct aaacattcta cagtgtataa aactgttcct cctgacaaag 46740
aatcatccag ccccaaatg tcagtagtgc tgagggttag aaaccctcct ttaactctt 46800
gggtttatth gctgaccttt acagtggatc agcttttatt tagttcatgt agaggtgaaa 46860
ttaatactag tgctcaaata tgtctttgta ttctggactt ggcctggatc ccccgaccaa 46920
atthgggaca agctcctgcc atgtgttgag gacctgaatt caggcagcta acaacagtat 46980
ttgaactgtg ttttcagtgg tgggagtga ggaatgagc cgacgtgcta gcaagcgcat 47040
agggttgcat gaggaaatag agagtaaagc tgcagcgtgg agccctgcta ttcagagtgt 47100
gcttgagaaa acagcagtgg aggcattact ggggagcttg atggaaatgc tcccctcaga 47160

cttgctgaat	caaaatcttt	aatttagcaa	gatccccagt	gaggcttgtg	catgtagaag	47220
ttagagaagc	acggggtaaa	ctcttctttt	ttttactttg	gaggaaaata	cacctttttt	47280
cttattatgg	ctctgacctt	tactagctgt	gtgaccttgg	ccaagttata	aaacctcact	47340
gcaccttatt	tgttttagct	ggaaaatgga	gatcataata	tcacctgtcc	tatgagattg	47400
ttgtaagaat	caaacaagct	tatttatgcc	aagaacccat	atggtaaaag	ctcaacaaac	47460
tgtcactagt	gataataaga	aaaagatcac	aaaagtagaa	aacattaggg	agacagctta	47520
ggctctaaat	ctcacagttg	tcgtccccc	acaatacttg	tattttttgca	gatccagttt	47580
ctctgaatac	taaaataaaa	cgggagtttc	ataaacttct	atagacagtg	gtccttgtca	47640
gtagcccaag	tggcagagag	tacatggatc	tggggacaaa	cagcctctac	tgtaggaat	47700
gttccatcct	cctggcctga	gttacacctg	ctcattgtga	ttccgaattt	gaaaggaaca	47760
cagtaggaat	tttcaagacc	ctgggaagag	gaaggctgtg	gtaaacagga	aggatgagat	47820
tagaagaagg	agtttaggtg	aggtgagccc	ttgttttact	agtagggttt	aagaatatcc	47880
aagtcagctg	gacatggtgg	ctcacacctg	taattctagc	actttgggag	gccgaggtgg	47940
gcagatcacc	tgaggtcagg	agttcgagac	cagtctggcc	aacatgggtg	aaccccgctc	48000
caactgaaaa	tacagaaatt	agcagggcat	ggtagggcat	gcctgtaatt	ccaactactc	48060
actcgggagt	ttgaggcagg	agaatcgctt	gaacttggga	ggtagagggt	gcagtgagcc	48120
aagattgggc	caccacactc	ccacctgggc	aacagaatga	gattccgtct	ccaaaaaaaa	48180
aaaaaaaaaga	aaaaaaaaaa	aagaatatcc	aggtcaaccc	cacctaaccc	tcagcggggc	48240
tcccttctgt	tgcttgggtg	ggctcctggg	tctcttgacg	cacacgagat	tgtgagagtg	48300
tatggaaaca	ctgccctcgc	tatcaggaca	gcgcctgcca	tgccagccag	aacacatcat	48360
aggaattgca	aaactctttt	gcaaaccagt	gagagatatg	cttccaatgt	gaggtaaagc	48420
agaacttta	tcacagctgc	agtgttcac	agaattccaa	gagccaagat	ggtaaaagaa	48480
taaaaaaaa	gaaaggaaag	ggctcaaatt	aaagacttca	agctgcagaa	taagattaaa	48540
taaaaggatt	caattgaact	gcatcatatt	cagtaatgac	taatcctaag	tatacagggt	48600
ttgggggtga	aaggatttgt	aagtgtttt	caggaaaata	ttttttccat	ctttcatttt	48660
aattagaata	gatttgcatt	attttttctt	agtttttatt	tttaaaatat	ttattgccac	48720
aaatttagaa	aatacaggga	aaacataaat	aacagtacat	gtaaaccaat	attttgtccc	48780
ttcttttgtt	caacagctat	ttctcaggca	cctgctgggt	gtcagcagct	gtgctcagtg	48840
tggtgaccaa	aaccttgtc	aacaaggcag	caaggttcta	acctgggttag	ggcttacagt	48900
tgagtagctg	aaattttgat	ttcttttctg	tgccctagt	aaagatatga	tagcaaaca	48960
taagagctat	tttttttatt	gtgttcttac	tctgtgttgg	gccctgttct	cagtgggtta	49020
tagcctatta	actcagtctc	tttaccacca	ctctgagggg	aggctctgtc	ataccactt	49080
gacagatcgg	gaagtggaag	catcaggagg	ttaagcaact	tgtaaaagat	cacaaaatca	49140
ataatgacag	agttttgatt	agaatcccag	cagcctgtct	ccagaacctg	ccctattaag	49200
tgcagtgcaa	ctgtactgcc	tttcataata	tgtatcaa	tgagatgata	ctttataatt	49260
tcaattcttg	cttttctatt	gaacagtaca	cagtaacatc	ctcctataat	gcatataaac	49320
ccccaaaaga	tgtagaattt	taattttatt	atttgtctga	taggctcata	atgaaataag	49380
actctataaa	gctgtgta	ttagatatag	gaaacatttg	gattatagtg	gtatgtagtg	49440
ggaacaaatg	gtcttctgaa	tcaggaagac	atgagttaga	gtatgccggg	gtacctcctt	49500
actcactgta	tgaccttggg	caagtttctg	aacttttagt	tcctttccag	gctaataatc	49560
gccttctgga	cttgtcatca	ggattaaatg	agtctaccta	tataaaatgc	ccagcgcagt	49620
gcccagcacg	tggtagaagg	tctgctagtg	gttactgtta	ctgctggcta	ttaaatacat	49680

tttaatcttc cttcagaata cctggccaga tagcacagtg gttaagaatg cacatgaaag 49740
ccagactggt ggggtccagt cctggctcga ctctctccta gctatgtgac attaggcaac 49800
ttacataaac tccttgttcc tcagtttgca tttctttaaa actgcatagt tatcataccc 49860
atgtcttaga gttttgtgag tgtaaattat tgtatataaa gctctgagaa cagtttggtta 49920
cacagtaggc actgtatgaa cattttctgt aattatcaat aatataatta ttaaataaca 49980
ttttcagaag gagataaaaa tattacacct taaaaagcag gtatctttta attcttcttc 50040
agctactgaa gttttgctta ctatttgaca tatcatttgt ttcacgtttg tggctcagac 50100
gtggcttatg ccaatgcata ttaacacagg aattttaaat ttggtgatat tattatatatt 50160
tatctgaatg aacagaattt gctgatttga cactgtgttt gaatgtgcat tttttgttga 50220
aaaatgacaa ttctggaatg ccgtctccct ttccagatta ttcagagctg ggagagcttc 50280
ccccacgac tccttttagaa ccagtttgtg aagatgggcc ctttggcccc ccaccagagg 50340
aaaagaaaag gacatctcgt gagctccgag agctgtggca aaaggctatt cttcaacaga 50400
tactgctgct tagaatggag aaggaaaatc agaagctcca aggttggttt gccatcttga 50460
tattgaacag gcctggctct atcttggctc tgaagttaat cacatcagac ataagcatgc 50520
tgtcttaaaa atacagcagc acgatagtct aatgtataca tctatctata tctgtttact 50580
ttttcagagt aatattaaca ctgtttactt tctggtgatc taatgatagt ttcaccaaca 50640
atattcatta ttctctatg gtcactgtta gtacagtgtt tagaacttct gagatccaag 50700
ctttaaatct aagctctaac acgctgaaag gtgcttttca ttttgttttg ttttccccctc 50760
tgtctctctc tctctctcta ctttatcctc agccatggtc tgtgcctgtg tgttaggtat 50820
gaacttttct tgtgtaagtc attaacatac gtaacttcac tctgtgtgct ttttcagtga 50880
tttgcaagta atctgaaaaa aaagaattag ctgagttcta cctgtactga tatcaatagt 50940
gtcaaaaatat gacatgaact ttgaaagttt agattttgtt catttcctgt ttccatgctg 51000
acactggaac caattaatgt tatcttcaaa gtagcttaag atgcaaagtt tacatactct 51060
ttggaaagag catgagtctt agggatatcta gagaactgcc cgggtgataaa gtagtgaaga 51120
ttttgagcag gaagtctgca taatctcttt caaagggag atgtagcaga tggttcagtc 51180
accctgccat tgccccagaa caattttgga attacagtac atttcattca gcatcattct 51240
tgattgcaaa ttttgatctt ttaaaatgac cttgatgctt gtatagagct aaaaagtcac 51300
taagacacca actctgagga ataagctcct gagaatgtgt tgcactctgt agtttcagtt 51360
gcatagctag tgatcatagc agtgataga cgttctctgt gcatgtccct acaatgcttg 51420
tgagttatga caaactgtg tacgagcaac atagtttctg cagttgaaaa gtacgaattc 51480
atagaatgta aagagatagt gtctatatct tttgactgaa aacagaaaat gagatataaa 51540
ggaataagac ctttcgacat gaaagtaacc ccacagttgg aataggctag taagctttcc 51600
aacatgcagt tttgaagctg agaaagacgg gtccctctcat cagggtgctg tggaagatga 51660
tagcacactg gggggcggtt agagcaggtg agtgctgttt tcttccaacc cagtttttct 51720
gccactttct tatgtttttg tgaaggtaat tttaaaagca gatgtctaaa agatgtttgg 51780
tagtgatggc attactgcat gtgtcatcag ttaaatgaca gctcgggagc acagcagtta 51840
tgttcgtgtg tatcttggga tttttgttga agagggaaaaa ggcagttatg ttcacatgt 51900
aggtcaaaact ttaatgcaa tactggccaa tattcttgca aatgacagcc atgtaaaatc 51960
agggcatagc tataaaatgg gaacggtgct cacagctggc ttctttgtgg tgaggacagc 52020
tataattggt gaggcaaaac cagtgtgcca caaaagcaga atacattctg ctgtgcaagc 52080
aatgaccaga cagactagaa tgaaaaggca agagtttctc aaggttacct ggaaccctt 52140
gccaggtgtt gcattaagtt tactggccct tgccaacatt cttctaattgc ttcctcattt 52200

catctggctt cttggcagtg ttcagttttt gtggtctttt atttttactg tttgacttca 52260
tttctcttct tagctctgta aagttccaca tgtgtttatc tttgtggtga aaacacaata 52320
aacttgctta atataatggt ggaagtatta atccattgta ttagtgtgta caggacctgg 52380
attgctgata aaaaaataac tagcaataac agcctgattg cttaaaaaata tttagtaagt 52440
tttgtcgggg tggattgggg cagggcagaa cttttacatt aaatatagat gcaagatttg 52500
ataagaatca gccagagtggt acagtaagta ttcacttaat gttgccaata ggttcatgga 52560
aactgcgaat ttaagcaaaa tgatgtataa tgaacaaaat tttactaagg gtttattgat 52620
aaaaacaaga gttaagttcc tatggcatat ttctgggcac aaaaacatca ccaaacttct 52680
aaataaagac ccaagacact tctaataatta aatattgatg taaacgtgag atatgcaaac 52740
atttaagcaa gattaataca aatatgataa ttattggctt ggcacagtgg ctactcctg 52800
taatcccagc actttgggag gctgagacag ggagatcacc tgaggtcagg ggttcgagac 52860
cagcctggcc aatgtggtga aaccctgtct ctactataat tacaaaaaaa aattagccag 52920
gtgtggttgt gcacacctgt aatcccagct acttggaaga gtgaggcagg agaatcgctt 52980
gaaccagga gacgaagggt gcagtgaagg aagatggtgc cactgcactc cagcctgggc 53040
aacacagtga gactccatct taaaaaaaaa aaaaaaaga aagaagtaat tatttttcca 53100
cttattccac ttcagggctc cagggggcca gaacctatcc ctacagcttg ggatgcaagg 53160
caggaaccag ccctggaccg aatgccattc catcttgggg tgactcacac acacactcag 53220
actgggacca tgtagacata ctgattaacc taatgtgcac atctttgaga tgtgggagga 53280
aactggagca cttggagaaa acccacacag acatgaagag aacacaaaact ccacacagat 53340
aatggccccg ggctaagaat ccattttttt cttgtcaaca ttataagaaa gcgacattga 53400
gcataaagac attatttgag gacctgctgt actatgtact tagagagata ggcattctat 53460
cttgagttcc ttttttttct cccttcttga aggaagggtta aattgcatct gagatggctc 53520
ttgaaattga tcaggggttc aagctgactt gcatactctt tgggaaagaa tttagaagga 53580
tgtgtatgag gaagttctta tggttaagcc tgtttcctga cttgaataga tgaatcaaat 53640
attttttact attctggaag catcgcattc tggaaagaac catactatgt catctcagtc 53700
tacctcactc cattgtaggc acttggaagc tgaagttgtg atttctccaa aattagatag 53760
ctaattttca ttggtgttag aacaaaaagc gctgcctctc tttgaagaca ccagtcctcc 53820
accgtcctcc tctgcaaggc cgttttcccc cccctttttt ttttttttga gacagagttt 53880
tgctcttggt gcttaggcta cagtacagtg gcacaatctc ggctcactgc aacctccgcc 53940
tcctgggttc aagcgattct cctgtcttag cctccagagt agattacagg caccaccac 54000
cacaccggc taatttttat tattagtagt agtagtagta gtagagatgg ggtttacca 54060
tgttggccag aatggtctcg aactcctgac ctccaggaat cctccacct tggcctcca 54120
aagttctggg attacaggca tgagccactg tgtccagcca atttttctgt atttttaa 54180
gaagatgtga gcagccta atgtaagatcac aacatgtgat tcaatacagc cgtggcttgg 54240
tgttgacatg ttattaccag ttgagcta atcatgtaact cagcatttta tgctttacta 54300
agattaaaat gatgtgataa cattaaattt tgaattacag ttgatgtttt ttatttaaaa 54360
aacatttttc ttagttaaat aatacatgat ggtttaaaaa tcaaataattc agtgcaattc 54420
ttctaaaatc tctgcaagtg tgggggtcat ttaattgctg agcctcccag cctattagct 54480
ttccattctg agctttcaag agatggtggc agctggcaag gcagttttgt ctgggaaagc 54540
cattgttaac agagcagaat tggggatgga gcagccatag cccaccacc agagtaggca 54600
caaatcagac ctgaacgtta tcacaaagtc caagttggct cagacatttg tgttaaatca 54660
taataaatat tttagagaac ttggttgcaa atttacattt gatctcagtc agtcctcttc 54720

ccctatctct acaagcttac aaaccgcatg ggtgtgtggg ggtcttattt aatattgcga 54780
acagctgggt cctgtatctg aagttcttgc cctggagcct ggggtgttgt tgtagtctg 54840
caccatctgc cttgggtgat aaggcatttt ggaggccact gatttttaggc agcagtgttg 54900
ttaggatacg gaaacagcag gatgtttgtg gattgagcct tttcagctga atcttctggc 54960
cagttctttc tggctgtgtg aagttgtgtc gactacagag caggatgctc atgttgccctg 55020
ctgggctctg ttaggggtggc cagacgtgct tgtagcagcc ttactgccag aggaacgtac 55080
gttggtcatcc agagtccagt gctgccgcca gttgcagtgc agcaaggcta gccccaaacc 55140
tgatttgctg caaggattag ctcaactcta gtgacattta ttgtgttttc tcatagccca 55200
aatcacagcc aaaaaaaaaa aaaaaaaaaat ctagggttga cattttttaa aattttttta 55260
aaaaacattt ttcttggtta aataatacat gatggtttaa aaatcaaata ttcattgcag 55320
ttctaaaatc tctgcaagtg tgggggtcat ttaattgctg agcctccag cctcttagct 55380
aaaaaatcta gggttgacat ttttaaaaat gtattcaaca gactacgagg gaaaagatta 55440
aagatgggtg atggaaaacc ataaaagctg agagggaaggc agcactgggc ttagagtcac 55500
ttggcttccc tctagctagt aaataaccag caccaaata cctgatcctc ctgaacttca 55560
gtttctgtgg ccatgaaata agaggttggg tccaggaatc aatgtaaatt gtcaatttaa 55620
catttccctt tattgatatt actccccct gggcttgata atttagttat aattcttcat 55680
gcagcttttag gttgagtaag tttgggtggg aacagtagct ctcttcatat atttgagaga 55740
tgtcatttga aaggggtaga tttattcagt ttaactcaa gaagcagaaa tgggacccat 55800
ggtagaagct accaaatgga ggtttggctc taaataagaa aacgatcttt ggagtgcctc 55860
tcctagttta gatgaaaaaa attgcatcaa gttgtaacca tgctagtcat tgggaatttt 55920
attaacaaca cgtagctcct gtccctgggga ggctcatagt ttgatagggg taagatggaa 55980
agaattgggc agatgtggat tatgtcttag cagtagagcc aacagagtat gttgggggtg 56040
aaggggtaag agaaatcaca tacctcctag gtttttagca ttttccaaaa tgaggaaaat 56100
gggtagaggc atggacagtg acttatattt agacgcgtta agccagttgt aactgcttga 56160
cgtctcagcg ggataacaag taggcagcca tgttgtgtaa tggaaattcc atagctgtag 56220
cctttactaa tgcgctctgg aatggctctat tccagcctct gaaaatgatt tgctgaacaa 56280
gcgcctgaag ctcgattatg aagaaattac tcctgtctt aaagaagtaa ctacagtgtg 56340
ggaaaagatg cttagcactc caggaagatc aaaaattaag tttgacatgg aaaaaatgca 56400
ctcggtgtt gggcaaggta agcttcattg ggaagcatct agtcaacctc acccctcatt 56460
ggtgattggg gagaagtgtg gaattaaaaa aaagtcaagt ctaatttttag tggccatctc 56520
ccttcttttc atcacatctt aatctatttc catatacctt acttaataga catgagtttc 56580
accacctttc atgattcctc ttaattaaaa ttcccagaag gccgggaaat aggaagaaga 56640
cagaaaaacc caagggtttt gttgcctata aactagataa tgatttgatg atatactttg 56700
aattaaatta taaactagaa actaattgta tggcttgtct ctgggtactc tagggagaca 56760
acatagtgtg gggagcacag acttcagaca ggtggtcttg ggcttaaata tcaggcctgc 56820
cacttacttt gcagtgtgat cttagacaaa tgccctaccc tctctgagct ccagtttcta 56880
caagtgtgag atgtgggtgc tgacagtggg tgttgtgagg agcacacagc atgtgtctgc 56940
tatactgtaa ggccttagag agcgggcagg attcactgtt ttttcagtga gatctgccag 57000
cccaaactgt tactgggtcca agaagagata agtacagaac ttgaaactaa gcttttgtaa 57060
atgtttccag caatgtgaca cagtgtacct aattaaaaat gtggacttat attttgtcca 57120
tctgtttttt tttaaatttt gtttttctac taatttattt ttactgtatc gtataaaaaat 57180
atcagcctgt agtagattgg aaaattttta aaaagaaaaa aaattgatgc ttcacagata 57240

gtttgagaac cgctatTTTTg aagcttacct tcagtcatta ttagtgttct agtcaaacaa 57300
tgatttcttt aaaaatatat gttaatgtct tctggcaaga gtaaaagcct gagtctaatac 57360
tgattctatg ctactgagtt ctgggtgagc tcatcatgaa taaccagggtg ttctgaataa 57420
gggtttcaag tatgtataga atgggttttt cctgagttta tcagttgtgc agtgggaaaa 57480
cgttgtatat gcactttttc ttttttgaga tgtagtttca ctcttggtgc ccaggctgga 57540
gtgcaatggc gcgatctcag ctcaactgcaa cccctgcctc ccaggtttaa gctattctcc 57600
tgcctcagcc tectgactag ctgggattac aggcgcgccg caccatgcct ggctaatttt 57660
ttgtgggttt tttttttttt ttttaagaca gagtcttgct ctgtcgccca ggctggaatg 57720
cagtggcgtg atctcagctc actgcaagct ctgcctcccg ggttcacacc attctcctgc 57780
ctcagcctcc tgagtagctg ggactacagg tgcgcgccac catgcccggg taaatttttt 57840
ttgtattttt agtagagatg gggtttact atgttagcca ggatgggtct gatctcctga 57900
cctcatgac caccacctt ggctcccaa agtgctggga ttacagggtg gagccaccgt 57960
gcccggccaa ttttttgtgt ttttactaga gacgggtttt cactgtgttg gcaaggctgg 58020
tcttgaactc tggacctcag gtgatctgcc tgctcggcc tcccaaagtg ctgggattac 58080
agatgtgagc cactgcaccc ggctgcata tgcatttttc atctctagga gcataaatgg 58140
aacaagcag tgttttttac tatagttttt taggcatttt taaccttttc tgaattttga 58200
catcaatttt agtaatcatg ggaagtattt gtttggtacg cttttccct ttctatggat 58260
aaggaaactt gggcttagag cagttgaata gtggcttagg gccacagagc tgggttcaca 58320
ccaccgtact gcactgcctc ctggtgaaca ggatctccag gtgcttatct cagaacacgt 58380
atgcagtggg gaagaccgaa gttctggatg gacaccagct ttcagtgtga ctttagcagg 58440
tacctcttt ctgggtctt gcccccttac tgatagaagg agagacttgc actgagtaga 58500
ggatcttgga gctgtcttg agttctaata ttccttgac ctgtactttt tcttgagggt 58560
tacctttaca ccaaagacc ccaaattgct gttttgaaaa gggagaaagc agagaaaaga 58620
atgagtctgt tcttcccca ttcacagtgt cctagatgat caccttcagg tgtctttgct 58680
tctgcgaaag gcaaattgca tgggtctgtg acagctattc caaatatttg agcttcttag 58740
aagcctggca cctggatatt tgtttttcac tgggcatatt ttgtgggggc taatagaaat 58800
actctaggaa tctggaccct gggtagtgaa agttgggcac agatgattga gcattctgta 58860
tactggagtg agctaaggct gacctggaat ttccttatgt gttgcctgac tttgccacat 58920
cactttttac tgcagaagct ctaaccataa agggggcttt gtcagtcagg tggttttaac 58980
acattaagat ttaacaactc caaacaatg agggcggctt attttggtt tcaagaataaa 59040
aatgtgaatc aaaaaatttg agcctaaatt tgaatcatat ctttgacctt tgaagtagag 59100
gccaactcac ctgagagacc ttgtaagaga ggacagttgt gtggattaag aggcccttcc 59160
tcatagtac ataaaagacc ctgaagtgat ggaaataaag gaatttataa aattttccca 59220
gttaaaatta gatgaggggc caggattagg gtatcaattt aggagaagat aacataatcc 59280
tatgacatta tagataattc agtttagtac acatcaaaat gatttctcta aagatatcta 59340
gatagaacct tataagctgg aatgtctttt ttaggaatgg gattgcagag gggctgcctg 59400
ggctgctgac agtaggggac agatgcaaac tctgcttgcc tttgaccgg caatgccatt 59460
tataaaaact tactctagaa actaatcagc caaaaatgta ctgcagtaag aatgcttatt 59520
gtgacattgt ttaatagtca aaacaaaaca aaaaacccaa catgtgacta tcccatgtca 59580
tattccttga aaatgacacc ataagtagat ctgtatttac tgacgtaaaa gatgtctaag 59640
ttgttaaag aaaagagtac agcatggctt cctgtactgt tgatatttcc atgtgcgtat 59700
acatggaaaa acaccacga tgcagatgtc caggttatag acaggatgac catagggccc 59760

aacctggcat agccctgggt tatgactcct gtcctggcaa aattattaat agccttcctc 59820
ctcttcactg tcaaaagctt cctgctttgg atggtaaata tatgcttatt ctagttatgg 59880
gtggtttttc actttcttct ttatacctct tgcatttcag aggttttttt gcaccacttt 59940
taaacagtga gtgtatatta ttttttaagt gagtaagaag ctatttacat gggggatgga 60000
ggaatggcct cctgccctcc cagaccctgc ctgcaagccg taggtgggct ccactgccag 60060
gtttctcttg ggtaggagt gaaggcagca ccatggtggg gaagggcatt ccaggccatt 60120
cttagcaaaa acattgggtc caacctgcat gatcctgtgc tttaaatcac agaattctaa 60180
cttactcctg aataccacaa tatctggtac tgtccagtga cacagccaat attcttttct 60240
ttcaaaaaat aaaggctctga taagacaatg ggaatgattt agtaatagga aattggacat 60300
ttcataactt gggaaaattt cccagtttga gaaaaagtat tttgtgaaaa aaagccccac 60360
tataaatcac ttatcatgct gactgttttc tagcccatat ttacttctca tcagcatttg 60420
aagtatttgt ggggaggggtg tgcgtgtgtg tgtatgtacc caggatatat ctatgagctg 60480
gaatagcaga gggagacaag aaatagaata atagtagaaa gcagagatca gggatatatt 60540
gcttcctgtt gctaccataa caagttacta caaaaatagt gactaaagca acagaaattc 60600
ttctctcagt gttctggagg ccatagctcc aaaaccatgc cgtagcttg tctgtacttg 60660
gcctcttcca gcttctgggtg tctgtcagct tcctagactt gtggtcacgg cactccagcc 60720
tctgcctcct tggtcacatt gattccccct ctcctctcct cctctgtatg tctattataa 60780
gaatgcttgt cactggatat agggcccatc tggataatcc aggatgctct cctcctccca 60840
aagtccttac ttaattatat ctgcaaagaa ggtaacattc acaagctcca gggattagga 60900
agtgaacaca tctcttttga ggggacacca ttcaactcac tctacagggt cattatatta 60960
atgctgagat aaaattacag aaggatatagg atgtggctcat ggtttacagg ggcctgtat 61020
ttcttctaca ggccaactta aaaaaaatga tacgtgaaag ggaaagaaga aagtacttac 61080
tacacagtaa gtattttcaa gaggtggccc agtgagactt ttgaatctgt taataaaatg 61140
attactattt ggttcaaatc cacagatggg tatttttatca ttaattgcaa gataggaaca 61200
caaaatattt tttctctagt cccatttga gtagcagcct tgtttgacat ttctgacatg 61260
gaggacacca agagaaaatg gcagtcagca tccctgggct gtcactcacc ggcctaataga 61320
cctagggcaa gggacctgtt ctactgcct ctcttttctt taccatgagg ataactatgt 61380
ttcccttaga gggttatgag tatggtatgg gccaatacac ataactgca tggaatggcg 61440
atgggtgcata gtggcctcgc aatcagtgtc atctgctgct gctacctgcc agagcagaaa 61500
cttttcccaa aggtggccag agacagaaac cagagaaacc atccttctgg acaggctgtc 61560
tgagtggcag ggcagggtac aaagcggcca ctttttttcc cggatggaaa gaaagatcaa 61620
tgcctaactt ggaggcttcc tttctcccaa aagacaagaa agacttggca tcttattctt 61680
cagtcttctt gctctccccc tttccacctt tttggccttg taatagctga gtaatgagct 61740
aaagaatttt ggttcaaaac tgtcacctt taaaattagg tttgccctaa ataacatcct 61800
tgactttaag agaattttct taagtttttag acatttttaa tcaactgtgag tattcaaatt 61860
aatcacatgc aaagcattag ttagaggctc ttggacattt tctgttttta gagctttgtt 61920
ggatgctcac atggcaatgt ctgtgcagtc agttcctacc cagcctctgg gctcttcttg 61980
cagcttatct tgcagaaaaga agcctcatca gaattccaga atctcagcta tgattagctt 62040
actccacctc agctcagaaa catgcatgat tccctggagc taccaaactg ggggcagggt 62100
tcttgccgtc aattttgcct ctcaacaata cccttccagc cttcttgcca gctgctctct 62160
tccacatgca cccttgtgcc tgaggcaaac tgaatcactc tcggttccct ctctcttgta 62220
ctttctctct ccttttccct catccttaag gctcggctca aatgaaggat tctgtggaac 62280

cttgattgct cagttagaaa tgagcaaact gtcgcaagga cagaaaacca aacatcgcat 62340
gttctcactc ataggtggga attgaacaat gagaacactt ggacacagga aggggaacat 62400
caaacactgg ggcctatcgt ggggtggggg gagcggggag ggatagcatt aggagatata 62460
cctaattgta aatgacgagt taatgggtgc agcacaccaa catggcacac ttatacatat 62520
gtaacaaacc tgcacgttgt gcacatgtac cctaaaattt aaagtataat aataataata 62580
ataaaaagaa atgagccagc ttctctttca tctgagctct acttcctttt gattctctct 62640
gctttctgag atcacatctt acatgacaat ttttcatact tggttttatt tccctagaat 62700
gttgtaatt ggcaccaggt tggagctcag gtcgtatact ttattccttg cagagtctga 62760
cagggtcaga acatgataac acatttgaga agtgagaaga agggaggaag gggccaggga 62820
agtgagggga gaataggggg tgggaagtagg ggaagaagca aatagggcaa ggttttagtt 62880
gcctcccttc tgttcttatg ctgttaatta ataatggaac cagtggccag gcatgatggc 62940
tcatccctgt aatcccagga ctttgagggc tgaggcagga gtatcgcttg agcccaggag 63000
tttgagacaa gcctggacag catagtgaga ccctgtctct acaaaaataa aaaaaaatt 63060
agccaggcat ggtggtgggc acctataatt tcagctactt gggaggccga ggtgggagga 63120
tcattggagc ccacaagggt gaggctgcag tgagatgtga ttgtgcctct gactgcagc 63180
tcgggtgaca aagccagact ctgtctcaaa aaaaaaaaaa aaggaacaag aatttgata 63240
aatggaacat gaaacacaat tcatttttat tattaagttg tattctgtgc ataaattatt 63300
tccatgtctt ctctcccttt taaagggtgt ccacgtcatc accgaggtga aatctgaaa 63360
tttctagctg agcaattcca ccttaaacac cagtttcca gcaaacagca gccaaaggat 63420
gtgccatata aagaactctt aaagcagctg acttcccagc agcatgcat tcttattgac 63480
cttggttaagt ctgtgccatc gattggagat gacaatggaa gtttactca catgaaaaat 63540
ctgaagagac tgtccaagtt atgtattgac ctgccttag gtttagcaat caaaatttac 63600
tactgagact tttaatttaa aaagccctag ggtaatcaca aatgtcatct tcaagcatat 63660
aaaaatctct gtattttcac tggggagctt gttaactttg cttggcatgg agggagggtg 63720
ttcattaagg ctgcagtcac aattgtggtt cagtcagta actcaaatat tgataggagg 63780
tttttacagt caaccgaagg aacatcctgg aaaacgtata gatgttcaga accgaggctt 63840
ggtttaatta caggagccac tccctcgttt ttactgctca caaacagaat tcatcagaaa 63900
aattgtagaa agcagtttgt gtgtgtgcct tgaatgattt tattttggaa actgggtggc 63960
acctgtctc ttgaatagtt tttaaaataa gaagatggga acaatataca gtcagccctc 64020
catatctatg ggttctgaat ttggggactc aaccaacctc agatggaaag tatttgggaa 64080
gaaaaatcaa tgaaaactaa acaataatat agattttaaa atatagtaac tatctatgta 64140
gaatttacat tgtattaggt gttataggta atctagagat gatttaagggt gtgtgggagg 64200
atgtggccgg gcacagtggc tcacgcctgt aatctcagca ctttgggagg ccaaggctgg 64260
tgatcatga agtcaggaga tcgagaccat cctggctgac acggtgaaac cctgtctcta 64320
ctaaaaatac aaaaaaatta gccaggcatg gtggtgggag cctatagtc cagctactca 64380
ggaggccgag gcaggagaat ggcgtgaacc caggaggcgg agcttgagc gagccaagat 64440
catgccacta cgctccagcc tgggtgacag aatgagactc tgtctcaaaa aaaaaaaaaa 64500
aaagtgtatg ggaggatgtg tgtaggttat gtgcaaacat agcaccatct tatagaaggg 64560
ccttgagcac cgtggatttt ggtgntctgt ggggactcct gcaacctatc ccccgaggat 64620
gccaagggat gactgtattg gatagatttg cagttgccac tgtgaaggac ttgttgaaact 64680
ggggtgtgat tatgatgcac agagggccct cctgacttgt cagtggccat gcacagggcc 64740
aggtggcaat gactccctgt ttgcctgccg cctatcacc aagctgctgt ctctactggt 64800

ggtagagctgg ctogatgtgg taggagatgg gccctgctgc ttttagagca tgtggccctg 64860
cttcagaat acctgttctg gttgcagctg ctgctgctga aggcctccaca gaacacacag 64920
tgctttgggg cctgcggtg gcccggttct ctgattgttc ctgcagccac gacagaggat 64980
gcagtgtgag ccgcatcagg cagtatgaag tcctttcctc tcaagccacg tagctagcct 65040
taaagggttaa tttcataacc ctttaagggtta tttttttttt ttaatttttt tttttgagac 65100
gggtgtctcgc tctgtcgccc aggcctagagt gcagtgggtg gatctcagct cactgcaagc 65160
tccgtctcct ggggttcacag cattctcctg cctcagctc ccaagtagct gggactacag 65220
gtgcccccca ccatgcctag ctaatttttt gtatttttag tagagacggg gtttcaccgt 65280
gttagccagg atgggtctcaa tctcctgacc tcgtgatccg ccgccttggt cctctcaaag 65340
tgctgggatt acaggcgtga gccaccacac ccggccccac ttaagggttat tcttttagctt 65400
gaacatcatc tctgagaaac tttccctgac tgtggtctcc tctccacct caagactgga 65460
tgagggtgtct tgctaagccc cctgtagcac ccacactct ccccatgggtg cgtatcacat 65520
ttctcatcat caccgttatc tgcttattat catcactgct gctgcctaac ttcaccttg 65580
gccaaatgtt gtgcaaaggg acttaaactc ctttctttaa tccttacaac atgatcaggt 65640
agatgttggt ctgtttctct ttagagttga gaaaatagaa acagacagggt tacgtaactt 65700
gctgaaagtg acacagccga tttgccgcta atcagtgtga cttcggaagc tgcacttttt 65760
tttcaacttt tatttttagat tccaggattg cgtatgcagg tttcttacia aggtgtattg 65820
tgtgatgctg aggattggag tgtgattgaa cttgtcaccc aggaaccaag catggtaccc 65880
aataggtagt ttttcaaccc ttgccttct cctccctct ccacccccca ggagtccctg 65940
gtgtctgttc tcatctttat gtccatgtgt acccagtgtt cagctctcat ttctaagtga 66000
gaacatgtga tgcttggttt ctgtttctga attagtttac ttagggtaat gacctgcagc 66060
tgcatccatg ttgctgcaaa ggacatgatt ttgtcccttt ctatggctgc agagtattgc 66120
atgggtgtcca tatatcacat tttctttatc cggttcactg ttactgggca cctgggttg 66180
ttccatgtct ttgcaattgt gaatagtgt gtgatgaacg tgtgagtaca tgtgtctttt 66240
tggtaggatg atttattttg ttttgagtat atactcagta atgggattgc agggtcgaat 66300
ggtaattcag ctcttagcag aacctgtatt tcttactcca cctccccgc ctgtccttag 66360
tatacagcag tggctcttta ttgccttttt ccttatagg atacagccct ctgaggactg 66420
ggctggggct gtttgccat tataccctcg gcttctagga cagtggctgt gacacagcag 66480
atgctcaaag aatatcttta agattcagag tgtgagacac tgcactagca ccgcatctc 66540
atgggcccctc acaacagccc tgggaagggtg gcctgcaccc tctctaagaa atgaagaaac 66600
tgaggtcaca tgttgaccat ggtcaciaaag tcacctgagg ggaggtgaca ggaactgaac 66660
ccactgtcac tctgtgtttc cctgggaccc tctgagcgca ggaggcccggt gttgctgtgc 66720
agtggcaggc caaggcaatg ctttggtgga gctggggccc atttgggcca ctgacctgag 66780
gaaagcagtt ttgtgaattg gcagttagctg catttgctga catggtgagt tacaggaaat 66840
gccatcatgt tcctatcatg tgaaacaaag tgagaaatag gtacagggtg ggaggctgaa 66900
aggagggaat gcagacagcc ccgctcccca cacttgctcc aaggctgggn aggaggaaacg 66960
ggaagggtgc tccccctctg gattcagtc ccttcttctc ttcattcccc tgcagtatcc 67020
cctcattctt ccacggacac gatcagcccc tgcttcttgt tgctcagatg tcatcacttt 67080
tctgcagagg gaaaagaaga gaccagatca gaacaagggc ctcggcgtgg ctgtgcactc 67140
cgaaggcact gtgtgtgcct gagccccacc acggcctccc ctgcagggtc caggcagcct 67200
tccttgagct ggcagtaggt ctgtgggagc ccggtccact ggcagggtct gctgcattca 67260
agtcctctcc atccctgct ctccccaccc tctccctctg nngccccctt ctctgacagt 67320

gctgaccccc ctctctcttc cccactcttt cccatcctcg cctggcctcc ggtttgatg 67380
ctgtccacac acttccccgag ggctgagag gacctccgtg tgaggcaatg catttcccag 67440
gtcacctctg tgtctctctc caggcttttt ccagggactc cccggggtca gtctctctc 67500
cccactggaa cgggggaaact gggattggcc tagaccggc agtggagtcc cagggtgccct 67560
gcctgcccgg ctgactccgc ccaggggaggc ctccacaga agctcctcca gactccacct 67620
gttacctccc ccaactcctct cacccaaggc tgtgctgtgg ccaagtcagt tgtttagtct 67680
acactttctg tttagtctac accatggcta cctcaaggcc cagtgaagggt gtgtagtata 67740
aagcaaaatc aaatccatat ttcagttttc cttaaaaagt gaccttcata ttctggccag 67800
aagaacagaa tgggtgggtg gatataatgt gagttttcat gggttttgt tttcctgcct 67860
cttggtatac tttctgaaat tggcttttag tctaaacagg tttttttttt tttttttttt 67920
tttttnggca atgtgttttc ctccaaagag taagaataat aggcctcatg gctgggtcgt 67980
gttctacagt ttgtgaatat tttctcaacc tttgtcaaat ttcacttta cacatcctgt 68040
gtgaaattgg gcacgtgccg ttattttcaa cttagagggg atgaatgagc ccttaagagc 68100
ttgagttctc tgcccacatt gcgagttact cagtgccaga aggagttctg gaaccagggt 68160
ctcctcagtc tccataccac atcccttcta gggcaccatg ttgcttctgt gtttcttggc 68220
tctgcccact ccatgccagc acaactctcc ccacccctgc tttgggtggaa tcatgttct 68280
ttggggtaga tcacaccagc cagaggcaac tgctctcagc ttagcagatg gtactcatca 68340
cattattctt gaagccttgg gtcaggagcc tgccccaccc atctgcatcc atttgtccag 68400
ccctcagaca attgccactg tttcatgtc tattctttga ctctctatcc tgggtagaca 68460
acatggactg ccagcatcc tgtcttctgt ctggggctcc cactgtcgtc ctgaccacgc 68520
tgggggctgc cagtgcact gggaaactcc cgagggaccc ccttcagggt tcacatcatc 68580
tgctccctcc ctagcatccc agcctagaac actttccagc catcagctgc attccccagt 68640
gaggcgtaga gcctctccca tgataggagg gcttcagccg aaagaacact tcaacaggcc 68700
cagaaaccca ggagcaccat tagatcagaa agcagaagca agaatgcac taatctcccc 68760
cacatcaatt gctatagttt tattaatctg catattatag gtcagtaagg ggatggcaca 68820
gtttataatc cctgcaagag tctgatgatc ttttggtgac cagaagtgcc attttttgat 68880
gggcttctag agatcctcca tcagggatac cagacatgtt tggcatgcct gtgctgccgc 68940
gagacgctaa gcgtgtgtcc agactacacg tgtgggtcat gggtcagca gcagagctgt 69000
catattgatt gtttgcttct actaaatgta taaagcctgc ctgggtgtcca gaagaaaaga 69060
aactataatc caatttttta gaatccataa aaggtaagaa gtaggagaac atttagaatc 69120
cacaaaagat gagaagtagg agaacrgttg gatttttttag aatccataaa agatgagaag 69180
waggagaacc tccaaaagga aggaatcagc tgagagtatt gaagatgacc aagtacaaac 69240
aggcagaggg gagcgcttcc ctttctctc tcccaggcgg tgggtgcct cgctcgcca 69300
ggacacacag agcagcatcg tgcrtttga ggggcagggt gagctgctca tcactagcag 69360
gggtgctggc ggggaccaca gtgttctctt ccatctttga gttgaagtcc tgtgtgagaa 69420
atgagaaacc ttcattggcaa aagacagaaa gggacctaga atgtaacatt cagcagctct 69480
gttatctcac gcacctgtct gtccagttgg ggacgttgct gtatggagggt cagttgaaca 69540
atcacagttg aggagcctaa tgaattcttg caccaccagc cacacacatt attctgaaga 69600
gtgagccatt gtctctgatc ttatcaggat cacatcgtgg gatcatattt atttgggtcat 69660
tctgaatata ccctttaagt ccaaagtga atactaaat gtcgttgata aaaggaaaga 69720
ataaagtggg gtatgatttc ctttcacaga ggtctggaat cttcctgcct ttttcaagtc 69780
agtcgggtgt gctggcaa atgtttaataac cagctcctct caccctcag aggaagccct 69840

tggtgttcag	tgtttgcaga	tttccattgt	gcaactagtc	ctcccacacc	ccattttaaa	69900
ctaccactt	gatgtcactg	gtcatggagt	tgggctcaca	gagccagtgg	gagtcaactg	69960
gagcagccac	tggactcatt	caagtgtttc	ccaaaacaat	ctgctcctag	aaggactctc	70020
ccttaatctc	ctaaccctgc	cattcaggat	gattccctgc	actctgggaa	gcacacgttc	70080
tagtgggaag	actgatactg	ggcaactgat	aaccaagtga	cttaaacttc	tgagggttac	70140
aaaggggtgt	tgtatcctca	gtgtctcatt	tcagattctg	ctcagagcta	aatgcaacaa	70200
tgtgagaaga	tgtagtatc	ccagatcttc	atccaggaag	gaatcttaga	gatcattagg	70260
ttgtagggtt	tctcttctgc	agaggagata	gagggtcggg	gtcagattgc	tggtttgcca	70320
gtaccactcc	ctggagaaaa	gagcaaaaga	aagaaacttg	ttagtcaact	gtgcagagcc	70380
accgtgagac	tgaatagctt	tgtgggtggc	cccgtgtttg	ctgcaagaga	cctctggcct	70440
cttgtagcag	ctgccacatg	gtaaacagag	ccgagatata	aggagtctcg	ctgaaaatgc	70500
agtcagatgg	gctctgaata	gaggaaggca	ggacactctt	gagatgggat	ggggtttctc	70560
acagcaccgt	acagggacca	cctgcaagat	ctcttgaggg	gcttgtgaaa	aacacatccc	70620
tgaggtcacc	attcttgacc	tgtctcttat	tgagtttctg	atgcctggga	tgtgcagggt	70680
taacaagccc	ccagatgata	ctaataggat	tcctgcctga	aaattgctgg	gtgaaggctc	70740
ttccccctcc	aagtgataaa	gaagggaaaag	attgatcctg	gaagaacatc	cgtagatga	70800
gcaaaatttt	gtggagcact	tcatgaagag	gaattactag	gtcatttaga	aatatgtttg	70860
aattgtggat	catctttag	gcctttctgg	catatttctc	cacttagatc	cacaagacac	70920
atcgaatgtc	tttttataaa	gggggttttt	aatgcccata	tttgaccctc	tccacttaac	70980
agtcccatcc	tcattttata	tgtgaaggta	atctgcttta	cagaaaaatg	taaaggacct	71040
gcacttctct	gctttgtggg	aagttgtaaa	atgcagttta	aagaggcagg	cctcatatcc	71100
tgatagattt	gtaggaagga	ttgcacagtt	ttaccacagt	tcctcagagt	ttggcagaaa	71160
ttagctttcc	ctgagctggg	gtcttcccca	gctagcatgc	ttctcctatg	gggtgtgtgg	71220
ccttctctcc	tgtctttttg	aggcagagct	tcaatctaga	atctgttcac	aaactgaaca	71280
aatgcaacaa	acagtaaaaca	gtcttttgct	catagttaag	gtgccttgag	ttgggtgtga	71340
ggggctgagt	gtgttctcag	gggtgctctg	cccacggctc	cggccaactg	ctgcaggtgc	71400
gcatcatatg	gggtgtcttt	gtggaatgcc	atcagcacta	gcttagtacc	tcctaaatgg	71460
gagctggagg	gctacagtgc	tcaacactgg	attatacgaa	tgtggattgt	ccaggaaatg	71520
cttttaatcc	ccctcatcca	ctctctaccc	acgtgacctg	cctctccctc	tttacttggg	71580
gtttactcag	gaatgtgggt	gagttgtcgt	gttagcctag	aacagccatt	ccaaaacttt	71640
gatggaagga	tgccattcac	tttgaaaatt	atcgagtagc	ccaaagagct	tctgtttcca	71700
tggataattc	ctatcaatat	ttactatat	accaattaat	taaaactgag	attagtattt	71760
atgtgattcg	tattttattt	tacatagcta	taataaactc	atacatataa	aaaattatta	71820
aaaaatgact	gttttccaaa	ataaaattag	ttagaaatgt	gacattgttt	ctacattgaa	71880
aaaatctctt	taatgtctga	tttaatagaa	tccgctgaat	tttatttgct	tcattcatcc	71940
tgttttgctt	gttatttgaa	gcatataaag	caaaagctga	ccttgacacag	atctatagta	72000
ggaaaagcag	ggggagggcc	tcatggaccc	ctaaaaggat	ctcagcgacc	tccaggggtc	72060
ctcaggctga	ccaaacattg	agaactattg	acctggaaga	atgtaaaata	ggaaaacagt	72120
gtctccccca	atagaatttc	gtgtaaaacg	tggactgtgt	tacaaagtca	gatgggtgca	72180
gttgtcctgc	ttaaccgcta	atcaggagct	gaaggccaga	gactcacagc	tgttccccag	72240
cctggtagtg	aaccagagg	cctgtcttgc	tgtgcagtgg	gacaggaagt	tgcatttggg	72300
agtctcatag	aacacactgg	aagatgtgtt	ttagcttggc	caggttcatg	caggacagat	72360

tttctgcata	aagaaaatca	atgacagttt	ctgaaactgc	atcctggaag	ccttgaccag	72420
tttgggtaat	aacaagagat	ttgaaagtgt	ggggtgtaca	ggtgttttgc	tgaatctagg	72480
tggtggtggt	gattattatt	atgttgaatt	cagctttcag	ttctacctgc	ttgtgagttc	72540
caaactttgt	gaaaattagt	tgcttggacg	aaacttttct	ttgcctctgg	aaggctgtca	72600
gaaagcgaga	tttcccagct	tatgtgcagt	gttatagtta	atagagtaat	ggctctgcaa	72660
agttgttcct	ttactttaaa	tgtaatttat	tttgcatttg	tgctacagaa	cggtcataag	72720
tgtgcccttt	tgctctcttg	tttggaaact	gggtttttat	aatgtgtgtg	gtctatccga	72780
agattattgc	ccattattga	acaccattca	tagcaaccat	ttgcattagg	cattgtacgt	72840
gtactctcca	ctctgcaaac	tatgtgttct	gtcccttttt	aaaaagagga	agctaagggt	72900
cagagaagct	aggtagtcca	ttctgagctt	cacgtgccag	aggccatttt	gtacttactt	72960
caaatgccat	tgaaataaat	gcacatcaga	gaattgttct	tagcataagg	ggcgctacat	73020
gtaacttttt	attagtgaat	tggatgatgt	tcaagggtcg	tggttgatta	gaaaggcgtc	73080
cagaccctgg	ctccaggggac	tatggagcag	aactcgaggc	cagtgcctgt	cgagcgggtc	73140
cccacactcc	atctgtgtga	cctgactgtg	gatggcctgg	ctctgccgtt	agattgccac	73200
ggtgccctcc	tctggttgaa	cctttctcga	gaagtgcctg	ttggaggctt	gagtgcagag	73260
cctgtgagaa	gctctatgtg	gttcctattg	cctgtcagct	tgctgataaa	ggtcattggt	73320
ttggcaaaat	ttggcccaag	gtttgccttc	tcataacata	ccactcggtg	gcaaggctgg	73380
gaggaagggt	gctatagcta	tttctggaag	ctgcttaggg	ggctgcctcc	ccctaaattg	73440
gtacataatt	tgcagggcct	attgcaagat	gaaaatgcag	aaccctttct	tgaaagatta	73500
ttaggaattt	caagacagag	acaacagagc	atgaagcctt	gtgcaagggt	cttctaagca	73560
cagagccagt	gtgaccgcac	agaacacaca	cccgtgaagc	cagctctgcc	cccaccatct	73620
gaccactctt	gagtggccaa	ttagcatagg	tcactcccca	ccctgctagg	cccaccctct	73680
taggaatggt	gtgaggctta	aataagaaat	agccactcta	caagcgggtg	caattagcat	73740
gggctctggt	ttctgtgtga	ggtagtttgc	taacatgaga	gggtatctga	ttagctaaaa	73800
cgataacact	gacagattaa	attcagaata	actaaacctt	ccctgtgttc	ctttatgcca	73860
catgactcct	gcatattctg	ctaccagcac	ctgtttgata	ccagacggag	gggtccattt	73920
gggatgggac	aggagcatca	gcagaaatgc	agaagtgggg	aagtgtctca	tcttcttgga	73980
agctgagctg	gcaagggtaa	tggaatgaaa	gagattgtga	atatttttga	gactatgagg	74040
aaaccagtac	actggtgttg	cccagtagag	aagccacatg	tggtgtttaa	gcacttgaga	74100
tgtggctact	ccaaattgag	gcgtgtgtgc	agtataaaga	acacactgga	tttcaaagac	74160
ttggcatgaa	aaaagaatgc	ctaattgtct	agtattttta	tattgattat	gttgaagtga	74220
tagtattttg	tgtatgttgg	gtgtaacaaa	atatctaatt	aaaattaact	tcacctgtta	74280
ttttctaatt	tgggtgctag	aaactgttac	atcctgcatg	ggggtcacat	tccagttcag	74340
ttgcatgtgc	tgctacccat	tgttctacac	acacacacac	acacacacag	ctgcacacaa	74400
cctagagggg	tcagagaccc	caggagcccc	tgcttctggt	gcccaggcta	agcgctggag	74460
tggaagataa	agctgggagg	gtgggtgaagg	agggtgagtc	acggagctcc	aggctaacag	74520
agtggataat	ttgttctttg	agcactgggg	agctatggat	tgcttactag	cagcaagggt	74580
acttgtgcag	gggtatatctg	ggggagggtt	actgggggaa	gagatagagg	aggcaagaag	74640
tgaatacaga	acgagaaatc	aggacagtgg	ttaggagacc	gtagctttcc	tcttgagtca	74700
agttcagata	acacatctgg	actgatgaaa	ttctttttca	ggaagctgag	gaagagccca	74760
tgaaaatatg	ttctccctg	tgctgagacc	gaataattgc	agtgaacaat	taacgtgtgg	74820
cctagatcca	ccttttgcc	tcgctgatcc	aagcaggttc	ataattcttg	cctgggcccc	74880

agcttggccc tggctgccag ctgcctggct ccagatgttt cttaatcggt tcaagtactt 74940
ctctgctccc tggaaacagg cactcccatt agtcacattc cagaggagga ggaagaggaa 75000
cttgacaagt atcagctaca aaagcctcct gaacaaaaga aatcctttaa gcctatttga 75060
ataacagttt tttgtgaaaa taatcaggat gttgagagct tttttttttt tcttttaaac 75120
tcttttttga aggttaacttt tgtgaaaaga aaacacctgc tgctcctcag gctgtttcaa 75180
aacactgcct atagtttgaa agtacggaga tatgcatgtg gtatgaagca tttgcaggca 75240
taatattgtg agtctgggaa aagcagatcc agagagtgtg tgtagtaagg cgaggccttt 75300
tagctgcatt tagatgatgc tgggattggg gtgggtgcag ggtgcagcag tggggaggaa 75360
gaactgtgtg tgttcctctt gagaataggg gttatgtcta gaggattaac agttttcttt 75420
tttctttttt tttttttttt ttggagtgtg agtttttctc ttgtctccca ggctggagtg 75480
cagtggcatg atctcagctc actgccatct ctgcctccca ggttcaagca attctcctgc 75540
ctcagcctcc cgagtagctg ggattacagg cacctgccac cagcctgac taattttttc 75600
tattcttagt agcagtgagg tttcgccatg ttgggcaggc tgggtctgaa ctctgacct 75660
caggcgatcc tcccgcctag gcctctgaaa gtgctgggat tacaggcatg agccaccaca 75720
cctggccaac agttttcttt tttcgattga agttcagcta tttgcaggac cgaaggtagt 75780
tctgattact ttcacctgta cttccaccaa aaaataaata aaacaacat gagtaattgc 75840
tgatttttaa ttgaaagcat tattccagga ataactggtg gacttcgttt gcagaggaag 75900
tggcaaagac tgattgatat tatgatccag cttctaaaga ttttgctgct taatctgaag 75960
cacattggat ttctggttca ataggctttc tttttttgtt tttattatta caactaatat 76020
gtattctttt cacagggcga acctttccta cacaccata cttctctgcc cagcttggag 76080
caggacagct atcgctttac aacattttga aggcctactc acttctagac caggaagtgg 76140
gatattgcca aggtctcagc tttgtagcag gcattttgct tcttcatatg agtgaggaag 76200
aggcgtttaa aatgctcaag tttctgatgt ttgacatggg gctgcggaaa cagtatcggc 76260
cagacatgat tattttacag gtatagagtg ttccttatgt ctttaataca acaaaatgct 76320
aagaatgttt cttatccctc tccagatgtg cctcaggagc tttttcaccg tcaggaatac 76380
ttgtaaatagc tgtcactgct gataaaggac tctgtgctag gcattattcc aagcgcttca 76440
tctgcacttc cctctaataa caggaagaca ctgttcacg tctcaatttg tagattggaa 76500
aactgagtct ccaagagatt ataaattggg ccagtcaca cagctagcaa gtgtcagagc 76560
tggactggaa acccaggcct ctctgactct agggccttcc ctcttgcccc catcagccat 76620
cagatgatct cagacctacc tcccagcctc tgcattctgt ctctctctgc ctcaccccca 76680
cccttgctat ctcaggttca gctcaaatat cacatcctgg gagaagctca ttctgactac 76740
cctgatgttg tgttccccac ttccaccttg ggcacactgc gtcacgttat cctggctgat 76800
ttcttttgca caacacagcc actgccagaa atgatcttgt ttccataatc atctccctgt 76860
ctattttctg atttttcata gcctgtgaac tttaggagag ggaagggatc ttacgggtct 76920
tggagccgag ttcttagttt ctgaaacagt gcgtgggttg aagtaggcac ccataagta 76980
tttggtgaat gaacaattct gtcagagaaa accaaacaca gtagcgtatt gcaaatacca 77040
cgtgctgctc ttgctgcctg tcagagggaa aactctggat cctgcttcag gaatattcct 77100
aaatgttgca gcacatgttg atatgttcat ttactaccag taagatacta tgccttcaga 77160
gctctagaga gtatcctggg agggaaataca ttagagccaa ggacttgctt tgagagcacc 77220
aaattatgtg attcaaaatc ttttcacctt gacctgtgaa catggaccac gtgaatgcaa 77280
atatcataga aggaactcat tcaactgaaag attttgacca cataacactt tccacatgta 77340
ctgtgaggtt cttcctacat tccctttatt aactttaaag acagtgggtca ccaggcagtg 77400

gaatttttga	gttttctata	atztatgtaa	cacacaactc	ttttggggtg	gtgccttttg	77460
ttgattagac	agtcttcgat	atgggagagc	cacagctggg	gcttatggga	ttatattatc	77520
tgagcctctg	aaaacggttt	tgttttcttt	ctctcagttt	agataggaca	tatccaactt	77580
gggtggatctt	agcggattct	gaccctctgt	aggttgttgt	ttcttttaggc	tcaggccgtg	77640
gcactgctca	gatctgggct	ggctctcggg	cctctgtgag	cctgtaactc	ttgggtggcac	77700
tactaggaac	tggcatgaga	tttctgccag	aatcatgtca	ttctgtgaag	ttggagttcc	77760
acttttagttg	gaaaaagttt	ttatttcctc	ttaagatgca	cacttgtctt	cttgttttta	77820
cttgccaggt	atctggatat	tccatatatt	atacaccaaa	agaaattatg	cttctcctgc	77880
ctattgagta	atttcagggg	tccagagggg	acttgctgag	tgaacatgta	caatggattc	77940
ctatggaatc	ataagatgcc	cctaattcag	tcttagtaaa	gagactggct	tcttatttct	78000
aattcctcca	ggcttgagtt	gtgcaaagag	tatgtatttg	taagagaatt	tatgaaatgt	78060
ttgcacaaga	cagattttta	gatcttctta	gtggaggaat	acaagggaac	aataaaaagg	78120
aagtggcagt	agaagaccca	gcgttagcgt	cctgggccta	caccagcca	gtgcctggca	78180
ccagcaggca	cttgggaagc	acttgttgga	tgaattagta	gctgagctca	gtggatcgca	78240
agccaaatcg	aatgttttaa	gttctagtaa	gtcttctctt	acaccacccc	tgtgagcagt	78300
aggcataact	ttattgctgt	ggcagatccc	taattctcag	cccttggtgg	tgtcttctctg	78360
cagatccaga	tgtaccagct	ctcgagggtg	cttcatgatt	accacagaga	cctctacaat	78420
cacctggagg	agcacgagat	cggccccagc	ctctacgctg	ccccctgggt	cctcaccatg	78480
tttgccctcac	agttcccgtc	gggattcgta	gccagagtct	ttggtgagca	ttagtaaatc	78540
tgtttgccag	aaccagcctt	ctcttattag	aggggaaaca	tttctgtctc	ctccttggtg	78600
attcttattt	ttatacctgt	agctcttacc	agaacagggt	attgtttgat	agtctaagat	78660
tagtcagggg	tgggttttgt	gactttggag	tcctccttaa	cttctgataa	tcacggggct	78720
tcccttagat	gccttcatct	tgtgggatgt	ggatccgata	cgtgtagata	cgatcgctca	78780
ccatgagggt	ctccctagag	cagacatttg	gaggacttgg	ctgaggagcc	acagggtgat	78840
gtttctcatg	aattgccttc	ctcagccact	ctgggttgtg	agtattgact	gatgctgact	78900
gtgggcctct	gggccccttc	tagattccct	tggcatctct	tcctcccctt	tctcttcttg	78960
ccctgccttc	ggctctacac	tttctcccaa	gtcactgtct	tggagaccag	tgtcaggacc	79020
ttgagtaaca	cctccgtgtg	gatggctcgc	tctccccgct	cagccttgac	acttcatgaa	79080
ggcctcttgc	ccctgagccc	acatgtcaca	gccactgcca	ctcccgtgcc	cccgtgttta	79140
accttgggtg	gttcacatgt	aaaacctgcc	tttatattct	tgatttactt	tttgagaaca	79200
ttgtcaaagt	taggtgagtg	ttcatacaca	aagccttcaa	cctgccttca	tatgcaggga	79260
tagggctgtc	cacgtgcgca	tcaggaaccg	agtggaatgt	tgtgagcatg	gtcagttcgg	79320
gcacagtttg	ttttccctac	tgcagaataa	aagtgatatt	tttgacaatt	caggttcttt	79380
tttttattgt	aaaggaggag	gctactaaaa	aaatgatagt	tattatatat	caaagtgttt	79440
taagcatcac	ttgacagctt	aaaaacatgt	gatcttttaa	aaatttgttt	ttatgattag	79500
agagcatctt	aagggaaatg	ttcaaagaca	ttgatactac	ttcagacatg	ctttgggttaa	79560
acatcttaaa	tatccaaatt	ctagaaatcc	taaaatttgc	tttttaatat	aagtgagcat	79620
ttaccttctt	tctctctttt	cctttccccc	caaatactag	atttttatta	ttcactttta	79680
tctacaagaa	ccttttaaga	gtttcccat	ttgctttact	ataagaattc	atattccttc	79740
ttttctgtcc	ctgaaaaaat	aaaatcacta	aattaaaata	gatacaaaaa	gctatctcct	79800
ggttgagcat	atcttttagtg	agagttcatg	aaggtttata	ccatgggttaa	aaaaaaaaaa	79860
agattaacta	aaagcctcaa	aattgtgtgc	ttagtttatt	aacaaaagag	ttacagaaac	79920

taaaatctca	agctctaggc	tttaagcttt	cttgccaata	acttctatgt	ttttgacttc	79980
tctaactctg	gaaattaaaa	gaaaattatt	aatctacctt	ccttacattt	tctccacatt	80040
ttagctatga	ttttcataca	gggtcatgaa	gaggagtga	gatggaaatg	gggaggaggg	80100
agcgctgttt	gttaatgggt	tgtaaacagc	tcaggcatta	aattacttgg	ttagtgaaga	80160
aaattctacc	aaggcaacca	ggctgaccac	agactggagg	gctgaggggt	catcactgag	80220
tcactctctg	cctggggccc	caggcactgg	agctgctgct	tgagaaaagt	tctggggctc	80280
tggaagagaa	atTTTTcctt	cggctcataa	atgggtaaaa	agacgttaac	aaacaagcag	80340
actccacttt	ggaaatgata	gcccttctat	tgagagtaa	tttgaagctc	tctgaagctc	80400
actctaata	cttcataaat	caaagctgca	gcttgtaaag	gtaagatatt	tttctgtaga	80460
ctttgtaggc	agtggtgaga	ctcggagttt	cataaacatt	atgcatagag	atgccagtgt	80520
ctacatcatt	cctggatccc	acagactcct	gctgtgctaa	gtgggtcggt	gtccagctgg	80580
ccaagggctc	ctgggattag	aggcgggaag	tgggatctca	aggccgcact	ggcttgtgat	80640
gtcactttgc	agaactgct	tttttttcca	cagtccatcc	catctttcag	tacttaaaaa	80700
cagaaaagat	aggtttttac	aaaccatttc	tatttttagc	actgatgact	tagagaatgg	80760
tgatggagat	agcttagttt	tatatctcaa	agcctgccat	tcagtcacta	tagtcttttt	80820
tggcctaggg	cccatttcat	tataagcctt	taagtctgga	taaactctaa	aaacatgtag	80880
aactttgttg	actaagtaaa	atatttcagt	ttgcaccacc	ttagctcata	tattagttaa	80940
taggttacct	ggcattaagt	atgtatctgc	tccttggagg	ggcggctgcc	agtgatgtgt	81000
gcgccttaac	cttcacatga	tactcacacc	ttgctgaatg	gcagttcttc	tacctgggtg	81060
cacataatgt	caatagcccc	ttcctgtatt	tttctagctt	gagtacagca	gggccctggg	81120
agagacgctt	gctctgttca	ctttctcatc	acatctacct	ttgggggaaa	aaaaatctaa	81180
aaaacaggac	ctggcttgct	cctgatggag	gaggaggctg	cagtgttcag	cctctgatgt	81240
ttttctatag	gacatgctgc	caaatagatg	agggaggagg	aggagtataa	aaactaaggg	81300
tttggaacaa	aacacagaag	ccacctgcaa	tatagtgaag	gcttcagaga	gacttttagga	81360
tgaaaataga	ctgaaaacaa	gattgtttct	gtggccagga	aaatctccag	ctattcaggt	81420
gacatgatgc	cgcgatgatga	tgagtgtgtc	cagtctgtct	gtgctgttgt	tctgcacagc	81480
actgtcatca	gccttcagcg	tcccttttac	ccgttactca	tagaatgtag	cggagccacg	81540
actggaggac	cgcagccttc	cagaagaaag	ttgagaaggc	tcagccttga	caaagacaaa	81600
ggtggctgga	aaaagagatg	cagtgcacac	tcacatagga	agattgcact	ttgagatcat	81660
gaagtttagt	ttcaaataga	gttccaatac	acagtaacgc	aataagaggg	ttgctgaaat	81720
gtcctcaaag	aaaagcagtt	cctttgtgtt	gttcccagcg	aatacagtgc	aaagtaatag	81780
agtcagctga	atttaagatt	cctatttcct	gccggataaa	acgtcttgcc	tgtttctagg	81840
tggcttgaaa	aaggagagga	gagaaggaag	aggcaggaga	aaagtccac	tgaaaggacg	81900
tgggctacag	tgtagtgagc	taggctactg	cctcactgcg	ctgggagggt	ccaacagttc	81960
actgtctcct	agggaggttt	tcaaatagcag	gacatttgct	cacttttcca	aggagagtta	82020
ttgttttttt	gttttgtttt	gttttgtttt	gttttaaaaa	attccagaat	gtaaatgtat	82080
aagataccgg	aagacaggca	aaataaaaaat	aattgggttg	gggcagtggg	tttataggta	82140
acatttttct	ttactatttt	tttaaaatta	gatgtgattt	aaaaaaattt	ccaaagccaa	82200
aaaacgggtac	aaatgcttta	aaggatgaag	atgttggtccc	caagtgtcat	cagacaaatt	82260
taggaggccc	ttcctcccaa	gcaaagcttc	ctgcagtcct	tccttcaact	ctgaattcaa	82320
gcacatttct	gcattgtgca	cccaaataat	ctcccgattt	aagacccctt	gtgtctcaca	82380
gaagcttctg	gggctgaact	ttctccggcc	ttggagggtt	ggacgctttg	aatgggagga	82440

gtggtggtga gtggagcatc tctggcagca ggcatttggg agtctctggc aggaatcaat 82500
cagcgtagtc tccaaagggtg gcctttctct gacactaact agcccttgca ggggtcatac 82560
ccataacctg catctcatta acatcatctc cttaccagtg cactgaccta gtgagaaaag 82620
gaacaacaag cattcagcga ctctgcggt gctccagggg aagttagaat tgcttggtg 82680
gggcagaggc ccctgggtgat ctggacctgc gtgcccccat ttgccacct tctgccctgc 82740
acaaccagtg cccctgcctt gccagccaga ctgtttttca ggctcctgca cacctccctg 82800
tattgacacc ctatttttct tttattcaga gtattaatcc tgaggctctga cctaggaaat 82860
tttcattggt tcttcaagca gtcaccttct tgtgggcctt ttcttttctc tttgttctcg 82920
taacaccctg ggcataactc taccgaacca gaactccttg gtgtctctgc agcgtgttct 82980
ttgtgttttg ctcatggctt aatctccaga gcctaataca gtgcctgatg tgtattagat 83040
gctcaataga tgctcattaa gttaaagtag aagacacctc tcagcagagt tctcttaagg 83100
tgttgtgaat agcattggga aagaacattt attttttaat tacattaaat acaaacagat 83160
ataataaaaat aaatcatatg cccagtgcct tgtcttaatt ttttaacata tcaataaaga 83220
gactttaaaa cacataacac caccctctcc cctccaaatt tcttttccgg gaaagtctcc 83280
ttttggaatc ataggaagca cttactaagt tgatttattg taaaaaaacc aagatcctaa 83340
taaatctcag aagatctcct gttaacctaa agagaccact gatgtggatt ctgtatttgg 83400
ttgtgctgac aaaagtttcc cagtaattgt ttatttttaat tggcgtagat gtggtactgt 83460
acctaattta aggcacttgt cctctgaga gtagagacca agctatagaa aatcactggt 83520
gttgtaggga aagcctttcc ccaggatccc tgcaaaaaag gtcttgattt ttattctgaa 83580
agatgccctc attttttgtt cagctataaa agttcatata ttgaaaggag gtctaggaag 83640
tctcactgtg taaaccactg aaacttcaaa ttacttttag agttttgttt ctggaaatgt 83700
catttctgtt taaaaataca tctttgttat agtattattt tagatctttt tattttctgt 83760
agtggggaat tatacaggta gactacattt tataaaccag atatttcaga ggaatattct 83820
tcaattggcc tgccttggtg tatgtaacac ttaccctgaa aagctctgat ttcaaagaca 83880
cagttagtct tctagtatat cttcccagcc tcaacaacca gacttaagaa ggaagtgaag 83940
gattcatctt tcccactttc ctgcgccac cctgagccat cagtagttgt gatgtttgtg 84000
gaaagagtgt ggaccctgag ctgggtggga gaagcaggct gatctcagcg ctggcatggc 84060
ttagggctgc acccatctca gctcacatgg ttaattaagg gttttgtggt gggtacagag 84120
gatctcgagg gctatcccag ccagcgggct cctgggtctg tcatccctgc ctgtgctttg 84180
ttcagaaact acagggatct agtttcccat ttgcacagca gcaccagtc tttgcttttc 84240
tgtttcttctg tggcttttaa atgttatcat attaaccatc tagagaggca ccctgcaagg 84300
ttattcctct cacctgcttt tgctttcctt gatttgatga aatttacagc ttctttctct 84360
cttcattat ctttcagcca aaagaaacag agaaaagaaa tactgacact tgctccaat 84420
tatatttcta ctctgatttt taaaattggt tttttcttat attattatc tagttattag 84480
gtaacctgcc tcagtttagt caaccaataa ttagttatcg tsgctctgct ttaaccacag 84540
gacatcagac tctttttttc cccagcagct tcaactctat gaggaagggt agacagggct 84600
ggggttgctg ctcgccgct tgcttggtg ggtgccctcc ctcttattct gcagtctgta 84660
tagaagttgc atccatttgc cagccactct aagaacaaaa tatggccaga actaggaagt 84720
aaccttgaca gagttcttga actcctcaga gggaaaaatg ttctttatct cattatcatg 84780
ttaaaaatca gtaaacttgt atttaacaaw gtacttctgc agttgtacag ctgttgatca 84840
gttttttaag atctttgaat tctattcctt gtttcaaac agaggaaaca gagacacttt 84900
ttcacttact ctatcttaat ttctgatgct ttatctataa aaatctttta gtgtgaccca 84960

taaaaacatg ttttagtata tccttttaaaa ccaggagca ttcctggaaa aatagactaa 85020
taaaaccttt ttccctttcc cagtttaact tttgaagcat gtttgaattt ttttttcaga 85080
gtaaaacata atttttaatg tttatgtact tttatttgca atactgtctt gacaacactg 85140
tctgagatat caggctctta aaaatgaaat aaagtttgca atgtggggct atgtctccca 85200
cactcctgct ctgtgatgtg tggaaaaggc aatggaatgg tattgctga gaaactggtc 85260
tggtttaact ttctgcattt ctgtgttttc tcagatatga tttttcttca gggaacagag 85320
gtcatattta aagtggcttt aagtctgttg ggaagccata agcccttgat tctgcagcat 85380
gaaaacctag aaaccatagt tgactttata aaaagcacgc taccacactt tggcttggtgta 85440
cagatggaaa agaccatcaa tcaggatatga gtcagtccaa accttgcaaa tgcttaagcc 85500
atcctagata tgtagaaact taaatctctc ttgagcagga actgtttcct accactttgt 85560
gttctgaacg gcattctgca tgatgcctgg catggaggag gcatatcaca aacgtgtgga 85620
atgatcgtga gtgtgtgtgt tgtgagcgtc atggtgaaat gccacatgga aacatggtgg 85680
caatgttttag ctgtagaaac cagcacaggt tattagtagt ttcttacatt taagagactt 85740
cagctctagt agcttgttct tctgaaacat atatatattt atgcaatgat gcaatgtagg 85800
gttttgtaca ttgagtgtt tgatttgtgt gtgtgtatgt tgaatggttt taattggaat 85860
tttctccaaa taattctttg ataacaaagt tatgataggg aacatatatt ctatgaattt 85920
gtttcatgat gtgtgtgtgt ggtttttttt taactgaatt cagttcaata tctgtggctt 85980
cattacctct ggtttcagta tacaatagaa ccattatcct ctgaagtgtt agaggctgag 86040
aggtgagttt tactgggaat tacaactaaa ctgatgggtg aatgccctgg gttgggcatg 86100
ggagcagatt ttgatcctgt acagtttaag gaagaacctg ccagtaatgg ctgctgatga 86160
tggaggacta tgctccttag tagagaaagg ggcattcattg gaagtgtcta gaggctgggt 86220
agctacgaaa gcaactggaag ggatttcctc attgagtcac tgcagagtca gcagtcaagc 86280
ctctcataag cctagaatct gctccagata gtcattgtgtc acttaataat agcaatacgt 86340
tttgagaact gcatcattag gtgatttcatt cattgtgcaa acttcataaa gtgtacttaa 86400
acccaagtgg tatagcctac tacacaccta cacatgttgt tcctaggccc taaaaatctg 86460
tatgacatgt gactatactg aatactgttg gtagttgtaa cacaagggtg agtatattgtg 86520
tatctaaaca taactaaaca tagaaaagg acaataaaaa tatggtattc taattcttatg 86580
ggaccacat catatatgca gttcgccatt gacccaaatg ttgtaatgca acacacaact 86640
gtgtaacgaa agcatagagc aatcaggcaa aaacaaatgg tgaaataaag ctatttttga 86700
aaaatcctta ctctggagat ttctgtagtc caaaagggaat ccatgattcc agtggattgc 86760
atccacatgc agtgtttgtg attttcattt gcagccacac cttagggtgtt aagcacagaa 86820
aaagatgcaa atttggcctg caaaagaaag aggtttcata ccagttgtta acttttagatt 86880
tctgtttgca cattgcatat gcccttatga aagaacagtt cttgtctgtt ctgcactcat 86940
ctttaattga gagcctctcc atctcttttc ctccctgga acactcttct tgatgtggat 87000
agctttggct tgttggggcc ttgctttttg ccagggtggg tgggcatcga aatatacgca 87060
tgatcccatt cagtgcaggc cgcactcctg agaggatata aagtgggttca ctgactgacc 87120
cacactcact ctgctgcaaa gtggaaagggt aggggttcaa actcaagtcc ctcccacctc 87180
aaagtgttt agcagctctc ctacactgcc aagagcctct ggagggtcatt taatttagag 87240
ttttcccta ttttaccagg attctaatac tgacttctcc acccttttga ttctttgatt 87300
tctggcattt tcattcatte tttctttcat tcattctctt cttacagctt ttgttgcag 87360
tacttactta catttacagc ttctagggca gacccccgag agccttggtt acctagactg 87420
agggctatat ccactacctg acatgtcacc ttgtcctgt cctcaggcc atcccagctg 87480

acattgttta	cctcctaagt	attgagcctc	agaaaaaaat	cccattgtct	cctattttct	87540
gtaaaaaaca	aaaaataaaa	cgtattgaga	atacttagga	tacatcaggt	gctgtttcag	87600
tgctggaaga	gtggaaatgg	acacagcatg	ggaagaaaat	agctgtgcgt	gtacctggtt	87660
tgtttcaagc	cgctatctgg	ctatttgga	gttgccattc	atttttccac	tgactttttt	87720
tttttttttt	tttttgagac	agagtcttgc	tctgctaccc	aggctggaat	gcggtggtgt	87780
gatcttggct	cactgcaacc	tccaccttct	gggctcaagc	attctcgtgc	ctcagcttcc	87840
tgagtacctg	ggactatagg	catgtgccac	cacgctcagc	taatttttgt	attttttagta	87900
gaggtggggg	ttcgccatgt	tggccaggct	ggctctgaac	tcctggcctc	aagtgatcca	87960
cctgcctcat	cctcccaaag	tgctgggatt	acatgcatga	gccactgtgc	ccgacctcca	88020
ctgacttaat	aactctagga	cataggtatt	ataattccta	tttttataga	tgaagctgag	88080
cagagagtaa	catgcccggc	cccctgtaga	aaggcagggt	ctgtgggagc	cagggtgtgt	88140
aggttggagc	tgaggtgttt	gagtcagct	ggacttaaaa	gatgacctaa	gatcggtgtg	88200
cagacatttt	ccaaagaagg	ccaggcagta	tatattttag	gcttcacggg	tcataacgtc	88260
acaactactc	acctctgcga	cagctactca	cctcagcctt	tgcagcacia	acaatccatg	88320
acagcatgta	aggggatgat	gggtcatgtt	gcaaaaaact	ttattttaaa	aactgtgcag	88380
cgtgatgggc	ttggtccaca	ggtggtggtt	tgctgatccc	tgaactaaag	gatcacagca	88440
tgtgagaagg	tacaggaatg	agagcagaga	gcaattctca	gaacctgagg	tggtcaacat	88500
tttgggggta	tcgggagcac	aaaaatttgc	atttagggcc	cgggttttta	tcagtgggtcc	88560
ttatagaaag	tagatctacc	cgcatctctc	ccctctttcc	ctctggtgtt	tctatctgaa	88620
cttgacatct	gagtgttctc	tgtcaggctt	tctgctttcc	cactgcccc	tcccatccag	88680
aggggtgctgt	agtcttaagt	tctacacaga	aagcagacat	ggggtccagc	atgattcctc	88740
tgcagcctta	gagatcccc	aggcccaa	cttgggggtc	ttcagagtag	atgtaatgga	88800
tgcttccatt	gcagatgggt	gagcatactg	ccctgtgtac	agatgggggt	gggcaggaag	88860
tgagggcatc	tgggtgggtcc	tctgtgatag	ctgtgggtcc	atttcattaa	atgccctccc	88920
tgtatagata	gtctcagcct	gagcaactgg	gacacagctg	gatcctgggt	ggaagaggtg	88980
gccaaaggaca	gggctctgga	gctctggctc	aatgtggaca	ggctgaaagc	agccagagag	89040
ggcaattcca	aaggtgacag	agcccaggac	agaagacca	gggtgtctga	agcctttgtg	89100
gcagtgtact	taccaaagga	cctgtccatt	gaataaacca	tctacatctt	ctgacccaag	89160
agaatggaa	actttgggaa	taattagtaa	caaaggaaag	gaggtcagtg	ttgtattctt	89220
gtcacagtgg	gtgctctgtg	gctgtgaagt	ctcagctcag	tttaaggaag	aaaaaaagga	89280
aggggtggctg	tgggaggcag	ggcaaatagc	ctattttcag	catcctttag	gctccactca	89340
gagcatggcc	tcagcccagc	attgtcatca	tatcatgtca	gagcttggtta	gaaactcagc	89400
atctggggcc	acacccgaga	cctgctgaat	cagaatctgc	atttcagtga	gatcaccagg	89460
ggattcacat	acacacaaac	agctgagaaa	ccctgctgtg	ggcaactctg	ttagaaacac	89520
aatgaacaaa	ggagccccctg	ttccagttga	gcttgtagg	tagaaaccag	ggttctctgta	89580
ttcagaagac	acacctcaaa	tcaggggcaa	aggtgcctct	tctgcctgtg	ggggagccgt	89640
cacttcttgg	gcagtttgca	ccgtggaaaa	ggagtagttt	tgtacgagga	caactggtgc	89700
cataccagga	gggtggggcg	tggcggggag	aagtggttta	ccactggcgt	tgttgaaaat	89760
tgctcacatg	cagtggtaat	aacaagcaga	gggactttta	gtgggtttga	tgttttttgt	89820
aattcactac	agatagtgtg	tgcccccttg	ttgctgatac	caggccgact	gttcccactc	89880
tccagccctt	ggtatgacaa	tgggaccagc	agattggagg	gcaggggggt	aggaaggcgg	89940
aagctctgtg	gcgagttctg	caaaccrtca	gggttcatga	ctttattaat	cagtgtccat	90000

ggactgtgaa gagaaatgct gagtctacaa tagcaaata gccaagaaca taaacagaca 90060
attcaccgaa gaggagatat ctagtaaaca aatatctggg aaagtatttg gcttcatgtg 90120
taattttaaac ttatgtaaca tataatgctt tactctacta gataatagaa agacatttct 90180
tgatgccagt acccagcacc aagggatatac tgtatgcaga acattagcat gttgctgatg 90240
gcagtgcaca ttgattagt gctgttggga gacaatttgg cgaaacatat cccaagccag 90300
taaaatattc ataccctttg actcagtcac cccgtttctt ggaatgtatc ctcaggaaat 90360
aatccaaaat atgaggggaag ccatatgtat aaggatattc tcctagactt gtcacttata 90420
ataacagaaa cttggaacta gatgtctaac acttgatgac tggattaata tgatgatggg 90480
aggttgagct ggtagaatat catgaagcca gttatatata gcgacatgaa aaagctctta 90540
tttgatacaa tgttaagtaa aggaaaagt ggtataggaaa ttttatgttg gttatgttta 90600
gaactagaaa aacatgcttt taggaaatag gaaatatagc tagatataaa agttgtattt 90660
gggtgattttt cttttatttt tcaagcttcc aataatgtag ctctattgct tcagtaactt 90720
aaaatagttt tatcttttcg gcaaaacatc gaaagtatgg aaatagtcac tcctactttg 90780
gcaaacagaa gagaaatttt cttcagtacc aaaattctgg aacttgactg aaaactatga 90840
agaacctaag agccaggatg acaggaaggc tctagatccc cagtaattac aactctagtg 90900
gaattgctct gagatggggc agcaagaaag aagatgagag ccagtccccc ttgcagaggg 90960
gccaggtacc ttgcagcttt gtgtagtgc cagtgtcag ggaacggctt aggcaagacc 91020
ctgggggagg tgggcactgc acttgtccag cctcaggagt gactcagacc agaaatgaaa 91080
acaccttaaa gtgtatatat cttgttttcc tatcaacacc tagtttttaa tattcgtctg 91140
ttttattcat ctgagacaac ataccaaagg attgggtttt taatgttagg ccttcctgct 91200
ctttctctgg ctgagaactg ctccctggcag tggatcactt gtgctgtcta agtgtgcaag 91260
gacaggcgcc cctcccaatt cttttctttc cccaagtaat tagcccaagg gctgaagccc 91320
tcgtccagt accagggttc tcctttgacc accagcctca tattgccatg gtttggggta 91380
aattcagggg cataactgca gaatgaaggg cctaggagtc ttggcagtca ggagatcatc 91440
aggcaattaa gcagagatga ttgtgacca ggggtggttc tagggattaa tggaggcctg 91500
gaagagttta tggctttggg tactgctgag agccattaac ttaacacaga acatcaatcc 91560
gtaggaaaag ccagagggtt tgttcccagg ctttccagg taggagatca cttaaactct 91620
tgtgaaagaa aaaaagtaat agtgtacatg acatttattc agcaccatat ttataattat 91680
acatgagtgc caacaatct cagttttaac atttgtggtt tttactgttc agactattca 91740
gaagttccat gacgtggcat ccattgatga tggttttgct gaggttgaaa tgtgaggggt 91800
ttgtggcagg tgtgtggaag cagatccctt agctaagaga gcgcctgctc aacctgccag 91860
gcgtgtgctg ggaacttcca ggagttgcct cgtttaattc tcacagccat cctgggaggt 91920
aggcgctgtc tgccccaggt gggaggccag gaactgtggc tgagagaggt taagtaccga 91980
gctcaagtca tcatagagt cgccagtgga gccgcaattt taaggctgac tcaaagcctc 92040
tgagccagtt acctcggtg tcaaatggag ataataatcc ctgctgacct cacggctgct 92100
ctgaaggaca aaagagaatc agttcaatcc agtaacaat tctctctctc cctcttactc 92160
ccctcgca caatgcacaca cccccacag atataatgga ttttagtttt taggcatcaa 92220
atgacttttg tgccttatta atattatcca ctgaatcaaa aacagcaagc tgaaaaattc 92280
atctcaagga agagaaaata agattgttgg gaatggtgag aaaggaaaca tgggtttttga 92340
aaattgattc cagggaagat aggctagttt gaatgccagt agggagccat cagaagaagt 92400
agttttacac tgatttttaa caatattgga gttgcttaag gcaatgcaat agagaggcag 92460
tttctgcccc tttaaagcct gacttcactt tctgaatgtg tgttctgac tagcagggtt 92520

tttttttttt	ttcttttaag	atgggtccag	cttgactgca	ttctcagatc	catcagataa	92580
acgttagggc	ttcactgctg	tgctgagagg	ccccagcccc	tggggttctc	tcatagaaac	92640
aactggaaa	aaaggaaatg	ccttgggcag	cagcagcagc	agctgtcttc	tgattctgct	92700
ttccgccctg	ccttccttac	caagagaaa	tacagacacg	gacggcttga	gtcacttagg	92760
cacttaggag	ttgtttttca	cacgtgtggg	gttttcgtca	ccattactat	tgtgggaaa	92820
aagacaactc	aggcatcggt	tcgtattcac	tcactctgtg	gggtgacatg	tgggttttgg	92880
ctcatttctg	catatttgtg	tgcaaaggag	agtttttttag	taaacagtcc	cattacttag	92940
ctgttcttgt	aactctgaaa	acccaactga	actataatta	aactttgact	tgggtgactct	93000
gcaaacaggc	tatgattctt	ttgtttcttt	tctcctttta	acccatagtt	gatgtatcta	93060
acctaacaga	atttttcagag	aaaagaagt	aaataagaac	taaaaataaa	tttttatgtc	93120
tttaaaaatg	agagggtttt	tttttttttt	tggcttttgg	aagggtgagta	tcaaaaacct	93180
gtacttaatg	ttaccttgga	attatttcta	gatgtttctt	atatcctttt	gtcccaagta	93240
aaattattac	cttctcagtg	cgtagttttt	cttattttatt	acttctagta	ccaagtgtag	93300
agctaagcgt	agaggagacg	cttcacaggt	gcgcattgtc	gtgattgcag	acgcctgcct	93360
gtacttgtgg	ggtttttctc	agtttttagta	cgtgatgact	tttctttcta	taacagggtat	93420
ttgaaatgga	catcgctaaa	cagttacaag	cttatgaagt	tgagtaccac	gtccttcaag	93480
aagaacttat	cgattcctct	cctctcagtg	acaaccaaag	aatggataaa	ttagagaaaa	93540
ccaacagcag	cttacgcaaa	cagaaccttg	acctccttga	acagttgcag	gtagagcata	93600
tttataaagc	agcttcctga	atcacaaata	tatggtagtt	cattaactca	ccaaaggcaa	93660
cagcaggctg	ggctttccca	tgaccagagg	acctttccca	cctgatctg	tttatagttg	93720
ggatcaaagg	tatcccggga	gaatgggtcc	tttttattat	ggagcagaca	gattgtcctt	93780
tgctaaggtc	aggcagtcct	agagctttct	gagaggctgt	ttctgcactt	aactctttta	93840
ggggacaggc	ccagagatga	acttggattc	aggatgccgt	ggcctgttag	ctgaatgccca	93900
gccgttgtca	ttactcaaag	agaatctaag	agcttttaac	ttctatgagc	aaaaccagct	93960
aggtccacag	agggatggta	aaggaggaaa	gtaacacaga	aataaatata	acaacaccaga	94020
agagatgata	attcttttgt	agtccttggt	gcatatacaa	agatttgatt	aatgaaggct	94080
tcagttctcc	cctctagaaa	cttccatttc	aacacggata	tactcaggtg	aggacataca	94140
gaagaaagac	cagttgagac	tgtgcacgca	ggaggggtgt	cagagcaagc	actgaggtgc	94200
agcacggaga	ccagagctgg	ccagggtccag	catcaccccc	accccccacat	caccaggga	94260
cactgcccac	aagaacacct	aactgcggag	tgtagctctt	ttgtcaatct	gatggcatga	94320
agcaaccata	tgttctactt	ttttctactt	tttttaaatgt	cacaagtgtg	tagcagtgtc	94380
gtccctgtta	aggagttgtt	ttgaggggtg	ttttaaaagt	tgtttttgag	tggctgtgga	94440
taaaaataca	tattttttgcc	gaaattttta	tgggtgttct	gggctgtcct	gagaataagt	94500
tccattctga	tctaagcctc	tgatttttct	tcatagaaa	atgagctttg	cagacacaag	94560
cttggcagca	aggtgagaaa	ggccagccta	gtgagtcaag	ctatctgaaa	tgcatctctc	94620
ccagcgggca	ttccatccca	gcatacccta	tcagatatgt	gaaagagagg	aaccaagacc	94680
gaatgctatt	cctgcccagc	cctaataacc	actcacattc	tgaaatttaa	cttctttttt	94740
ttccctaaga	tagagatgtc	ctaactgaaa	atatgcctgt	atacaattta	ccctggaagt	94800
ctcagccatc	actcaaggga	agtctccaga	gggtgaagag	cctgtctggc	ctgtaggggt	94860
acacagtgtg	ggtgggtcatt	ttaaatggct	tccaagccaa	tgataggtcc	ctgaaatata	94920
acatgggtga	aacttcta	aaagctcaca	tttgcattga	agtgttttagc	ttgttaagat	94980
aggcagttct	caaataaaa	gtttgtttta	ttgggtaaat	gacctgttag	ttttttgggtg	95040

acagagcata gaaagtaatt tcatgctgct cctgtgctat tgtttttgtg aagacagggg 95100
gctgtgaaaa actgcttagc tacctacatt cctcaataaa ggcacagac agtaattggg 95160
gattacagat gttctccctg gaatggtcgt tctcttgacc aagtagtcct acacttctgg 95220
aaggatcatt cagaactgtg gtctatgcc aaccaccagt agttcctgag tccctgcagt 95280
gctgagtgtg gggggccacc aagttgagta agacactgca gctctcaaag agttggatct 95340
aggattgtat tgtatcgatt tgtgggtgtt ggatatagtt tttccatgat cccctacgaa 95400
aatatgcctc tcatatgtaa gaatcatgcc tctccgtgt acacttttca gacactgaca 95460
aggaaggggtg ttcaatacag tactgaattt tcatatagct tttctggggg ggccaaaata 95520
ccaaaatcaa cccatttcct acctttattc tgtccataaa attgttagaa atatcaaaat 95580
cccatttcat tccctgttaa atacatgtga acgttgtcta gacgtggag agcaaattct 95640
accacctctt ttgttcagca gtacatcaga cgattgcata gacgtgccag atggaaccaa 95700
ataataatgc acatggattt gtcataatcc gtacaagtca ttgacgcca cactgagcca 95760
gggtgctgtg gagacaggac catgtgtgaa agagaagaca tgcttgcttc tataaaagca 95820
tcggtgttat tgaggagacc tgacattaat gcagaatagc aaatgaccat gcaaattaat 95880
tcactactaa ctaagctgca ggttgcacct cggaatgcag aggggcttca aagtgatgag 95940
ggtagggcct gagccaggcc ctgatgatgg gtggattttg aggatcagag agtacagctt 96000
agagagatac cccaagtggg accaccctt gccagtagg ctgacaaact aaggctcttg 96060
gtcccttttt catatttttg gtgttctagt ggccagcca gagctagact tcgagtcag 96120
aattttctgg cacaagtgtt gtcacattca aaaaagtatt ttctttgtt gaaaaatgaa 96180
aaatatatat atatgtgtgt gtgtatatat atatgtgtgt gtgtgtgtat atatgtgtgt 96240
gtgtatatat acgtgtgtgt gtatatgtgt gtgtgtgtgt gtgtgtgtgt gtgtatatat 96300
gtgtatatat ataccatttt tcccaccta aatggagcat ggcaaactg gactggatta 96360
gtgagataga ccaagtcaca gagcactcca ggatgcagct gtgagctggg gaacagggtca 96420
gaaaggcctc agggacatca gcatacatgt tggagtttct gcagtttct tagggaacct 96480
tttaatgtca ctagagctaa cacacttgct acctgggaag caagcctgcc agagcaaatt 96540
agagagacga gggacagttg ctagaagac acacctggaa gttctattta actagcatta 96600
attatgtgct aggtgctgag gatgtagact gagtgagatc ctcatctctc ctctgtaggg 96660
tggaagagag gatattgctt gtctccatgg ctctagtga acagtcagt agaccaggca 96720
cataaacctc ttagcagaac acttggcctt tctaaggact ccatatgtgt tccggggtaa 96780
atgcctgtgt ttcttgacgt agtgatgtct tgttctcta gacatcacta actttacaca 96840
gtagctttag atggcgtgga cgtgaataaa tgcaacttag gttttcttgt tggtttcttt 96900
ttgagtatca ttgtgtttgt aaagaatttc agattagagg attgttacca cgtgggcctt 96960
caggaggaaa ctgttttgag tttttgtcag cccgaaatcg atttgtgcgt ttaagtatat 97020
gtgctcatca aaacaggcca ggctctgctg cagtaacaaa cttacaagtc tccgaggctt 97080
ggcacaacgg aagtctttgt gtcactgacg cccacttcag ctttgtgttg ctgaagcatt 97140
ctttggcct atccgagctg tccctctggg ggtgggtgct gggggtttg gttccctctg 97200
tgttgtgatc caccatctca gcacggctt ccacagcagc catagcagga gaagaaaatg 97260
ctgggggctc tcagggtgct tttaagggcc tggccaccga cctgcaaggg gtgcgagttg 97320
ccttcctgtg tggccagaac tgatgataaa ctgtagactc atccctgctg aaactcggt 97380
ccagagtgt cccaaggctg gacagcgtgt gggcactgga tcccacctgt gttagcactg 97440
gcaattgtat tctcatttct tcttttattc tccaggtggc aaatggtagg atccaaagcc 97500
ttgaggccac cattgagaag ctctgagca gtgagagcaa gctgaagcag gccatgctta 97560

ccttagaact ggagcggtcg gccctgctgc agacgggtgga ggagctgcgg cggcggagcg 97620
cagagcccag cgaccgggag cctgagtgcg cgcagcccga gccacggggc gactgacagc 97680
tctgcaggag agattgcaac accatcccac actgtccagg ccttaactga gagggacaga 97740
agacgctgga aggagagaag gaagcgggaa gtgtgcttct cagggaggaa accggcttgc 97800
cagcaagtag attcttacga actccaactt gcaattcagg gggcatgtcc cagtgttttt 97860
tttgttgttt ttagatacta aatcgtccct tctccagtcc tgattactgt acacagtagc 97920
tttagatggc gtggacgtga ataaatgcaa cttatgtttt cttgttggtt cctttttgag 97980
tgtcactgtg tttgtaaaga gcattcacia tacgggtggaa tttcaaaagc tgggaagagct 98040
cgagatcatg cctcaggcaa aggcgtgggt ccatcgttct tccgagaggg tttgtgtggc 98100
gactacaccc tcagcgtccc tggcaagggt cagttggctc tcgcccattc ttgttatgga 98160
aacctaagat gatcattggg aagatcagtg atcttgggtc attgatccct ggctcagagg 98220
atagcggttt ccatcataaa ccaagatgat gagttcagcc tttatccctc gtggttccac 98280
tagatgtaac ttaaaggagt taacatttga ggactttgtt ctacatcaga ttttactatt 98340
tgaatgttta agatcacttt attgaatttg aagatcatca aattaaataa aatgatttat 98400
ttaatttggg tatcctgatc actgtcaagt gaaatggatc tctctctttg gtatttaagg 98460
aagtttgtct ttaaaaaaaaa aatagagtgt tttcatacat ttttgcttat cccataagta 98520
cagttgatca aagtcatagt aggtaaatgc tttatgggac agctgacacc ttttagaccc 98580
taccagggtat tgctagcatg tgagctgcag ttgtggggtc tgagatatatt ctttgtggta 98640
gtttcatacc catactatag agtcatgtat ttattttttgc ctgttgtgtg atgtaatgca 98700
atcatgttcc tttgagtctc catcccttgg aaatctgact tcttgcagaa ggagtaggca 98760
catcaagata ttcaggggtg cccaagagt ctgggacttt caaaaaaaaa agatcaggct 98820
nnaactgcag tcagatttat gacagctgac agtttttccag aggtcgcaca cagtgactct 98880
cctctctcag gatgacgagg acctgtgcct tcaacaagca aaatgctgct cacggttgtc 98940
ctgcttgcag ccagtcactg tgtaaagcct ctctgatgtg cacttaagag tgggttgcct 99000
tctcaciaag atggggttct gtgcagtcac aggtcacttc cttgacaaca caatcatttc 99060
tgatctttat cactgtaacc acgtcttcta ttccatagga gtttcttttg attctctcag 99120
ttgcgggggg catctcttaa tcttggggta aaaggagaga ttgccatact tagactcact 99180
gtgagtctcc cgggccattt cacgaggaga ccacagtgtc gccaccagtg cctaaacagg 99240
tggctggcat tcgagacttc ctctgttcc ctgggtcaga ggatagcggg ttccatcata 99300
aaccaagatg atgagttcag cttttatccc tcgtgggttc gctagatgta acttatagga 99360
gttaacattt gaggactttg ttctgcatca gatcttacta tttgaatgtt tactgttgga 99420
ttttgggcat cttattactg ttactcaaaa acattgactc tgcatacaaga aagaaacaag 99480
aaagcaataa aacaagaaat aattcatgct cacattttta tgggtggtttt tttttttttt 99540
ttaactttgg atttttgctt ttcagcccag gagtaaagga atgccttatg aacacctgtg 99600
gcctacgtgt ggtcatgacc caaccatcag tgagattatt tgagatattg gtgtctgcat 99660
ccagtgttgt tatctgagtg tttattacgt aagttgtaac acctctacac aggggtgtgag 99720
tttagcactg atgagaccag ctccatcatt gtatgtggca gtgagtcctg ttacgagatt 99780
gggttgggca gaaaggactg ttgacatgag cctgtggatg taggttgga agtctcagcc 99840
tgtgactgac taggcaagga gcggagaggc aactgtgtga ggattctcag agccaaattt 99900
ttaagccatg ttttgggtta tatttcccc aacactcatt tgtgcacttg gtgggtgtcaa 99960

<211> 3983

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> 171..3725

<220>

<221> polyA_signal

<222> 3942..3947

<223> AATAAA

<220>

<221> misc_feature

<222> 36

<223> n=a, g, c or t

<400> 3

```

ccaggccgtc cccaggatgc cccaagcac ctgcgngtcc cggcccggcc ccgggctctg      60
agcgcgccgc ggcacagggt tctgcatatg aagtgtgtaa aatagattgc ttgatccaaa      120
acagaaaaac agtgataact gttttgctga gttcccagac ccttcccaag atg gaa      176
                                         Met Glu
                                         1
cca ata aca ttc aca gca agg aaa cat ctg ctt cct aac gag gtc tcg      224
Pro Ile Thr Phe Thr Ala Arg Lys His Leu Leu Pro Asn Glu Val Ser
      5              10              15
gtg gat ttt ggc ctg cag ctg gtg ggc tcc ctg cct gtg cat tcc ctg      272
Val Asp Phe Gly Leu Gln Leu Val Gly Ser Leu Pro Val His Ser Leu
      20              25              30
acc acc atg ccc atg ctg ccc tgg gtt gtg gct gag gtg cga aga ctc      320
Thr Thr Met Pro Met Leu Pro Trp Val Val Ala Glu Val Arg Arg Leu
      35              40              45              50
agc agg cag tcc acc aga aag gaa cct gta acc aag caa gtc cgg ctt      368
Ser Arg Gln Ser Thr Arg Lys Glu Pro Val Thr Lys Gln Val Arg Leu
      55              60              65
tgc gtt tca ccc tct gga ctg aga tgt gaa cct gag cca ggg aga agt      416
Cys Val Ser Pro Ser Gly Leu Arg Cys Glu Pro Glu Pro Gly Arg Ser
      70              75              80
caa cag tgg gat ccc ctg atc tat tcc agc atc ttt gag tgc aag cct      464
Gln Gln Trp Asp Pro Leu Ile Tyr Ser Ser Ile Phe Glu Cys Lys Pro
      85              90              95

```

cag cgt gtt cac aaa ctg att cac aac agt cat gac cca agt tac ttt 512
Gln Arg Val His Lys Leu Ile His Asn Ser His Asp Pro Ser Tyr Phe
100 105 110

gct tgt ctg att aag gaa gac gct gtc cac cgg cag agt atc tgc tat 560
Ala Cys Leu Ile Lys Glu Asp Ala Val His Arg Gln Ser Ile Cys Tyr
115 120 125 130

gtg ttc aaa gcc gat gat caa aca aaa gtg cct gag atc atc agc tcc 608
Val Phe Lys Ala Asp Asp Gln Thr Lys Val Pro Glu Ile Ile Ser Ser
135 140 145

atc cgt cag gcg ggg aag atc gcc cgg cag gag gag ctg cac tgc ccg 656
Ile Arg Gln Ala Gly Lys Ile Ala Arg Gln Glu Glu Leu His Cys Pro
150 155 160

tcc gag ttc gac gac acg ttt tcc aag aag ttc gag gtg ctc ttc tgc 704
Ser Glu Phe Asp Asp Thr Phe Ser Lys Lys Phe Glu Val Leu Phe Cys
165 170 175

ggc cgc gtg acg gtg gcg cac aag aag gct ccg ccg gcc ctg atc gac 752
Gly Arg Val Thr Val Ala His Lys Lys Ala Pro Pro Ala Leu Ile Asp
180 185 190

gag tgc atc gag aag ttc aat cac gtc agc ggc agc cgg ggg tcc gag 800
Glu Cys Ile Glu Lys Phe Asn His Val Ser Gly Ser Arg Gly Ser Glu
195 200 205 210

agc ccc cgc ccc aac ccg ccc cat gcc gcg ccc aca ggg agc cag gag 848
Ser Pro Arg Pro Asn Pro Pro His Ala Ala Pro Thr Gly Ser Gln Glu
215 220 225

cct gtg cgc agg ccc atg cgc aag tcc ttc tcc cag ccc ggc ctg cgc 896
Pro Val Arg Arg Pro Met Arg Lys Ser Phe Ser Gln Pro Gly Leu Arg
230 235 240

tcg ctg gcc ttt agg aag gag ctg cag gat ggg ggc ctc cga agc agc 944
Ser Leu Ala Phe Arg Lys Glu Leu Gln Asp Gly Gly Leu Arg Ser Ser
245 250 255

ggc ttc ttc agc tcc ttc gag gag agc gac att gag aac cac ctc att 992
Gly Phe Phe Ser Ser Phe Glu Glu Ser Asp Ile Glu Asn His Leu Ile
260 265 270

agc gga cac aat att gtg cag ccc aca gat atc gag gaa aat cga act 1040
Ser Gly His Asn Ile Val Gln Pro Thr Asp Ile Glu Glu Asn Arg Thr
275 280 285 290

atg ctc ttc acg att ggc cag tct gaa gtt tac ctc atc agt cct gac 1088
Met Leu Phe Thr Ile Gly Gln Ser Glu Val Tyr Leu Ile Ser Pro Asp
295 300 305

acc aaa aaa ata gca ttg gag aaa aat ttt aag gag ata tcc ttt tgc 1136
Thr Lys Lys Ile Ala Leu Glu Lys Asn Phe Lys Glu Ile Ser Phe Cys
310 315 320

tct cag ggc atc aga cac gtg gac cac ttt ggg ttt atc tgt cgg gag	1184
Ser Gln Gly Ile Arg His Val Asp His Phe Gly Phe Ile Cys Arg Glu	
325 330 335	
tct tcc gga ggt ggc ggc ttt cat ttt gtc tgt tac gtg ttt cag tgc	1232
Ser Ser Gly Gly Gly Gly Phe His Phe Val Cys Tyr Val Phe Gln Cys	
340 345 350	
aca aat gag gct ctg gtt gat gaa att atg atg acc ctg aaa cag gcc	1280
Thr Asn Glu Ala Leu Val Asp Glu Ile Met Met Thr Leu Lys Gln Ala	
355 360 365 370	
ttc acg gtg gcc gca gtg cag cag aca gct aag gcg cca gcc cag ctg	1328
Phe Thr Val Ala Ala Val Gln Gln Thr Ala Lys Ala Pro Ala Gln Leu	
375 380 385	
tgt gag ggc tgc ccc ctg caa agc ctg cac aag ctc tgt gag agg ata	1376
Cys Glu Gly Cys Pro Leu Gln Ser Leu His Lys Leu Cys Glu Arg Ile	
390 395 400	
gag gga atg aat tct tcc aaa aca aaa cta gaa ctg caa aag cac ctg	1424
Glu Gly Met Asn Ser Ser Lys Thr Lys Leu Glu Leu Gln Lys His Leu	
405 410 415	
acg aca tta acc aat cag gag cag gcg act att ttt gaa gag gtt cag	1472
Thr Thr Leu Thr Asn Gln Glu Gln Ala Thr Ile Phe Glu Glu Val Gln	
420 425 430	
aaa ttg aga ccg aga aat gag cag cga gag aat gaa ttg att att tct	1520
Lys Leu Arg Pro Arg Asn Glu Gln Arg Glu Asn Glu Leu Ile Ile Ser	
435 440 445 450	
ttt ctg aga tgt tta tat gaa gag aaa cag aaa gaa cac atc cat att	1568
Phe Leu Arg Cys Leu Tyr Glu Glu Lys Gln Lys Glu His Ile His Ile	
455 460 465	
ggg gag atg aag cag aca tcg cag atg gca gca gag aat att gga agt	1616
Gly Glu Met Lys Gln Thr Ser Gln Met Ala Ala Glu Asn Ile Gly Ser	
470 475 480	
gaa tta cca ccc agt gcc act cga ttt agg cta gat atg ctg aaa aac	1664
Glu Leu Pro Pro Ser Ala Thr Arg Phe Arg Leu Asp Met Leu Lys Asn	
485 490 495	
aaa gca aag aga tct tta aca gag tct tta gaa agt att ttg tcc cgg	1712
Lys Ala Lys Arg Ser Leu Thr Glu Ser Leu Glu Ser Ile Leu Ser Arg	
500 505 510	
ggt aat aaa gcc aga ggc ctg cag gaa cac tcc atc agt gtg gat ctg	1760
Gly Asn Lys Ala Arg Gly Leu Gln Glu His Ser Ile Ser Val Asp Leu	
515 520 525 530	
gat agc tcc ctg tct agt aca tta agt aac acc agc aaa gag cca tct	1808
Asp Ser Ser Leu Ser Ser Thr Leu Ser Asn Thr Ser Lys Glu Pro Ser	
535 540 545	

gtg tgt gaa aag gag gcc ttg ccc atc tct gag agc tcc ttt aag ctc	1856
Val Cys Glu Lys Glu Ala Leu Pro Ile Ser Glu Ser Ser Phe Lys Leu	
550 555 560	
ctc ggc tcc tcg gag gac ctg tcc agt gac tcg gag agt cat ctc cca	1904
Leu Gly Ser Ser Glu Asp Leu Ser Ser Asp Ser Glu Ser His Leu Pro	
565 570 575	
gaa gag cca gct ccg ctg tcg ccc cag cag gcc ttc agg agg cga gca	1952
Glu Glu Pro Ala Pro Leu Ser Pro Gln Gln Ala Phe Arg Arg Arg Ala	
580 585 590	
aac acc ctg agt cac ttc ccc atc gaa tgc cag gaa cct cca caa cct	2000
Asn Thr Leu Ser His Phe Pro Ile Glu Cys Gln Glu Pro Pro Gln Pro	
595 600 605 610	
gcc cgg ggg tcc ccg ggg gtt tcg caa agg aaa ctt atg agg tat cac	2048
Ala Arg Gly Ser Pro Gly Val Ser Gln Arg Lys Leu Met Arg Tyr His	
615 620 625	
tca gtg agc aca gag acg cct cat gaa cga aag gac ttt gaa tcc aaa	2096
Ser Val Ser Thr Glu Thr Pro His Glu Arg Lys Asp Phe Glu Ser Lys	
630 635 640	
gca aac cat ctt ggt gat tct ggt ggg act cct gtg aag acc cgg agg	2144
Ala Asn His Leu Gly Asp Ser Gly Gly Thr Pro Val Lys Thr Arg Arg	
645 650 655	
cat tcc tgg agg cag cag ata ttc ctc cga gta gcc acc ccg cag aag	2192
His Ser Trp Arg Gln Gln Ile Phe Leu Arg Val Ala Thr Pro Gln Lys	
660 665 670	
gcg tgc gat tct tcc agc aga tat gaa gat tat tca gag ctg gga gag	2240
Ala Cys Asp Ser Ser Ser Arg Tyr Glu Asp Tyr Ser Glu Leu Gly Glu	
675 680 685 690	
ctt ccc cca cga tct cct tta gaa cca gtt tgt gaa gat ggg ccc ttt	2288
Leu Pro Pro Arg Ser Pro Leu Glu Pro Val Cys Glu Asp Gly Pro Phe	
695 700 705	
ggc ccc cca cca gag gaa aag aaa agg aca tct cgt gag ctc cga gag	2336
Gly Pro Pro Pro Glu Glu Lys Lys Arg Thr Ser Arg Glu Leu Arg Glu	
710 715 720	
ctg tgg caa aag gct att ctt caa cag ata ctg ctg ctt aga atg gag	2384
Leu Trp Gln Lys Ala Ile Leu Gln Gln Ile Leu Leu Leu Arg Met Glu	
725 730 735	
aag gaa aat cag aag ctc caa gcc tct gaa aat gat ttg ctg aac aag	2432
Lys Glu Asn Gln Lys Leu Gln Ala Ser Glu Asn Asp Leu Leu Asn Lys	
740 745 750	
cgc ctg aag ctc gat tat gaa gaa att act ccc tgt ctt aaa gaa gta	2480
Arg Leu Lys Leu Asp Tyr Glu Glu Ile Thr Pro Cys Leu Lys Glu Val	
755 760 765 770	

act aca gtg tgg gaa aag atg ctt agc act cca gga aga tca aaa att	2528
Thr Thr Val Trp Glu Lys Met Leu Ser Thr Pro Gly Arg Ser Lys Ile	
775 780 785	
aag ttt gac atg gaa aaa atg cac tcg gct gtt ggg caa ggt gtg cca	2576
Lys Phe Asp Met Glu Lys Met His Ser Ala Val Gly Gln Gly Val Pro	
790 795 800	
cgt cat cac cga ggt gaa atc tgg aaa ttt cta gct gag caa ttc cac	2624
Arg His His Arg Gly Glu Ile Trp Lys Phe Leu Ala Glu Gln Phe His	
805 810 815	
ctt aaa cac cag ttt ccc agc aaa cag cag cca aag gat gtg cca tac	2672
Leu Lys His Gln Phe Pro Ser Lys Gln Gln Pro Lys Asp Val Pro Tyr	
820 825 830	
aaa gaa ctc tta aag cag ctg act tcc cag cag cat gcg att ctt att	2720
Lys Glu Leu Leu Lys Gln Leu Thr Ser Gln Gln His Ala Ile Leu Ile	
835 840 845 850	
gac ctt ggg cga acc ttt cct aca cac cca tac ttc tct gcc cag ctt	2768
Asp Leu Gly Arg Thr Phe Pro Thr His Pro Tyr Phe Ser Ala Gln Leu	
855 860 865	
gga gca gga cag cta tcg ctt tac aac att ttg aag gcc tac tca ctt	2816
Gly Ala Gly Gln Leu Ser Leu Tyr Asn Ile Leu Lys Ala Tyr Ser Leu	
870 875 880	
cta gac cag gaa gtg gga tat tgc caa ggt ctc agc ttt gta gca ggc	2864
Leu Asp Gln Glu Val Gly Tyr Cys Gln Gly Leu Ser Phe Val Ala Gly	
885 890 895	
att ttg ctt ctt cat atg agt gag gaa gag gcg ttt aaa atg ctc aag	2912
Ile Leu Leu Leu His Met Ser Glu Glu Glu Ala Phe Lys Met Leu Lys	
900 905 910	
ttt ctg atg ttt gac atg ggg ctg cgg aaa cag tat cgg cca gac atg	2960
Phe Leu Met Phe Asp Met Gly Leu Arg Lys Gln Tyr Arg Pro Asp Met	
915 920 925 930	
att att tta cag atc cag atg tac cag ctc tcg agg ttg ctt cat gat	3008
Ile Ile Leu Gln Ile Gln Met Tyr Gln Leu Ser Arg Leu Leu His Asp	
935 940 945	
tac cac aga gac ctc tac aat cac ctg gag gag cac gag atc ggc ccc	3056
Tyr His Arg Asp Leu Tyr Asn His Leu Glu Glu His Glu Ile Gly Pro	
950 955 960	
agc ctc tac gct gcc ccc tgg ttc ctc acc atg ttt gcc tca cag ttc	3104
Ser Leu Tyr Ala Ala Pro Trp Phe Leu Thr Met Phe Ala Ser Gln Phe	
965 970 975	
ccg ctg gga ttc gta gcc aga gtc ttt gat atg att ttt ctt cag gga	3152
Pro Leu Gly Phe Val Ala Arg Val Phe Asp Met Ile Phe Leu Gln Gly	
980 985 990	

```

aca gag gtc ata ttt aaa gtg gct tta agt ctg ttg gga agc cat aag      3200
Thr Glu Val Ile Phe Lys Val Ala Leu Ser Leu Leu Gly Ser His Lys
995                      1000                      1005                      1010
ccc ttg att ctg cag cat gaa aac cta gaa acc ata gtt gac ttt ata      3248
Pro Leu Ile Leu Gln His Glu Asn Leu Glu Thr Ile Val Asp Phe Ile
                      1015                      1020                      1025
aaa agc acg cta ccc aac ctt ggc ttg gta cag atg gaa aag acc atc      3296
Lys Ser Thr Leu Pro Asn Leu Gly Leu Val Gln Met Glu Lys Thr Ile
                      1030                      1035                      1040
aat cag gta ttt gaa atg gac atc gct aaa cag tta caa gct tat gaa      3344
Asn Gln Val Phe Glu Met Asp Ile Ala Lys Gln Leu Gln Ala Tyr Glu
                      1045                      1050                      1055
gtt gag tac cac gtc ctt caa gaa gaa ctt atc gat tcc tct cct ctc      3392
Val Glu Tyr His Val Leu Gln Glu Glu Leu Ile Asp Ser Ser Pro Leu
                      1060                      1065                      1070
agt gac aac caa aga atg gat aaa tta gag aaa acc aac agc agc tta      3440
Ser Asp Asn Gln Arg Met Asp Lys Leu Glu Lys Thr Asn Ser Ser Leu
1075                      1080                      1085                      1090
cgc aaa cag aac ctt gac ctc ctt gaa cag ttg cag gtg gca aat ggt      3488
Arg Lys Gln Asn Leu Asp Leu Leu Glu Gln Leu Gln Val Ala Asn Gly
                      1095                      1100                      1105
agg atc caa agc ctt gag gcc acc att gag aag ctc ctg agc agt gag      3536
Arg Ile Gln Ser Leu Glu Ala Thr Ile Glu Lys Leu Leu Ser Ser Glu
                      1110                      1115                      1120
agc aag ctg aag cag gcc atg ctt acc tta gaa ctg gag cgg tcg gcc      3584
Ser Lys Leu Lys Gln Ala Met Leu Thr Leu Glu Leu Glu Arg Ser Ala
                      1125                      1130                      1135
ctg ctg cag acg gtg gag gag ctg cgg cgg cgg agc gca gag ccc agc      3632
Leu Leu Gln Thr Val Glu Glu Leu Arg Arg Arg Ser Ala Glu Pro Ser
                      1140                      1145                      1150
gac cgg gag cct gag tgc acg cag ccc gag ccc acg ggc gac tga      3677
Asp Arg Glu Pro Glu Cys Thr Gln Pro Glu Pro Thr Gly Asp *
1155                      1160                      1165
cagctctgca ggagagattg caacaccatc ccacactgtc caggccttaa ctgagagggga      3737
cagaagacgc tggaaggaga gaaggaagcg ggaagtgtgc ttctcagggg ggaaccggc      3797
ttgccagcaa gtagattctt acgaactcca acttgcaatt cagggggcat gtcccagtgt      3857
tttttttggt gtttttagat actaaatcgt cccttctcca gtccgtgatta ctgtacacag      3917
tagctttaga tggcgtggac gtgaataaat gcaacttatg ttttaaaaaa aaaaaaaaaa      3977
aaaaaa                                                    3983

```

<210> 4

<211> 3988

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> 176..3730

<220>

<221> polyA_signal

<222> 3947..3952

<223> AATAAA

<220>

<221> misc_feature

<222> 1..458

<223> homology with Genset 5' EST in ref : A35235

<400> 4

ataataggca ctgaagacat gttaatggaa ggtggatttg tgattcagaa cctctagact	60
acctgggcga gtctttttaa atgtttctgc atatgaagtg tgtaaaatag attgcttgat	120
ccaaaacaga aaaacagtga taactgtttt gctgagttcc cagacccttc ccaag atg	178
	Met
	1
gaa cca ata aca ttc aca gca agg aaa cat ctg ctt cct aac gag gtc	226
Glu Pro Ile Thr Phe Thr Ala Arg Lys His Leu Leu Pro Asn Glu Val	
5 10 15	
tcg gtg gat ttt ggc ctg cag ctg gtg ggc tcc ctg cct gtg cat tcc	274
Ser Val Asp Phe Gly Leu Gln Leu Val Gly Ser Leu Pro Val His Ser	
20 25 30	
ctg acc acc atg ccc atg ctg ccc tgg gtt gtg gct gag gtg cga aga	322
Leu Thr Thr Met Pro Met Leu Pro Trp Val Val Ala Glu Val Arg Arg	
35 40 45	
ctc agc agg cag tcc acc aga aag gaa cct gta acc aag caa gtc cgg	370
Leu Ser Arg Gln Ser Thr Arg Lys Glu Pro Val Thr Lys Gln Val Arg	
50 55 60 65	
ctt tgc gtt tca ccc tct gga ctg aga tgt gaa cct gag cca ggg aga	418
Leu Cys Val Ser Pro Ser Gly Leu Arg Cys Glu Pro Glu Pro Gly Arg	
70 75 80	
agt caa cag tgg gat ccc ctg atc tat tcc agc atc ttt gag tgc aag	466
Ser Gln Gln Trp Asp Pro Leu Ile Tyr Ser Ser Ile Phe Glu Cys Lys	
85 90 95	
cct cag cgt gtt cac aaa ctg att cac aac agt cat gac cca agt tac	514

Pro	Gln	Arg	Val	His	Lys	Leu	Ile	His	Asn	Ser	His	Asp	Pro	Ser	Tyr	
	100						105					110				
ttt	gct	tgt	ctg	att	aag	gaa	gac	gct	gtc	cac	cgg	cag	agt	atc	tgc	562
Phe	Ala	Cys	Leu	Ile	Lys	Glu	Asp	Ala	Val	His	Arg	Gln	Ser	Ile	Cys	
	115						120					125				
tat	gtg	ttc	aaa	gcc	gat	gat	caa	aca	aaa	gtg	cct	gag	atc	atc	agc	610
Tyr	Val	Phe	Lys	Ala	Asp	Asp	Gln	Thr	Lys	Val	Pro	Glu	Ile	Ile	Ser	
	130						135				140				145	
tcc	atc	cgt	cag	gcg	ggg	aag	atc	gcc	cgg	cag	gag	gag	ctg	cac	tgc	658
Ser	Ile	Arg	Gln	Ala	Gly	Lys	Ile	Ala	Arg	Gln	Glu	Glu	Leu	His	Cys	
					150					155				160		
ccg	tcc	gag	ttc	gac	gac	acg	ttt	tcc	aag	aag	ttc	gag	gtg	ctc	ttc	706
Pro	Ser	Glu	Phe	Asp	Asp	Thr	Phe	Ser	Lys	Lys	Phe	Glu	Val	Leu	Phe	
			165					170				175				
tgc	ggc	cgc	gtg	acg	gtg	gcg	cac	aag	aag	gct	ccg	ccg	gcc	ctg	atc	754
Cys	Gly	Arg	Val	Thr	Val	Ala	His	Lys	Lys	Ala	Pro	Pro	Ala	Leu	Ile	
			180					185				190				
gac	gag	tgc	atc	gag	aag	ttc	aat	cac	gtc	agc	ggc	agc	cgg	ggg	tcc	802
Asp	Glu	Cys	Ile	Glu	Lys	Phe	Asn	His	Val	Ser	Gly	Ser	Arg	Gly	Ser	
		195					200				205					
gag	agc	ccc	cgc	ccc	aac	ccg	ccc	cat	gcc	gcg	ccc	aca	ggg	agc	cag	850
Glu	Ser	Pro	Arg	Pro	Asn	Pro	Pro	His	Ala	Ala	Pro	Thr	Gly	Ser	Gln	
	210					215				220				225		
gag	cct	gtg	cgc	agg	ccc	atg	cgc	aag	tcc	ttc	tcc	cag	ccc	ggc	ctg	898
Glu	Pro	Val	Arg	Arg	Pro	Met	Arg	Lys	Ser	Phe	Ser	Gln	Pro	Gly	Leu	
					230				235			240				
cgc	tcg	ctg	gcc	ttt	agg	aag	gag	ctg	cag	gat	ggg	ggc	ctc	cga	agc	946
Arg	Ser	Leu	Ala	Phe	Arg	Lys	Glu	Leu	Gln	Asp	Gly	Gly	Leu	Arg	Ser	
			245					250				255				
agc	ggc	ttc	ttc	agc	tcc	ttc	gag	gag	agc	gac	att	gag	aac	cac	ctc	994
Ser	Gly	Phe	Phe	Ser	Ser	Phe	Glu	Glu	Ser	Asp	Ile	Glu	Asn	His	Leu	
		260					265				270					
att	agc	gga	cac	aat	att	gtg	cag	ccc	aca	gat	atc	gag	gaa	aat	cga	1042
Ile	Ser	Gly	His	Asn	Ile	Val	Gln	Pro	Thr	Asp	Ile	Glu	Glu	Asn	Arg	
		275					280				285					
act	atg	ctc	ttc	acg	att	ggc	cag	tct	gaa	gtt	tac	ctc	atc	agt	cct	1090
Thr	Met	Leu	Phe	Thr	Ile	Gly	Gln	Ser	Glu	Val	Tyr	Leu	Ile	Ser	Pro	
	290					295				300				305		
gac	acc	aaa	aaa	ata	gca	ttg	gag	aaa	aat	ttt	aag	gag	ata	tcc	ttt	1138
Asp	Thr	Lys	Lys	Ile	Ala	Leu	Glu	Lys	Asn	Phe	Lys	Glu	Ile	Ser	Phe	
					310					315				320		
tgc	tct	cag	ggc	atc	aga	cac	gtg	gac	cac	ttt	ggg	ttt	atc	tgt	cgg	1186

Cys Ser Gln Gly Ile Arg His Val Asp His Phe Gly Phe Ile Cys Arg	
325 330 335	
gag tct tcc gga ggt ggc ggc ttt cat ttt gtc tgt tac gtg ttt cag	1234
Glu Ser Ser Gly Gly Gly Gly Phe His Phe Val Cys Tyr Val Phe Gln	
340 345 350	
tgc aca aat gag gct ctg gtt gat gaa att atg atg acc ctg aaa cag	1282
Cys Thr Asn Glu Ala Leu Val Asp Glu Ile Met Met Thr Leu Lys Gln	
355 360 365	
gcc ttc acg gtg gcc gca gtg cag cag aca gct aag gcg cca gcc cag	1330
Ala Phe Thr Val Ala Ala Val Gln Gln Thr Ala Lys Ala Pro Ala Gln	
370 375 380 385	
ctg tgt gag ggc tgc ccc ctg caa agc ctg cac aag ctc tgt gag agg	1378
Leu Cys Glu Gly Cys Pro Leu Gln Ser Leu His Lys Leu Cys Glu Arg	
390 395 400	
ata gag gga atg aat tct tcc aaa aca aaa cta gaa ctg caa aag cac	1426
Ile Glu Gly Met Asn Ser Ser Lys Thr Lys Leu Glu Leu Gln Lys His	
405 410 415	
ctg acg aca tta acc aat cag gag cag gcg act att ttt gaa gag gtt	1474
Leu Thr Thr Leu Thr Asn Gln Glu Gln Ala Thr Ile Phe Glu Glu Val	
420 425 430	
cag aaa ttg aga ccg aga aat gag cag cga gag aat gaa ttg att att	1522
Gln Lys Leu Arg Pro Arg Asn Glu Gln Arg Glu Asn Glu Leu Ile Ile	
435 440 445	
tct ttt ctg aga tgt tta tat gaa gag aaa cag aaa gaa cac atc cat	1570
Ser Phe Leu Arg Cys Leu Tyr Glu Glu Lys Gln Lys Glu His Ile His	
450 455 460 465	
att ggg gag atg aag cag aca tcg cag atg gca gca gag aat att gga	1618
Ile Gly Glu Met Lys Gln Thr Ser Gln Met Ala Ala Glu Asn Ile Gly	
470 475 480	
agt gaa tta cca ccc agt gcc act cga ttt agg cta gat atg ctg aaa	1666
Ser Glu Leu Pro Pro Ser Ala Thr Arg Phe Arg Leu Asp Met Leu Lys	
485 490 495	
aac aaa gca aag aga tct tta aca gag tct tta gaa agt att ttg tcc	1714
Asn Lys Ala Lys Arg Ser Leu Thr Glu Ser Leu Glu Ser Ile Leu Ser	
500 505 510	
cgg ggt aat aaa gcc aga ggc ctg cag gaa cac tcc atc agt gtg gat	1762
Arg Gly Asn Lys Ala Arg Gly Leu Gln Glu His Ser Ile Ser Val Asp	
515 520 525	
ctg gat agc tcc ctg tct agt aca tta agt aac acc agc aaa gag cca	1810
Leu Asp Ser Ser Leu Ser Ser Thr Leu Ser Asn Thr Ser Lys Glu Pro	
530 535 540 545	
tct gtg tgt gaa aag gag gcc ttg ccc atc tct gag agc tcc ttt aag	1858

Ser Val Cys Glu Lys Glu Ala Leu Pro Ile Ser Glu Ser Ser Phe Lys	
550 555 560	
ctc ctc ggc tcc tcg gag gac ctg tcc agt gac tcg gag agt cat ctc	1906
Leu Leu Gly Ser Ser Glu Asp Leu Ser Ser Asp Ser Glu Ser His Leu	
565 570 575	
cca gaa gag cca gct ccg ctg tcg ccc cag cag gcc ttc agg agg cga	1954
Pro Glu Glu Pro Ala Pro Leu Ser Pro Gln Gln Ala Phe Arg Arg Arg	
580 585 590	
gca aac acc ctg agt cac ttc ccc atc gaa tgc cag gaa cct cca caa	2002
Ala Asn Thr Leu Ser His Phe Pro Ile Glu Cys Gln Glu Pro Pro Gln	
595 600 605	
cct gcc cgg ggg tcc ccg ggg gtt tcg caa agg aaa ctt atg agg tat	2050
Pro Ala Arg Gly Ser Pro Gly Val Ser Gln Arg Lys Leu Met Arg Tyr	
610 615 620 625	
cac tca gtg agc aca gag acg cct cat gaa cga aag gac ttt gaa tcc	2098
His Ser Val Ser Thr Glu Thr Pro His Glu Arg Lys Asp Phe Glu Ser	
630 635 640	
aaa gca aac cat ctt ggt gat tct ggt ggg act cct gtg aag acc cgg	2146
Lys Ala Asn His Leu Gly Asp Ser Gly Gly Thr Pro Val Lys Thr Arg	
645 650 655	
agg cat tcc tgg agg cag cag ata ttc ctc cga gta gcc acc ccg cag	2194
Arg His Ser Trp Arg Gln Gln Ile Phe Leu Arg Val Ala Thr Pro Gln	
660 665 670	
aag gcg tgc gat tct tcc agc aga tat gaa gat tat tca gag ctg gga	2242
Lys Ala Cys Asp Ser Ser Ser Arg Tyr Glu Asp Tyr Ser Glu Leu Gly	
675 680 685	
gag ctt ccc cca cga tct cct tta gaa cca gtt tgt gaa gat ggg ccc	2290
Glu Leu Pro Pro Arg Ser Pro Leu Glu Pro Val Cys Glu Asp Gly Pro	
690 695 700 705	
ttt ggc ccc cca cca gag gaa aag aaa agg aca tct cgt gag ctc cga	2338
Phe Gly Pro Pro Pro Glu Glu Lys Lys Arg Thr Ser Arg Glu Leu Arg	
710 715 720	
gag ctg tgg caa aag gct att ctt caa cag ata ctg ctg ctt aga atg	2386
Glu Leu Trp Gln Lys Ala Ile Leu Gln Gln Ile Leu Leu Leu Arg Met	
725 730 735	
gag aag gaa aat cag aag ctc caa gcc tct gaa aat gat ttg ctg aac	2434
Glu Lys Glu Asn Gln Lys Leu Gln Ala Ser Glu Asn Asp Leu Leu Asn	
740 745 750	
aag cgc ctg aag ctc gat tat gaa gaa att act ccc tgt ctt aaa gaa	2482
Lys Arg Leu Lys Leu Asp Tyr Glu Glu Ile Thr Pro Cys Leu Lys Glu	
755 760 765	
gta act aca gtg tgg gaa aag atg ctt agc act cca gga aga tca aaa	2530

Val	Thr	Thr	Val	Trp	Glu	Lys	Met	Leu	Ser	Thr	Pro	Gly	Arg	Ser	Lys	
770					775				780					785		
att	aag	ttt	gac	atg	gaa	aaa	atg	cac	tcg	gct	ggt	ggg	caa	ggg	gtg	2578
Ile	Lys	Phe	Asp	Met	Glu	Lys	Met	His	Ser	Ala	Val	Gly	Gln	Gly	Val	
				790					795					800		
cca	cgt	cat	cac	cga	ggg	gaa	atc	tgg	aaa	ttt	cta	gct	gag	caa	ttc	2626
Pro	Arg	His	His	Arg	Gly	Glu	Ile	Trp	Lys	Phe	Leu	Ala	Glu	Gln	Phe	
				805					810					815		
cac	ctt	aaa	cac	cag	ttt	ccc	agc	aaa	cag	cag	cca	aag	gat	gtg	cca	2674
His	Leu	Lys	His	Gln	Phe	Pro	Ser	Lys	Gln	Gln	Pro	Lys	Asp	Val	Pro	
				820					825					830		
tac	aaa	gaa	ctc	tta	aag	cag	ctg	act	tcc	cag	cag	cat	gcg	att	ctt	2722
Tyr	Lys	Glu	Leu	Leu	Lys	Gln	Leu	Thr	Ser	Gln	Gln	His	Ala	Ile	Leu	
				835					840					845		
att	gac	ctt	ggg	cga	acc	ttt	cct	aca	cac	cca	tac	ttc	tct	gcc	cag	2770
Ile	Asp	Leu	Gly	Arg	Thr	Phe	Pro	Thr	His	Pro	Tyr	Phe	Ser	Ala	Gln	
850					855					860				865		
ctt	gga	gca	gga	cag	cta	tcg	ctt	tac	aac	att	ttg	aag	gcc	tac	tca	2818
Leu	Gly	Ala	Gly	Gln	Leu	Ser	Leu	Tyr	Asn	Ile	Leu	Lys	Ala	Tyr	Ser	
				870					875					880		
ctt	cta	gac	cag	gaa	gtg	gga	tat	tgc	caa	ggg	ctc	agc	ttt	gta	gca	2866
Leu	Leu	Asp	Gln	Glu	Val	Gly	Tyr	Cys	Gln	Gly	Leu	Ser	Phe	Val	Ala	
				885					890					895		
ggc	att	ttg	ctt	ctt	cat	atg	agt	gag	gaa	gag	gcg	ttt	aaa	atg	ctc	2914
Gly	Ile	Leu	Leu	Leu	His	Met	Ser	Glu	Glu	Glu	Ala	Phe	Lys	Met	Leu	
				900					905					910		
aag	ttt	ctg	atg	ttt	gac	atg	ggg	ctg	cgg	aaa	cag	tat	cgg	cca	gac	2962
Lys	Phe	Leu	Met	Phe	Asp	Met	Gly	Leu	Arg	Lys	Gln	Tyr	Arg	Pro	Asp	
				915					920					925		
atg	att	att	tta	cag	atc	cag	atg	tac	cag	ctc	tcg	agg	ttg	ctt	cat	3010
Met	Ile	Ile	Leu	Gln	Ile	Gln	Met	Tyr	Gln	Leu	Ser	Arg	Leu	Leu	His	
930					935					940				945		
gat	tac	cac	aga	gac	ctc	tac	aat	cac	ctg	gag	gag	cac	gag	atc	ggc	3058
Asp	Tyr	His	Arg	Asp	Leu	Tyr	Asn	His	Leu	Glu	Glu	His	Glu	Ile	Gly	
				950					955					960		
ccc	agc	ctc	tac	gct	gcc	ccc	tgg	ttc	ctc	acc	atg	ttt	gcc	tca	cag	3106
Pro	Ser	Leu	Tyr	Ala	Ala	Pro	Trp	Phe	Leu	Thr	Met	Phe	Ala	Ser	Gln	
				965					970					975		
ttc	ccg	ctg	gga	ttc	gta	gcc	aga	gtc	ttt	gat	atg	att	ttt	ctt	cag	3154
Phe	Pro	Leu	Gly	Phe	Val	Ala	Arg	Val	Phe							

Gly Thr Glu Val Ile Phe Lys Val Ala Leu Ser Leu Leu Gly Ser His
 995 1000 1005
 aag ccc ttg att ctg cag cat gaa aac cta gaa acc ata gtt gac ttt 3250
 Lys Pro Leu Ile Leu Gln His Glu Asn Leu Glu Thr Ile Val Asp Phe
 1010 1015 1020 1025
 ata aaa agc acg cta ccc aac ctt ggc ttg gta cag atg gaa aag acc 3298
 Ile Lys Ser Thr Leu Pro Asn Leu Gly Leu Val Gln Met Glu Lys Thr
 1030 1035 1040
 atc aat cag gta ttt gaa atg gac atc gct aaa cag tta caa gct tat 3346
 Ile Asn Gln Val Phe Glu Met Asp Ile Ala Lys Gln Leu Gln Ala Tyr
 1045 1050 1055
 gaa gtt gag tac cac gtc ctt caa gaa gaa ctt atc gat tcc tct cct 3394
 Glu Val Glu Tyr His Val Leu Gln Glu Glu Leu Ile Asp Ser Ser Pro
 1060 1065 1070
 ctc agt gac aac caa aga atg gat aaa tta gag aaa acc aac agc agc 3442
 Leu Ser Asp Asn Gln Arg Met Asp Lys Leu Glu Lys Thr Asn Ser Ser
 1075 1080 1085
 tta cgc aaa cag aac ctt gac ctc ctt gaa cag ttg cag gtg gca aat 3490
 Leu Arg Lys Gln Asn Leu Asp Leu Leu Glu Gln Leu Gln Val Ala Asn
 1090 1095 1100 1105
 ggt agg atc caa agc ctt gag gcc acc att gag aag ctc ctg agc agt 3538
 Gly Arg Ile Gln Ser Leu Glu Ala Thr Ile Glu Lys Leu Leu Ser Ser
 1110 1115 1120
 gag agc aag ctg aag cag gcc atg ctt acc tta gaa ctg gag cgg tcg 3586
 Glu Ser Lys Leu Lys Gln Ala Met Leu Thr Leu Glu Leu Glu Arg Ser
 1125 1130 1135
 gcc ctg ctg cag acg gtg gag gag ctg cgg cgg cgg agc gca gag ccc 3634
 Ala Leu Leu Gln Thr Val Glu Glu Leu Arg Arg Arg Ser Ala Glu Pro
 1140 1145 1150
 agc gac cgg gag cct gag tgc acg cag ccc gag ccc acg ggc gac tga 3682
 Ser Asp Arg Glu Pro Glu Cys Thr Gln Pro Glu Pro Thr Gly Asp *
 1155 1160 1165
 cagctctgca ggagagattg caacaccatc ccacactgtc caggccttaa ctgagagggga 3742
 cagaagacgc tggaaggaga gaaggaagcg ggaagtgtgc ttctcagggga ggaaaccggc 3802
 ttgccagcaa gtagattctt acgaactcca acttgcaatt cagggggcat gtcccagtg 3862
 tttttttgtt gtttttagat actaaatcgt cccttctcca gtcttgatta ctgtacacag 3922
 tagctttaga tggcgtggac gtgaataaat gcaacttatg ttttaaaaaa aaaaaaaaaa 3982
 aaaaaa 3988

<210> 5

<211> 1168

<212> PRT

<213> Homo sapiens

<400> 5

Met Glu Pro Ile Thr Phe Thr Ala Arg Lys His Leu Leu Pro Asn Glu
1 5 10 15
Val Ser Val Asp Phe Gly Leu Gln Leu Val Gly Ser Leu Pro Val His
20 25 30
Ser Leu Thr Thr Met Pro Met Leu Pro Trp Val Val Ala Glu Val Arg
35 40 45
Arg Leu Ser Arg Gln Ser Thr Arg Lys Glu Pro Val Thr Lys Gln Val
50 55 60
Arg Leu Cys Val Ser Pro Ser Gly Leu Arg Cys Glu Pro Glu Pro Gly
65 70 75 80
Arg Ser Gln Gln Trp Asp Pro Leu Ile Tyr Ser Ser Ile Phe Glu Cys
85 90 95
Lys Pro Gln Arg Val His Lys Leu Ile His Asn Ser His Asp Pro Ser
100 105 110
Tyr Phe Ala Cys Leu Ile Lys Glu Asp Ala Val His Arg Gln Ser Ile
115 120 125
Cys Tyr Val Phe Lys Ala Asp Asp Gln Thr Lys Val Pro Glu Ile Ile
130 135 140
Ser Ser Ile Arg Gln Ala Gly Lys Ile Ala Arg Gln Glu Glu Leu His
145 150 155 160
Cys Pro Ser Glu Phe Asp Asp Thr Phe Ser Lys Lys Phe Glu Val Leu
165 170 175
Phe Cys Gly Arg Val Thr Val Ala His Lys Lys Ala Pro Pro Ala Leu
180 185 190
Ile Asp Glu Cys Ile Glu Lys Phe Asn His Val Ser Gly Ser Arg Gly
195 200 205
Ser Glu Ser Pro Arg Pro Asn Pro Pro His Ala Ala Pro Thr Gly Ser
210 215 220
Gln Glu Pro Val Arg Arg Pro Met Arg Lys Ser Phe Ser Gln Pro Gly
225 230 235 240
Leu Arg Ser Leu Ala Phe Arg Lys Glu Leu Gln Asp Gly Gly Leu Arg
245 250 255
Ser Ser Gly Phe Phe Ser Ser Phe Glu Glu Ser Asp Ile Glu Asn His
260 265 270
Leu Ile Ser Gly His Asn Ile Val Gln Pro Thr Asp Ile Glu Glu Asn
275 280 285
Arg Thr Met Leu Phe Thr Ile Gly Gln Ser Glu Val Tyr Leu Ile Ser
290 295 300
Pro Asp Thr Lys Lys Ile Ala Leu Glu Lys Asn Phe Lys Glu Ile Ser

305	310	315	320
Phe Cys Ser Gln Gly Ile Arg His Val Asp His Phe Gly Phe Ile Cys			
	325	330	335
Arg Glu Ser Ser Gly Gly Gly Gly Phe His Phe Val Cys Tyr Val Phe			
	340	345	350
Gln Cys Thr Asn Glu Ala Leu Val Asp Glu Ile Met Met Thr Leu Lys			
	355	360	365
Gln Ala Phe Thr Val Ala Ala Val Gln Gln Thr Ala Lys Ala Pro Ala			
	370	375	380
Gln Leu Cys Glu Gly Cys Pro Leu Gln Ser Leu His Lys Leu Cys Glu			
385	390	395	400
Arg Ile Glu Gly Met Asn Ser Ser Lys Thr Lys Leu Glu Leu Gln Lys			
	405	410	415
His Leu Thr Thr Leu Thr Asn Gln Glu Gln Ala Thr Ile Phe Glu Glu			
	420	425	430
Val Gln Lys Leu Arg Pro Arg Asn Glu Gln Arg Glu Asn Glu Leu Ile			
	435	440	445
Ile Ser Phe Leu Arg Cys Leu Tyr Glu Glu Lys Gln Lys Glu His Ile			
	450	455	460
His Ile Gly Glu Met Lys Gln Thr Ser Gln Met Ala Ala Glu Asn Ile			
465	470	475	480
Gly Ser Glu Leu Pro Pro Ser Ala Thr Arg Phe Arg Leu Asp Met Leu			
	485	490	495
Lys Asn Lys Ala Lys Arg Ser Leu Thr Glu Ser Leu Glu Ser Ile Leu			
	500	505	510
Ser Arg Gly Asn Lys Ala Arg Gly Leu Gln Glu His Ser Ile Ser Val			
	515	520	525
Asp Leu Asp Ser Ser Leu Ser Ser Thr Leu Ser Asn Thr Ser Lys Glu			
	530	535	540
Pro Ser Val Cys Glu Lys Glu Ala Leu Pro Ile Ser Glu Ser Ser Phe			
545	550	555	560
Lys Leu Leu Gly Ser Ser Glu Asp Leu Ser Ser Asp Ser Glu Ser His			
	565	570	575
Leu Pro Glu Glu Pro Ala Pro Leu Ser Pro Gln Gln Ala Phe Arg Arg			
	580	585	590
Arg Ala Asn Thr Leu Ser His Phe Pro Ile Glu Cys Gln Glu Pro Pro			
	595	600	605
Gln Pro Ala Arg Gly Ser Pro Gly Val Ser Gln Arg Lys Leu Met Arg			
	610	615	620
Tyr His Ser Val Ser Thr Glu Thr Pro His Glu Arg Lys Asp Phe Glu			
625	630	635	640
Ser Lys Ala Asn His Leu Gly Asp Ser Gly Gly Thr Pro Val Lys Thr			

645 650 655
Arg Arg His Ser Trp Arg Gln Gln Ile Phe Leu Arg Val Ala Thr Pro
660 665 670
Gln Lys Ala Cys Asp Ser Ser Ser Arg Tyr Glu Asp Tyr Ser Glu Leu
675 680 685
Gly Glu Leu Pro Pro Arg Ser Pro Leu Glu Pro Val Cys Glu Asp Gly
690 695 700
Pro Phe Gly Pro Pro Pro Glu Glu Lys Lys Arg Thr Ser Arg Glu Leu
705 710 715 720
Arg Glu Leu Trp Gln Lys Ala Ile Leu Gln Gln Ile Leu Leu Leu Arg
725 730 735
Met Glu Lys Glu Asn Gln Lys Leu Gln Ala Ser Glu Asn Asp Leu Leu
740 745 750
Asn Lys Arg Leu Lys Leu Asp Tyr Glu Glu Ile Thr Pro Cys Leu Lys
755 760 765
Glu Val Thr Thr Val Trp Glu Lys Met Leu Ser Thr Pro Gly Arg Ser
770 775 780
Lys Ile Lys Phe Asp Met Glu Lys Met His Ser Ala Val Gly Gln Gly
785 790 795 800
Val Pro Arg His His Arg Gly Glu Ile Trp Lys Phe Leu Ala Glu Gln
805 810 815
Phe His Leu Lys His Gln Phe Pro Ser Lys Gln Gln Pro Lys Asp Val
820 825 830
Pro Tyr Lys Glu Leu Leu Lys Gln Leu Thr Ser Gln Gln His Ala Ile
835 840 845
Leu Ile Asp Leu Gly Arg Thr Phe Pro Thr His Pro Tyr Phe Ser Ala
850 855 860
Gln Leu Gly Ala Gly Gln Leu Ser Leu Tyr Asn Ile Leu Lys Ala Tyr
865 870 875 880
Ser Leu Leu Asp Gln Glu Val Gly Tyr Cys Gln Gly Leu Ser Phe Val
885 890 895
Ala Gly Ile Leu Leu Leu His Met Ser Glu Glu Glu Ala Phe Lys Met
900 905 910
Leu Lys Phe Leu Met Phe Asp Met Gly Leu Arg Lys Gln Tyr Arg Pro
915 920 925
Asp Met Ile Ile Leu Gln Ile Gln Met Tyr Gln Leu Ser Arg Leu Leu
930 935 940
His Asp Tyr His Arg Asp Leu Tyr Asn His Leu Glu Glu His Glu Ile
945 950 955 960
Gly Pro Ser Leu Tyr Ala Ala Pro Trp Phe Leu Thr Met Phe Ala Ser
965 970 975
Gln Phe Pro Leu Gly Phe Val Ala Arg Val Phe Asp Met Ile Phe Leu

980 985 990
Gln Gly Thr Glu Val Ile Phe Lys Val Ala Leu Ser Leu Leu Gly Ser
995 1000 1005
His Lys Pro Leu Ile Leu Gln His Glu Asn Leu Glu Thr Ile Val Asp
1010 1015 1020
Phe Ile Lys Ser Thr Leu Pro Asn Leu Gly Leu Val Gln Met Glu Lys
1025 1030 1035 1040
Thr Ile Asn Gln Val Phe Glu Met Asp Ile Ala Lys Gln Leu Gln Ala
1045 1050 1055
Tyr Glu Val Glu Tyr His Val Leu Gln Glu Glu Leu Ile Asp Ser Ser
1060 1065 1070
Pro Leu Ser Asp Asn Gln Arg Met Asp Lys Leu Glu Lys Thr Asn Ser
1075 1080 1085
Ser Leu Arg Lys Gln Asn Leu Asp Leu Leu Glu Gln Leu Gln Val Ala
1090 1095 1100
Asn Gly Arg Ile Gln Ser Leu Glu Ala Thr Ile Glu Lys Leu Leu Ser
1105 1110 1115 1120
Ser Glu Ser Lys Leu Lys Gln Ala Met Leu Thr Leu Glu Leu Glu Arg
1125 1130 1135
Ser Ala Leu Leu Gln Thr Val Glu Glu Leu Arg Arg Arg Ser Ala Glu
1140 1145 1150
Pro Ser Asp Arg Glu Pro Glu Cys Thr Gln Pro Glu Pro Thr Gly Asp
1155 1160 1165

<210> 6

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<221> misc_binding

<222> 1..18

<223> sequencing oligonucleotide PrimerPU

<400> 6

tgtaaaacga cggccagt

18

<210> 7

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<221> misc_binding

<222> 1..18

<223> sequencing oligonucleotide PrimerRP

<400> 7

caggaaacag ctatgacc

18